Letters and comments

Contributions to this section are asked to make their comments brief and to the point. Letters should comply with the notice printed on the inside back cover. Tables and figures should only be included if absolutely essential and no more than five references should be given. The Editor reserves the right to shorten letters and subedit contributions to ensure clarity.

LETTER TO THE EDITOR

doi 10.1308/147870805X42572

Impact of junior doctor education on drug allergy documentation

AN MORRITT¹, DJ ALEXANDER²

¹Department of Plastic & Reconstructive Surgery, St James's University Hospital, Leeds, UK

²Department of General Surgery, York District Hospital, York, UK

CORRESPONDENCE TO:

Mr AN Morritt, Senior House Officer, Department of Plastic & Reconstructive Surgery, St James's University Hospital, Leeds LS9 7TF, UK. E: andrewmorritt@hotmail.com

Inaccurate documentation of drug allergies can result in patients being given drugs to which they are allergic.¹ This presents serious risk of harm to patients, as well as serious medicolegal implications. Prescribing errors account for up to 25% of litigation claims in medical practice.² The UK Department of Health has aimed to reduce by 40% the number of serious errors in the use of prescribed drugs by the end of 2005.²

We performed a snapshot audit of current practice in drug allergy documentation. Consultant approval was obtained though none of the junior doctors concerned had prior knowledge of the audit. The study group comprised of all general surgery (colorectal, breast and vascular surgery) and urology in-patients. Information about drug allergies was collected from: the patient; the clerking for the current admission; and the drug allergies' box on the drug chart. Patients with drug allergies were checked for the presence of a drug allergies' wristband. Drug allergy documentation in the medical notes and on the drug charts was recorded as correct or incorrect

depending on whether or not the information correlated with what the patient said.

A letter was sent by a consultant surgeon to all PRHOs and SHOs in general surgery and urology (12 doctors). It summarised the results of the first audit and stated: (i) it was their responsibility to document patients drug allergies in the medical notes and on the drug chart; (ii) that it was insufficient to leave drug allergies boxes blank on the premise that in a court of law 'if it isn't documented it hasn't been asked'; and (iii), finally, that they must check the allergies' box on the drug chart before prescribing any drugs. The letter also warned that there would be a re-audit in the future; this was performed 3 months after sending the letter.

The first audit (92 patients) showed that 20% of notes and 24% of drug charts contained inaccurate drug allergy documentation. Twenty-three patients (25%) had one or more drug allergies (most commonly to penicillin); 20/23 patients (87%) with a drug allergy were wearing a drug allergy bracelet. The second audit (89 patients) showed that the documentation of drug allergies was accurate in all notes and drug charts. Twenty-one (24%) had one or more drug allergies (most commonly to penicillin); 18/21 (86%) of patients with drug allergies were wearing a drug allergy wrist bracelet.

The first audit showed that documentation of drug allergies in the medical notes and on the drug charts for general surgery and urology inpatients was imperfect. The considerable improvement in the documentation was due to the education of junior doctors about their roles and responsibilities, and also due to the threat of re-audit. There was little difference between documentation of the patient's drug allergies on the drug allergy wristband between the first and second audits. This is probably because documentation of drug allergies on the patient's wristband is performed by the nursing staff who did not receive the letter sent to the junior doctors. Reducing errors in the documentation of patient's drug allergies should reduce the chance of patients being given medications to which they are allergic. This, in turn, will reduce the number of adverse drug reactions suffered. As a result of this audit, junior doctors will receive a letter stating their responsibilities to document drug allergies accurately before they start work in the department.

In summary, documentation of drug allergies in the medical notes and on drug charts on the general surgery and urology wards at a district general hospital was imperfect, but was improved by education of junior doctors and the prospect of re-audit.

References

1. McDonnell PJ, Jacobs MR. Hospital admissions resulting from preventable adverse drug reactions. *Ann Pharmacother* 2002; **36**: 1331–6.
2. Wright J, Hill P. *Clinical Governance*. London: Churchill Livingstone, 2003.

COMMENTS ON:

Deliss L. Basic surgical training tips. *Ann R Coll Surg Engl* 2003; **85**: 417–9

Deliss L. Designing a basic surgical training timetable. *Ann R Coll Surg Engl* 2003; **85**: 420–1

doi 10.1308/147870805X42518

Implementing a 'New Deal' compliant SHO rota

AW LAMBERT

Department of General Surgery, Derriford Hospital, Plymouth, UK

CORRESPONDENCE TO:

Mr AW Lambert, Consultant General & Vascular Surgeon, Department of General Surgery, Derriford Hospital, Plymouth PL6 8DH, UK
E: anthony.lambert@phnt.swest.nhs.uk

Having struggled to implement a 'New Deal' compliant SHO rota locally and only just having managed to do so with severe implications for training, I read with interest these thoughts and suggestions for SHO training. As they stand, the suggestions of Deliss do not achieve compliance with the New Deal. The total number of hours each week would be 61, including a half-day off after being on-call (56 allowed). Minimum rest periods must be achieved overnight (a minimum of 5 hours undisturbed rest between 22:00 and 07:00 on 70% of occasions). The SHO normal working day has also been shortened to 08:30–16:30.

In terms of training opportunities, the 'learning experiences' are a maximum of 2.5 hours long, meaning the SHO would have to arrive late or leave early, be it a theatre session or out-patient clinic, so losing valuable experience.

The European Working Time Directive^{1,2} will eventual mean that no junior doctors can work more than 48 hours a week, with all time spent in the hospital being counted as work. This will lead to the implementation of full-shift rotas, as has already occurred in our hospital. We believe that these constraints are not in the best interests of patients, consultants or trainees. However, they are unavoidable and we should develop timetables, working practices and educational programmes that will satisfy both hours' requirements and training needs the better.

References

1. Council Directive 93/104/EC of the European Parliament and Council. *Official Journal of the European Community* 2000; **L307**: 18–24.
2. Council Directive 2000/34/EC of the European Parliament and Council. *Official Journal of the European Community*. 2000; **L195**: 41–5.

doi 10.1308/147870805X42527

Timetable design

D LOVEDAY

Department of Orthopaedics, Arrowe Park Hospital, Wirral, UK

CORRESPONDENCE TO:

Dr D Loveday, Department of Orthopaedics, Arrowe Park Hospital, Wirral CH49 5PE, UK
E: davidloveday@hotmail.com

I read this article with great interest as I am a senior house officer (SHO) on a basic surgical training rotation. The 'best practice' timetable fails to address the main problem with devising SHO timetables. I am amazed it includes an 'on-call' day of 24 hours. With the European Working Time Directive (EWTd) coming further into force in August 2004, very few SHOs are able to work an on-call day. SHO working patterns include full shift, partial shift and on-call rotas, all with minimum rest periods. This is the problem for SHO timetables and planning an ideal timetable for SHOs needs to address the EWTd.

AUTHOR'S RESPONSE

doi 10.1308/147870805X42536

LOUIS DELISS

Suffolk Nuffield Hospital, Suffolk, UK

CORRESPONDENCE TO:

Louis Deliss, Orthopaedic, Upper Limb and Hand Surgeon, Playford Mount, Great Bealings, Woodbridge, Suffolk IP13 6PE, UK
Tel: +44 (0)1473 735600; Fax: +44 (0)1473 735106; E-mail: louisjdeliss@aol.com

I thank the authors for their interest and would encourage them to keep trying to maintain teaching standards within the new rules. My suggested timetable is not perfect but is a template that needs adapting for local use and staffing patterns. 'Learning opportunities' are designed as parts of clinical sessions specifically organised for teaching. This leaves some time for 'service workload' commitments. I do not believe that 'full-shift' working is inevitable or desirable. It would appear that we are still not certain of the exact details of the EWTd rules as applied to doctors in training.

I am pleased that my paper has started a lively debate, and would urge others to publish their solutions (possibly on the College web-site). The publications on the Government web-sites are disappointing and mostly unhelpful.

COMMENT ON:

Spencer J, Smith A, Woods D. The effect of time delay on infection in open long-bone fractures: a 5-year prospective audit from a district general hospital. *Ann R Coll Surg Engl* 2004; **86**: 108–12

doi 10.1308/147870805X42581

A significant bias in the results?

JR GIBBS, D RICKETTS

Orthopaedic Department, The Princess Royal Hospital, Haywards Heath, West Sussex, UK

CORRESPONDENCE TO:

Mr JR Gibbs, Secretary, Journals Club, Orthopaedic Department, The Princess Royal Hospital, Lewes Road, Haywards Heath, West Sussex RH16 4EX, UK

We read this article with interest and would like to raise the following points. First, 27 patients out of 130 were not included in the final analysis, giving an exclusion rate of 20%. Amongst this group of 27 were patients who had sustained multiple injuries, a severe head injury or died from their injuries. Excluding this group of patients may have influenced the final results. We feel it would be beneficial to detail the fractures sustained in this group along with the Gustilo and Anderson grade of the fracture.

Second, as the authors state, there is no randomisation regarding the timing of operation. The table of infected cases seems to illustrate that wound debridement, lavage and fracture stabilisation took place earlier for the more severe grades of fracture. This, we feel, would bias the final results.

In light of the above, we believe this paper does not have a sufficient evidence base to change the current management of open fractures.

AUTHOR'S RESPONSE

doi 10.1308/147870805X42563

JONATHAN SPENCER

Nuffield Orthopaedic Centre, Oxford, UK

CORRESPONDENCE TO:

Mr Jonathan Spencer, NDOS, Nuffield Orthopaedic Centre, Windmill Road, Oxford OX3 7LD, UK
Tel: +44 (0)1865 227645; E-mail: Jonmfs Spencer@aol.com

We would agree with the authors in their observations. However, this is a very difficult cohort of patients to study, as there are many confounding factors at all levels of the study that may lead to the introduction of bias into the results. This point was explored in the discussion.

We initially attempted to follow-up patients who received primary treatment in other hospitals. We found that records were very difficult to obtain and the quality of the information to be of such poor quality as to be misleading. We, therefore, felt that data on these patients should not be included in the analysis.

Randomisation of the timing of surgery of these patients would have been impractical and unethical. It was, therefore, not done.

We also believe this paper does not offer, on its own, sufficient evidence to warrant a change in the current management of open fractures. We also acknowledge that the hypothesis that 'a delay to surgical debridement of open fractures increases the infection' is very difficult to prove statistically due to multiple confounding factors. Further work in this area is needed.

We do, however, propose that experienced teams should treat these injuries in normal working hours if at all possible in order to optimise the outcome for the patient.

COMMENT ON:

Ray S, Mackie C. Positioning the patient for abdominoperineal excision of the rectum (APER). *Ann R Coll Surg Engl* 2003; **85**: 281

doi 10.1308/147870805X42509

High risk of lower limb neuropraxia and litigations for a better view of the perineum

HAK-MIEN QUAH, AJAI KUMAR ABDUL SAMAD, DAVID J HAY, ANDREW MAW

Department of General Surgery, Glan Clwyd Hospital, Rhyl, UK

CORRESPONDENCE TO:

Mr Andrew Maw, Department of General Surgery, Glan Clwyd Hospital, Rhyl, Denbighshire LL18 5UJ, UK
Tel: +44 (0)1745 534957; Fax: +44 (0)1745 534561; E-mail: Andrew.Maw@cd-tr.wales.nhs.uk

We agree with Ray and Mackie that it is important to optimise the position of the patient for such procedure and to minimise the time in which the patient's legs are in the Lloyd-Davies position. However, we are strongly against putting the legs on the inside of padded lithotomy poles as there are risks of direct compression on the peroneal nerve and its branches in the lateral calf. Although the authors have commented that they have seen no cases of lower limb neuropraxia or compartment syndrome, we ourselves are aware of such cases using the lithotomy and Lloyd-Davies position. Although the particular position that they describe may not place the nerves in close proximity to the lithotomy poles, patients may alter their position during the procedure, and some patient's body habitus may also directly place the nerves at risk.

We have recently been involved in a case where if a nerve injury occurred with the use of the lithotomy poles, it would be deemed to be negligent by default. We would instead recommend the use of the modified 'powered' Alan Boots (Yellowfins, Amatech, Massachusetts, USA), which have inherent protective structures for the nerves, provide a decent exposure to the perineum and can be adjusted easily during the procedure.

COMMENT ON:

Sotheran WJ, Rainsbury RM. Skin-sparing mastectomy in the UK – a review of current practice. *Ann R Coll Surg Engl* 2004; **86**: 82–6

doi 10.1308/147870805X42545

Reconstruction after skin sparing mastectomy

E ARUMUGAM, J PEREIRA, PM GILBERT

Department of Plastic Surgery, Queen Victoria Hospital, East Grinstead, West Sussex, UK

CORRESPONDENCE TO:

Mr PM Gilbert, Consultant Plastic and Reconstructive Surgeon, Queen Victoria Hospital, East Grinstead, West Sussex RH19 3DZ, UK

The authors describe the use of skin sparing mastectomy in the UK and subsequent reconstruction stating that the TRAM flap has been established as the 'gold standard' for breast reconstruction.¹ Surgery has, however, not stood still in the last few years. The TRAM flap was first described by Hartrampf *et al.* in 1982; however, there were problems with the vascularity of this flap.

The anatomical studies into the blood supply of the anterior abdominal wall skin by Taylor *et al.* indicated the basis for these problems. Amez *et al.* described the use of the free TRAM flap in 1991. Blood supply to the flap tissue was improved but of course the technique required experience in microvascular surgery. A further advance of the free TRAM flap was the fact that only small portions of the rectus muscle were excised thereby reducing the chances of later abdominal wall herniation. It was almost inevitable that the flap would be developed to a point where no muscle at all needed sacrifice.

The deep inferior epigastric perforator (DIEP) flap was originally described by Holmstrom² as long ago as 1979 but was not taken up by other surgeons until it was popularised by Allen and Treece in 1994 and also by Blondeel in the same year. Blondeel's report of 100 DIEP flaps in 1999 and subsequent report from the same unit of more than 500 cases has established it as the new 'gold standard'. What are the implications of this? To achieve the best possible results after various types of mastectomy now requires the use of techniques employing microvascular reconstruction. At

present, only plastic surgeons have been trained in these reconstructions; however, there is no reason why breast surgeons should not be trained to do it also. As an illustration, a few years ago most head and neck microvascular reconstruction was being done by plastic surgeons assisting maxillofacial surgeons. The latter speciality has very successfully incorporated microvascular surgery into its armamentarium. If breast surgeons do not do the same, then either they will do inadequate reconstructions or they will have to rely on the availability of plastic surgeons with all the logistical problems that this involves.

References

1. Malata CM, McIntosh SA, Purushotham AD. Immediate breast reconstruction after mastectomy for cancer. *Br J Surg* 2000; **87**: 1455–72.
2. Holmstrom H. The free abdominoplasty flap and its use in breast reconstruction. *Scand J Plast Reconstr Surg* 1979; **13**: 423–7.

AUTHORS' RESPONSE

doi 10.1308/147870805X42554

WJ SOTHERAN¹, RM RAINSBURY²

¹Portsmouth Hospitals Trust, Hampshire, UK

²Royal Hampshire County Hospital, Winchester, UK

CORRESPONDENCE TO:

Wendy Sotheran, 14 West Street, Titchfield, Hampshire PO14 4DH, UK. E: wsotheran@aol.com

We quoted from Malata *et al.* that the TRAM flap had become the 'gold standard' for breast reconstruction and commented that limitations to training were likely to account for the infrequency with which it was offered by our respondents.

Arumugam and colleagues argue that the DIEP flap should be considered the new 'gold standard' and we agree that this procedure can lead to excellent results. Training in microvascular surgery would be welcomed by many surgeons offering breast reconstruction and is likely to become integrated into oncoplastic training programmes in the future. However, to give the 'best possible results' in reconstruction, the extended operating time and the significant risk of total flap loss must be weighed against the benefits of a preserved rectus. The DIEP flap is an excellent procedure in centres with a high caseload and sufficient theatre capacity. For many surgeons working outside these centres, the pedicled TRAM or latissimus dorsi reconstruction provide a safe and acceptable alternative with high patient satisfaction and low morbidity.

To describe all these cases as 'inadequate reconstructions' implies that the technical excellence of a procedure is paramount, regardless of the patients' experience or expectations, or what the NHS can afford to provide.