Letters and comments

Contributions to this section are asked to make their comments brief and to the point. Letters should comply with the notice printed on the inside back cover. Tables and figures should only be included if absolutely essential and no more than five references should be given. The Editor reserves the right to shorten letters and subedit contributions to ensure clarity.

COMMENTS ON:

M Bailey, A Kingsnorth. Controversial Topics in Surgery: Inguinal hernia – laparoscopic versus open repair. *Ann R Coll Surg Engl* 2005; **87**: 57–60

RESPONSE 1

doi 10.1308/003588405X50985

Radical retropubic prostatectomy and inguinal hernia repair

NIKESH THIRUCHELVAM, ANDREW ADAMSON

Department of Urology, Royal Hampshire County Hospital, Winchester, UK

CORRESPONDENCE TO

Nikesh Thiruchelvam, Department of Urology, Royal Hampshire County Hospital, Romsey Road, Winchester SO22 6DG, UK

T: +44 (0)1962 8663535; E: n.t@talk21.com

We read with interest the recent *Controversial Topics in Surgery* debate regarding inguinal hernia – laparoscopic versus open repair.^{1,2} The benefits of both methods were eloquently presented; however, as urologists, we feel that an important consideration has been ignored in this debate.

Increasing numbers of younger men are now being diagnosed with early prostate cancer. Though optimal management of this patient group has yet to be established, it is likely that many men will continue to opt for ablative surgery of the prostate. Our personal experience of attempted radical retropubic prostatectomy being near impossible following laparoscopic hernia repair is confirmed by others who substantiate it to be incredibly difficult, if not impossible.^{3,4} Regardless of whether the laparoscopic repair is via the transperitoneal or

extraperitoneal approach, dissection of the obliterated retro pubic space, to allow mobilisation of the prostate is difficult because of the secondary fibrotic reaction caused by the mesh repair. Fibrosis around the anterior surface of the bladder and the prostate, often extending laterally to the endopelvic fascia, obliterates the tissue planes compromising oncological cancer principles and, most probably, outcomes. The same problems will face the laparoscopic radical prostatectomist and also those who favour the perineal approach.

We feel that general surgeons who regularly perform laparoscopic inguinal hernia repairs should be aware of the consequences of this approach, as decision making in patients who are subsequently diagnosed with early prostate cancer will be influenced by their actions. Indeed, perhaps there is a case for more careful prostate assessment or ‘screening’ of patients undergoing laparoscopic hernia repair.

References

1. Bailey M. Inguinal hernia – laparoscopic or open repair? The case for laparoscopic repair. *Ann R Coll Surg Engl* 2005; **87**: 57–8.
2. Kingsnorth A. Inguinal hernia – laparoscopic or open repair? The case for open repair. *Ann R Coll Surg Engl* 2005; **87**: 59–60.
3. Cook H, Afzal N, Cornaby AJ. Laparoscopic hernia repairs may make subsequent radical retropubic prostatectomy more hazardous. *BJU Int* 2003; **91**: 729.
4. Katz EE, Patel RV, Sokoloff MH, Vargish T, Brendler CB. Bilateral laparoscopic inguinal hernia repair can complicate subsequent radical retropubic prostatectomy. *J Urol* 2002; **167**: 637–8.

RESPONSE 2

doi 10.1308/003588405X50994

Single layer Shouldice repair with mesh inlay and onlay

D. HEGDE

Department of Surgery, A J Institute of Health Sciences, Mangalore, India

CORRESPONDENCE TO

D Hegde, Dean and Professor of Surgery, A J Institute of Health Sciences, Mangalore 575004, India

T: +91 (0)824 9343562024; E: drdhegde@hotmail.com

There can be little argument that there is a need for consensus and common purpose, as we approach the daily

problem of inguinal herniae. While a laparoscopist will vehemently argue about the virtues of laparoscopic method (without waiting for the long-term results), most others may argue for the open method, because it is time-tested, and avoids the risks and uncertainties of a totally different method. While in olden days we accepted the inevitability of pain after an operation, today we are sensitising our patients to even minor pain, in an effort to recruit them for some of the minimally invasive procedures. Reduced pain after such procedures does not make the patient immune to another pain, for instance a tooth ache. Pain is part of human existence and we should alleviate it wherever possible. It is not in the long-term interest of the patient and the doctor to sensitise patients to an emotive issue like pain.

I am convinced that the open method of hernia surgery will remain the popular method, just as newer technologies have not replaced the good old razor. Just as we have better razor blades, now we have to improve on the open method and achieve nil recurrence.

The three major techniques, now in practice – the Shouldice method, the Lichenstein repair and the laparoscopic technique – offer good results in specialist centres. An average surgeon uniformly gets inferior results, whatever technique he practices; in fact, even specialist centres do get recurrences.

For the purist, who believes that hernia is an anatomical defect, the Shouldice repair offers the most ideal method of correcting inguinal anatomy. This precise but invasive technique is difficult to practice and leaves an ugly scar. In these days of minimal invasion, it is fascinating to watch patients queuing up to undergo surgery in a Shouldice Clinic. What a patient wants after all is a guaranteed repair and he has no time for the experimenters with promises. I have received mail from Dr Degani, chief surgeon, during this month confirming that his Shouldice Clinic has a longer waiting list, than ever before.

I am convinced that the Shouldice method is the gold standard for inguinal hernia repair. However, it is impossible for lesser surgeons to get brilliant results with this technique, which is very demanding and meticulous. The main secret is the use of stainless steel wire instead of Prolene. It is a pity that surgeons are generally uncomfortable with wire, which is stronger, has reduced tendency for tightening and tissue necrosis, and the knot, though difficult for a beginner, is unlikely to slip. Tight sutures and slipped knots following use of nylon probably account for most of our failures.

I started doing Shouldice repair in the early 1980s. Having now observed the technique in the Shouldice Clinic, my dissection was not so meticulous and my repair using Prolene was of inferior quality. Within 10 years of practicing it, I was frustrated by a recurrence rate far higher than in the Shouldice Clinic. While I was convinced of the soundness of Shouldice dissection, I felt mere tissue was not adequate. In

the early 1990s, I started placing a mesh inlay to reinforce the preperitoneal layer. Following a standard Shouldice dissection, I placed a mesh in the preperitoneal space to cover the peritoneal bulge. It was fixed to available tissues posteriorly. A two-layer (double breasting) repair of transversalis fascia was now done, covering the mesh. This appeared to be the ultimate repair as it included: (i) the goodness of Shouldice dissection; (ii) it avoided a difficult four-layer repair; (iii) a mesh in the preperitoneal space never gets infected. It was felt that with the additional strength of mesh, we now have a technique superior to Shouldice with the prospect of no recurrence. Now, I had the confidence to do bilateral direct hernia surgery, in one sitting (not done in the Shouldice clinic even today).

My euphoria was short-lived and I soon started having a rare recurrence medially. It was obvious that my two-layer tissue repair using Prolene and my fixation of mesh medially had their imperfections. By now, I was willing to be convinced by the Lichenstein experience – that only an onlay mesh can be stitched beyond pubic tubercle medially, and can be split laterally to enclose the cord, putting a final lid on possible areas of recurrence.

My ultimate and refined technique of a guaranteed repair of posterior wall after a standard Shouldice dissection, that produces a long upper leaf of transversalis fascia and a shorter lower leaf, the peritoneal bulge with extraperitoneal fat is separated all round posteriorly to create a space for placement of a piece of mesh, which now runs beyond what eye can see. The visible part of the mesh is sutured to back of structures – conjoined tendon above, rectus sheath medially, lower leaf of transversalis below, making sure that the mesh overhangs beyond the stitches from inside. A mere approximation of the upper leaf of transversalis to inguinal ligament covers the inlay mesh. A single layer, Prolene mesh is now sutured as onlay beyond pubic tubercle medially and split to enclose, cord laterally (as in the Lichenstein technique).

What is described above can be called 'single layer Shouldice repair with mesh inlay and onlay'. It corrects the laxity of posterior wall and strengthens it both posteriorly and anteriorly. It takes about 30 min to do. Mere placement of mesh preperitoneally (as done in the laparoscopic method) or as onlay (as in the Lichenstein method), without correcting the laxity of posterior wall, is theoretically unsound and should result in recurrence. Merely relying on tissue repair, however meticulous, as in the Shouldice method, is difficult to justify. Finally, we should take into account a reality, well known to surgeons, that a technique, meticulously executed by a master, always gives a good result. We need to offer a technique that can be practiced universally.

I have had no recurrence, or infection in 200 cases that I did in the last 2 years. A delayed infection is possible, in which event the onlay can be removed – still leaving the inlay, which in my experience never gets infected. For the same reason, I

believe strongly, in separating the onlay from the inlay. The 'hernia systems' now available, with a connecting bridge, require removal of the entire piece.

A trainee requires to do 25–50 cases under supervision to get the hang of the technique. But at the end of that period would have learnt the ultimate technique with no prospect of recurrence. Merely stitching a mesh anteriorly, though easy, cannot give the same satisfaction and result. There is also the danger of our trainees totally losing out on the experience of an inguinal dissection – all surgeons will gratefully remember how an inguinal dissection contributed to their eventual mastery of surgery.

AUTHOR'S RESPONSE

doi 10.1308/003588405X51001

MICHAEL BAILEY

Department of Gastrointestinal, Hepatobiliary & Minimal Access Surgery, Royal Surrey County Hospital, Guildford, UK

CORRESPONDENCE TO

Michael Bailey, Professor of Surgery and Director Minimal Access Therapy Training, Department of Gastrointestinal, Hepatobiliary & Minimal Access Surgery, Royal Surrey County Hospital, Egerton Road, Guildford, Surrey GU2 7XX, UK
T: +44 (0)1483 555838; F: +44 (0)1483 555836; E: mebailey@uk-consultants.co.uk

Thiruchelvam and Adamson refer to the possible difficulty of radical retropubic prostatectomy after laparoscopic hernia repair. The quoted paper by Cook *et al.*¹ is a single patient. The paper by Katz *et al.*² also refers to a single patient. A review of the literature reveals two more substantive papers. Erdogru *et al.*³ have a much larger group of 20 patients which was compared with a group of 20 matched patients. Their conclusion was that laparoscopic herniorrhaphy does not adversely affect the operative outcomes or functional results of radical prostatectomy. Stolzenburg *et al.*⁴ in a group of 9 patients, concluded that minimally invasive hernia repair with mesh placement was not a contra-indication to endoscopic extraperitoneal radical prostatectomy.

Hegde's letter relates mainly to his own modifications of established open techniques. I agree that whatever operation is performed needs to be done well. However, I disagree with him that postoperative pain is unimportant and I am sure that we should strive to offer patients procedures which minimise postoperative pain and provide a rapid return to full activity. Laparoscopic hernia repair achieves both of these objectives.

In terms of length of follow-up, it should be remembered that in Lichtenstein's⁵ paper follow-up was between 2–14

years, a very similar length of follow-up to my own laparoscopic series.

References

1. Cook H, Afzal N, Cornaby AJ. Laparoscopic hernia repairs may make subsequent radical retropubic prostatectomy more hazardous. *BJU Int* 2003; **91**: 729.
2. Katz EE, Patel RV, Sokoloff MH, Vargish T, Brendler CB. Bilateral laparoscopic inguinal hernia repair can complicate subsequent radical retropubic prostatectomy. *J Urol* 2002; **167**: 637–8.
3. Erdogru T, Teber D, Frede T, Marrero R, Hammady A, Rassweiler J. The effect of previous transperitoneal laparoscopic inguinal herniorrhaphy on transperitoneal laparoscopic radical prostatectomy. *J Urol* 2005; **173**: 769–72.
4. Stolzenburg JU, Ho KM, Do M, Rabenalt R, Dorschner W, Truss MC. Impact of previous surgery on endoscopic extraperitoneal radical prostatectomy. *Urology* 2005; **65**: 325–31.
5. Lichtenstein IL. Herniorrhaphy: a personal experience with 6,321 cases. *Am J Surg* 1987; **153**: 553–9.

COMMENT ON:

CJ Webb. Removal of ruptured cysts, *Ann R Coll Surg Engl* 2004; **86**: 384

doi 10.1308/003588405X51010

NS AWSARE, GB BOUSTEAD

Department of Urology, Lister Hospital, Stevenage, Herts, UK

CORRESPONDENCE TO

NS Awsare, Clinical Research Fellow, Department of Urology, University Hospital of Wales, Cardiff CF4 4XW, UK
M: +44 (0)7799 008021; E: ninaad_awsare@hotmail.com

We read this *Technical Tip* with interest and employ a similar technique using a balloon catheter (such as a Fogarty or Foley, depending on the size of cyst). The technique initially described for vesicovaginal fistula repair,¹ and excision of urethral diverticuli,² is equally suitable for controlled removal of cysts. The catheter is passed into the cyst through a small opening made in its wall between two stay sutures or a purse-string suture. The balloon is then inflated and sutures are tied to the catheter for security. This method has the added advantage of allowing manipulation and mild traction to aid dissection further. We find the method exceptionally useful in excision of urethral diverticuli and circumferential dissection of fistulous tracts.

References

1. Dmochowski R. Surgery for vesicovaginal fistula, urethrovaginal fistula, and urethral diverticulum. In: Walsh PC, Retik AB, Vaughan ED, Wein AJ. (eds) *Campbell's Urology*, 8th edn. Philadelphia, PA: Saunders, 2002; 1200.
2. Moore TD. Diverticulum of the female urethra. An improved technique of surgical excision. *J Urol* 1952; **68**: 611–6.

COMMENT ON:

H Ismail *et al.* A different frontal sinus drain. *Ann R Coll Surg Engl* 2004; **86**: 306

doi 10.1308/003588405X51029

A CHARKIN¹, J ROWE-JONES¹, E KERSHAW²

¹ENT Department and ²Department of Anaesthetics, Royal Surrey County Hospital, Guildford, UK

CORRESPONDENCE TO

A Charkin, ENT Department, Royal Surrey County Hospital, Egerton Road, Guildford GU2 7XX, UK
M: +44 (0)7747 792518; E: alexcharkin@hotmail.com

We read this *Technical Tip* with interest. An 8-year-old boy presented to us with a Potts Puffy tumour. CT scan demonstrated frontal sinusitis with osteomyelitis of the anterior table and a subperiosteal collection. He underwent combined approach of endoscopic fronto-ethmoidal recess dissection and open drainage using a spectacle incision. The small size of the right frontal sinus, 13 mm x 3 mm on the CT scan, prevented the use of standard drains which were all too large. An epidural catheter, 16-G, clear, three lateral eyes (Portex Ltd, Hythe, Kent CT21 6SL, UK) with its luer lock proved ideal. This was sutured to the skin and provided an effective and simple way to irrigate the sinus.

COMMENT ON:

R Persaud *et al.* Life-threatening necrotizing fasciitis of the neck: an unusual consequence of a sore throat. *Ann R Coll Surg Engl* 2004; **86**: 253–7

doi 10.1308/003588405X51038

JP GARNER, NJ EVERITT

Department of General Surgery, Chesterfield and North Derbyshire Royal Hospital, Chesterfield, UK

CORRESPONDENCE TO

NJ Everitt, Department of General Surgery, Chesterfield and North Derbyshire Royal Hospital, Calow, Chesterfield S44 5BL, UK
E: jeffgarner@doctors.org.uk

We are relieved that Persaud *et al.*'s patient recovered from her severe streptococcal illness, but would like to express concerns regarding their report. We do not believe that, on the evidence presented, their diagnosis of necrotizing fasciitis is substantiated and that to advocate direct closure of the skin in the presence of substantial, deep-seated infection is both inappropriate and potentially dangerous.

No bacterial species, other than a streptococcus, was isolated, making synergistic necrotizing fasciitis (Type I) unlikely. Type II necrotizing fasciitis, traditionally called streptococcal gangrene is a well-defined entity with several typical features, notably early and rapid progression of skin involvement with formation of an overlying eschar and marked systemic toxicity.¹ Yet, as the authors admit, none of these features were present. Bisno and Stevens² widely accepted definition of necrotizing fasciitis describes infection that results in the progressive destruction of fascia and fat in the subcutaneous plane, but none of these features was reported at the first operation; indeed, there is no mention of necrosis at all. We doubt, therefore, that the case reported was necrotizing fasciitis of either type.

Rather, we believe that the authors describe a severe streptococcal pharyngitis with involvement of the lateral pharyngeal compartment³ and consequent inflammation of the overlying skin. Given the degree of involvement of the neck muscles, it is not surprising that tenderness exceeded the severity of cutaneous cellulitis, a feature common to many deep-seated infections.

Regardless of the precise diagnosis, severe soft tissue sepsis was present and we are surprised that the skin was closed primarily at the initial operation. Our opinion is supported by the need for debridement of necrotic tissue 6 days later following discharge of pus from the wound. Whether muscle necrosis was due to streptococcal necrotizing myositis⁴ or a compartment-type syndrome after primary closure is unclear. It seems likely to us that soft tissue destruction could have been minimised with an acceptable cosmetic result had delayed primary closure been adopted.

The mainstay of treatment for necrotizing fasciitis is radical debridement⁴ with further planned examination and debridement as required.⁵ To our knowledge, no authority recommends primary closure. We are concerned that, through incorrect classification, this report might persuade the inexperienced to attempt primary closure when necrotizing fasciitis is actually present with catastrophic results.

References

1. O'Rourke I, Carson P, Lum G. Abscess, cellulitis and necrotizing bacterial infection. In: Morris PJ, Wood WC. (eds) *Oxford Textbook of Surgery*, 2nd edn. Oxford: Oxford University Press, 2000.
2. Bisno AL, Stevens DL. Streptococcal infections of skin and soft tissues. *N Engl J Med* 1996; **354**: 240–5.
3. Ballenger JJ. Infection of the fascial spaces of the neck and floor of the mouth. In: Ballenger JJ. (ed) *Diseases of the Nose, Throat, Ear, Head and Neck*, 14th edn. Philadelphia, PA: Lea and Febiger, 1991.
4. Burge TS, Watson JD. Necrotizing fasciitis: be bloody, bold and resolute. *BMJ* 1994; **308**: 1455–4.
5. Green RJ, Dafoe DC, Raffin TA. Necrotizing fasciitis. *Chest* 1996; **110**: 219–29.

AUTHORS' RESPONSE

doi 10.1308/003588405X51047

R PERSAUD, D KRAHÉ, C GEORGALAS, S KANIYUR, S DAS, G ALUSI

St Bartholomew's Hospital, West Smithfield, London EC1A 7BE, UK

CORRESPONDENCE TO

R Persaud E: randm.persaud@ukgateway.net

We strongly believe that the evidence presented was more compatible with necrotizing fasciitis than with the diagnosis of a severe streptococcal pharyngitis, involving the lateral pharyngeal compartment and overlying skin. The latter diagnosis would not explain the extensive necrosis present along the fascial planes and adjacent muscles, seen on CT scanning and twice intra-operatively.

The general surgeons highlighted that Type II necrotizing fasciitis, traditionally called streptococcal gangrene, is a well-defined entity with several typical features, notably early and rapid progression of skin involvement with formation of an overlying eschar and marked systemic toxicity. The condition is characterised by rapid progression of necrosis along the fascial planes; skin necrosis, although common, is not an absolute defining feature.¹ The absence of the general features, and more specifically skin necrosis at the initial debridement, does not preclude the diagnosis of necrotizing fasciitis especially when other compatible signs and symptoms are present. It is also worth mentioning that an infective disease occurring in the exceptionally high vascular head and neck region will behave differently to the same condition present in other parts of the body, such as the abdomen or limbs. Some authors have even suggested that cervical necrotizing fasciitis is a different clinicopathological entity to craniofacial necrotizing fasciitis, with the former being associated with a 4-times higher mortality rate.² The young age of the patient, her general good health prior to the infection and pre-admission antibiotic therapy would also have contributed to a different disease progression.

It is important to reiterate that necrotizing fasciitis spreads along fascial planes, secondarily affecting muscle, subcutaneous tissue and skin. It follows that early detection and treatment of the condition could result in the skin being minimally affected. We opted for primary closure of the incisions after the two debridement procedures partly because the skin, although inflamed, looked viable and partly because of our intention to employ continuous suction and daily irrigation of the wound with hydrogen peroxide via three redi-vac drains left *in situ*. Following the publication of our article, another case of cervical necrotizing fasciitis has been reported in which the authors mentioned skin closure and insertion of a drain after initial surgical debridement.³ The presence of the suction drains in our

patient, coupled with the absence of muscle fascia, made it difficult to develop compartment syndrome, which, in any case, would have been manifested quite differently with local compression symptoms.

Whilst we did not specifically use the term 'necrosis' to describe our findings at the first operation, the pathological process of liquefaction necrosis is implicated when we stated 'a thin custard-coloured fluid was found along all the fascial planes from the skull base to the upper mediastinum'. The extensive debridement of the fasciae and adjacent muscle undertaken, as described, during the first procedure testifies to the above.

Although we achieved a successful outcome with primary closure, we are not advocating early skin closure in necrotizing fasciitis except in special, carefully considered circumstances. Furthermore, if primary skin closure is considered, we recommend the use of multiple continuous suction drains and daily irrigation of the wound with hydrogen peroxide (which acts as a chemical debriding agent). In retrospect, we feel that our management strategy did not compromise the care of the patient and in fact produced a relatively quick resolution of the condition as well as a good cosmetic result.

References

1. Cheshom J, Halstensen A, Haga T, Hoiby EA. Necrotizing fasciitis due to group A streptococci in Western Norway: incidence and clinical features. *Lancet* 1994; **344**: 1111–5.
2. Banerjee AR, Mutty GE, Moir AA. Cervical necrotizing fasciitis: a distinct clinicopathological entity? *J Laryngol Otol* 1996; **110**: 81–6.
3. Al-Ammar A, Mir SM. Cervical necrotizing fasciitis with facial nerve paralysis. *J Laryngol Otol* 2004; **118**: 573–5.

LETTER TO THE EDITOR

doi 10.1308/003588405X51056

Abdominal pus swabs – an on-going practice, should it continue?

N SARFRAZ, ARIF KHAN

Department of Surgery, Macclesfield District General Hospital, Macclesfield, Cheshire, UK

CORRESPONDENCE TO

N Sarfraz. M: +44 (0)7753 618104; E: dmoaman@email.com

It is common practice to send a culture swab when pus is encountered in the peritoneal cavity. This practice consumes staff time, effort and resources but do the results actually influence patient management? An audit was, therefore, conducted to assess the clinical utility of these intra-operative pus swabs and whether they had any effect on clinical decisions and patient management.

The clinical records of all patients undergoing emergency abdominal surgery from December 2000 to June 2001 were retrospectively analysed. Forty patients (age, 8–84 years; male:female ratio, 19:21) had had peritoneal pus swabs sent. A general policy had been followed of performing this investigation only if any free fluid or pus was identified. Of these 40 patients, 26 had appendicitis.

The results were analysed on a standard proforma to determine the site of the swab, the type and duration of antibiotics used, the extent of swab positivity, result availability before discharge, and the bacteriological spectrum. The most important aspect assessed was whether the antibiotics were changed subsequently during the hospital stay and if this decision was based on clinical grounds or on the basis of culture and sensitivity in keeping with the results of the initial pus swab. Patients who had subsequent wound or abdominal swabs were again analysed as to whether the swabs had any bearing on management.

Two major groups emerged according to site, with 26 of the 40 swabs from appendicectomies and 14 from cases of generalised peritonitis. Of the swabs, 55% had positive cultures: *Escherichia coli* (72%), anaerobes (31%) and mixed faecal organisms (31%) were the commonest isolates.

The 26 peri-appendicular swabs incidentally constituted 36.6% (26/71) of all appendicectomies over the same time period. Of these patients, 80% (21/26) were discharged home even before the culture report was available. Of patients who had appendix swabs sent, 76% (20/26) had a hospital stay of less than 4 days. The other six were detained either for social reasons or postoperative pyrexia (which persisted for 1–3 days in this group). Of the appendicectomy group, 89% (23/26) received antibiotics and 50% (13/26) had a positive culture. Interestingly, 81% (21/26) of all these patients were discharged before the culture report was available on day 5. No patient had antibiotics after the fourth day, and the culture result was not consulted when it was available. One patient came back with a minor wound infection, which was dealt with by wound drainage. A swab was sent but never followed up.

On evaluating pus swabs from the laparotomies for peritonitis (14/40 cases), nearly all (10/14) had a hospital stay of more than 7 days. Of these, 93% (13/14) had antibiotics except one patient who had a perforated duodenal ulcer; 64% (9/14) had positive cultures and all the reports were available before discharge. None had a change in antibiotics in view of the culture and sensitivity result. Thirteen of the above patients had faecal peritonitis and 29% developed wound infection, had a subsequent wound swab sent but these were all managed by the basic surgical principle of laying the wound open. Antibiotics were continued for more than 3 days in 10/14 patients, one patient was transferred to another ITU and 3 died due to the high mortality associated with faecal peritonitis.

In summary, we found that 90% of the 40 patients received empirical antibiotics, 80% of the appendicectomies were discharged before the culture results were available and in none of the 40 patients were the antibiotics changed based on a culture result.

Abdominal infections are treated by drainage, control of the source of infection, and antimicrobial agents active against expected pathogens.¹ There is controversy over the value of diagnostic cultures and whether expanded antimicrobial coverage is required.² The spectrum of bacterial isolates from the peritoneum is very constant across the world, the only changes being slight variations in the percentages of the various organisms. Intestinal flora are not permanently affected by short-term antibiotic therapy and older antibiotics are appropriate first-line therapeutic agents for infections caused by normal intestinal flora.³

It has also been shown that the sensitivity of the organisms to the antibiotics given does not influence the incidence of later infective complications or the outcome. Results from a prospective open study of 51 patients with secondary peritonitis suggested that micro-organisms invading the peritoneal cavity and respective antibiotic treatment are not the major determinants of the clinical course of peritonitis. Rather, the determinants are the degree of the systemic inflammatory response judged by scoring systems, the length of history, and local and systemic concentrations of endotoxin. Intra-abdominal microbiological findings did not correlate with severity of illness, the later incidence of infective complications, or the final outcome.⁴

The anatomical origin of bacterial contamination and microbiological findings are not major predictors of outcome. The pre-operative physiological derangement, the surgical clearance of the infectious focus and the host response to treatment are established prognostic factors.⁵ Identification of the exact microbiology is only necessary in severely ill patients with complicated infections who can be identified by scoring systems.⁶ On the other hand, these studies do not mention whether they would recommend the routine use of culture and sensitivity in uncomplicated intra-abdominal infections.⁷

In our small study in patients undergoing emergency abdominal surgery for localised/generalised peritonitis, the pus swab sent at the time of operation did not influence clinical decision-making to alter the empirical or subsequent antibiotic regimens or change patient management and could not justify its usefulness in this era of financial stringency.

References

1. Bohnen JM. Antibiotic therapy for abdominal infection. *World J Surg* 1998; **22**: 152–7.
2. Krepel CJ, Gohr CM, Edmiston CE, Condon RE. Surgical sepsis: constancy of antibiotic susceptibility of causative organisms. *Surgery* 1995; **117**: 505–9.
3. Schoeffel U, Jacobs E, Ruf G, Mierswa F, von Specht BU, Farthmann EH. Micro-organisms and the severity of peritonitis. *Eur J Surg* 1995; **161**: 501–8.
4. Farthmann EH, Schoeffel U. Epidemiology and pathophysiology of intraabdominal infections (IAI). *Infection* 1998; **26**: 329–34.
5. Wilson SE, Faulkner K. Impact of anatomical site on bacteriological and clinical outcome in the management of intra-abdominal infections. *Am Surg* 1998; **64**: 402–7.
6. Koperna T, Schulz F. Prognosis and treatment of peritonitis. Do we need new scoring systems? *Arch Surg* 1996; **131**: 180–6.
7. Burnett RJ, Haverstock DC, Dellinger EP, Reinhart HH, Bohnen JM, Rotstein OD *et al.* Definition of the role of enterococcus in intraabdominal infection: analysis of a prospective randomized trial. *Surgery* 1995; **118**: 716–3.