TECHNICAL SECTION:

Technical tutorials, notes and tips

Orthopaedic screws

Use of a Vicryl lasso during insertion of locking screws

HV Dabke, DJ Woodnutt

Department of Orthopaedics, Morriston Hospital, Swansea, UK

During nailing of long bones, retrieval of locking screws that get 'lost' in the soft tissues can be difficult especially in obese patients. We describe a simple technique to avoid this problem.

A loop of No. 1 Vicryl is 'lassoed' around the neck of the screw and a clip is attached to its end. The screw is inserted as usual. If the screw gets mal-tracked in the soft tissues, a tug on this Vicryl loop will help to get the screw out. Just before full tightening of the screw, the Vicryl loop is either pulled out or cut. Since it is absorbable, cutting the loop does not result in any problem.

This method may be used when screws are inserted in 'difficult' areas (e.g. arthroscopically) or while locking anteroposteriorly during femoral nailing, where formal exposure is not desirable.

Correspondence to: Mr HV Dabke, 2/1 Teal House, St Anne's Close, Lincoln LN2 5TA, UK

Lost locking screws – an avoidance technique

Stephen M Blake, Mark D Brinsden, Simon Smith

Department of Trauma and Orthopaedic Surgery, Royal Cornwall Hospital Treliske, Truro, Cornwall, UK

Insertion of locking screws into intramedullary nails can often be troublesome in patients with large limbs. We describe a simple technique to ensure the screw is not lost in the soft tissues. Secure an O-Vicryl tie around the neck of the screw.

Place the screw onto the screwdriver and hold the tie under tension. This ensures the screw is engaged securely and, if it were to displace, it can easily be retrieved by withdrawing the tie. After the proximal cortex has been engaged, the tie can be cut and the screw fully advanced.

Correspondence to: Mr SM Blake, Orthopaedic Registrar, North Devon District Hospital, Raleigh Park, Barnstaple, Devon EX31 4JB, UK
E-mail: stephenblake_uk@yahoo.co.uk

A simple technique for removal of screws with damaged heads

Mohammed Aqeel Bhutta

Rochdale, UK

Removal of metalwork is a relatively straight-forward procedure; however, screw removal can be complicated by 'stripping' or worn screw heads. Accepted methods for removing screws in this situation include 'overdrilling', removal from the opposite side, and coring out of the screw heads. Our technique involves the use of a small swab wrapped around the tip of the screwdriver which creates an interference fit between the screwdriver and the screw head creating more torque for removal. The swab thickness can be varied with the number of folds used. This is a simple, yet effective, technique.

Correspondence to: Dr Mohammed Aqeel Bhutta, 8 Edinburgh Way, Rochdale OL11 2NQ, UK

General techniques

Easy hand tying of a short suture

Kevin J Daly, Francis Dix, Charles N McCollum

Academic Surgery Unit, South Manchester University Hospital, Manchester, UK

We report a useful technique for hand tying a surgical knot when the remaining available suture would normally be too short.

A spare tie or length of suture is passed through a loop of the suture to be tied (Fig. 1). Both ends of the spare supporting suture are then held together under tension in the tying hand. The end of the suture can then be tied more easily with or without the aid of a needle holder to hold the needle (Fig. 2). This method is simple and avoids an instrument tie of material prone to fracture such as polypropylene. This technique can be used at the end of a

Figure 1 A supporting suture is passed through the loop of the suture to be tied.

Figure 2 A suture being tied with the aid of a supporting suture loop.

wound or in tightening a slack loop in a vascular anastomosis after the ends have been tied. In the latter case, a nerve hook may help remove the slack allowing the supporting suture to be passed through the loop and tied to a new interrupted suture.

Acknowledgement

We would like to thank Helen Carruthers for providing the illustrations.

Correspondence to: Mr Kevin Daly, Lecturer in Surgery, Room ATR4, Education and Research Centre, South Manchester University Hospital, Southmoor Road, Manchester M23 9LT, UK.

Tel: +44 161 291 5805; Fax: +44 161 291 5806; E-mail: kdaly@fs1.with.man.ac.uk

Letter in response to technical tip by Fang, Moran and Paul

Re. The left-handed use of scissors

Ann R Coll Surg Engl 2003; **84**: 357

Michael Pegum

Charlemont Clinic, Charlemont Mall, Dublin, Ireland

I was disappointed to see this well-known technique,¹ for the use of scissors with the left hand, put forward as being satisfactory. While it may be acceptable for cutting sutures, it certainly does not give adequate control for sharp dissection, and should not be encouraged. All surgeons should be ambidextrous.

Most scissors are made for right-handed use, so that pressure with the radial side of the thumb on the loop in the handle, away from the plane of the hand, will push the blades together as they are closed, and ensure efficient cutting.

When in use in the left hand, with the scissors held in the same way as in the right hand, the thumb loop of the scissors must be pulled towards the plane of the hand with the ulnar side of the thumb, while the other handle loop is pushed with the back of the distal phalanx of the little finger.

Once learned, this technique (known, I am sure, to every left-handed surgeon) is easy to use and gives full control of the instrument.

Reference

1. Fang C, Moran M, Paul A. The left-handed use of scissors. *Ann R Coll Surg Engl* 2002; **84**: 357.

Correspondence to: Mr Michael Pegum, Orthopaedic Surgeon, Charlemont Clinic, Charlemont Mall, Dublin 2, Ireland. Tel: +353 1 269 7514; Fax: +353 1 260 0569; E-mail: mpegum@eircom.net

Abdominal surgery

A simple method for locating the linea alba in midline laparotomy incisions

OJH Harley, TF Browne

Department of Vascular Surgery, Broomfield Hospital, Chelmsford, Essex, UK

In making a midline abdominal incision, the linea alba is occasionally difficult to locate when the anterior rectus sheath is reached. We describe a safe, simple and quick way of identifying the linea alba. The rectus sheath is entered and probed transversely between the rectus muscle fibres and the anterior rectus sheath with closed scissors. Unimpeded passage of the probe for several centimetres suggests direction away from the midline. Abrupt obstruction of the

probe reveals the position of the midline. This method is not useful at levels inferior to the arcuate line. In the patient who has previously had a paramedian incision, the technique may not reliably locate the linea alba.

Correspondence to: TF Browne, Consultant Vascular Surgeon, Department of Vascular Surgery, Broomfield Hospital, Chelmsford, Essex CM1 7ET, UK
Tel/Fax: + 44 1245 515222

Hyaluronidase in the reduction of prolapsed colostomy

Arindam Chaudhuri, Amit Prasai

Department of General Surgery, Peterborough District Hospital, Peterborough, UK

Stomal prolapse may be difficult to reduce, especially if long-standing, due to oedema of the prolapsed segment. The authors describe a method of easily decreasing the oedema, thereby facilitating reduction. Hyaluronidase (3000 units in 20 ml of water) is injected submucosally into the prolapsed segment of the colon. Excess fluid can be milked away within the next few minutes. Once the volume of the prolapsed segment has diminished, it can be successfully reduced. Long-standing prolapsed stomas need prompt reduction prior to development of ischaemic changes at the apex of the prolapsed segment. Other methods such as application of desiccating agents such as sugar may be successful,¹ but are messy and time-consuming, while this method achieves quite rapid results. Hyaluronidase (hyaluronoglucosidase) is a widely present endoglycosidase that cleaves hexosamine bonds, acting on both hyaluronic acid and chondroitin sulphate. It acts by depolymerising hyaluronic acid, with resultant breakdown of the extracellular matrix. This allows oedema fluid to be milked through the microscopic gaps created in the extracellular compartment.² It is contraindicated in the presence of infection and malignancy.² One author has successfully applied hyaluronidase in the reduction of difficult rectal prolapse.³ The authors recommend this technique in similar scenarios where effective and rapid reduction of oedema is necessary to reduce the prolapsed organ.

References

1. Coburn 3rd WM, Russell MA, Hofstetter WL. Sucrose as an aid to manual reduction of incarcerated rectal prolapse. *Ann Emerg Med* 1997; **30**: 347–9.
2. DeVries CR, Miller AK, Packer MG. Reduction of paraphimosis with hyaluronidase. *Urology* 1996; **48**: 464–5.
3. Chaudhuri A. Hyaluronidase in the reduction of incarcerated rectal prolapse: a novel use. *Int J Colorectal Dis* 1999; **14**: 264.

Correspondence to: Mr A Chaudhuri, Specialist Registrar, Department of General Surgery, Peterborough District Hospital, Thorpe Road, Peterborough PE3 6DA, UK
Tel: +44 1733 874000; Fax: +44 1733 875013, E-mail: arindam@chaudhuri.freemove.co.uk

A simple technique for fenestration of an hepatic cyst

Paul Glen, Paul G Horgan

Department of Academic Surgery, Queen Elizabeth Building, Glasgow Royal Infirmary, Glasgow, UK

A window is made into the cyst and the compressed liver tissue making up the roof is serially coagulated with the Ligasure electrocautery device (Autosuture, United States Surgical Corp., Norwalk, CT, USA). The sealed tissue may then be divided with scissors without fluid loss. This procedure is continued around the circumference of the cyst until it is completely de-roofed.

Large hepatic cysts compress the surrounding parenchyma, ducts and vessels into a dense capsule. Division of this tissue may result in bile leak or major haemorrhage. The Ligasure device has proved useful in achieving hepatic parenchymal division and has contributed to a low blood loss for hepatectomies in this unit (425–700 ml).¹

Reference

1. Horgan PG. A novel technique for parenchymal division during hepatectomy. *Am J Surg* 2001; **181**: 236–7.

Correspondence to: Mr Paul G Horgan, Department of Academic Surgery, Queen Elizabeth Building, Glasgow Royal Infirmary, Glasgow G31 2ER, UK
Tel: +44 141 211 4871, Fax: +44 141 552 3229