Use of the Shirodkar suture in shoulder surgery

S Cooper, D Acton, MJ Curtis

Department of Orthopaedic and Trauma Surgery, Kingston Hospital NHS Trust, Kingston, Surrey UK

Displaced four-part proximal humeral fractures in the elderly are usually treated by shoulder hemiarthroplasty. Achieving pain relief and functional range of movement depends on restoration of humeral anatomy and reconstruction of the rotator cuff to permit early mobilisation.¹ In these patients, the greater and lesser tuberosities are often osteoporotic and fragmented. After insertion of the prosthesis, standard surgical technique involves drilling holes through the greater and lesser tuberosities through which monofilament wires or heavy non-absorbable sutures are passed to fix the tuberosities together around the fin of the prosthesis. This procedure can be awkward and with poor quality bone the wires may cut out, compromising fixation and subsequent rotator cuff function. We have found that the use of the Shirodkar suture (Ethicon RS21, Johnson & Johnson International) considerably improves the ease and outcome of surgery. Originally designed for use in gynaecological surgery for cerclage of the incompetent cervix, the suture is a 5-mm wide, polyester (non-absorbable) tape with double-ended, blunt round-bodied 65-mm needles. The needles are passed

from the articular side of the rotator cuff at the bone-tendon junction and the tape tied over the tuberosities, the width of the tape spreading the loads across the tuberosities and providing reliable fixation. We have used this tape for tuberosity re-attachment in a series of 40 shoulder hemiarthroplasties without loss of fixation, with no local reaction or sinus formation and can recommend its use.

Reference

1. Cofield RH. The shoulder and prosthetic arthroplasty. In: Evans CM. (ed) *Surgery of the Musculoskeletal System*, 2nd edn. New York: Churchill Livingstone, 1990; 1571–91.

Correspondence to: Sarah Cooper, Department of Orthopaedic and Trauma Surgery, Kingston Hospital NHS Trust, Kingston, Surrey, KT2 7BE, UK

Use of adrenaline and bupivacaine to reduce bleeding and pain following harvesting of bone graft.

GA Cheeseman, A Chojnowski

Department of Orthopaedics and Trauma, Norfolk and Norwich University Hospital, Norwich, UK

Following bone grafting procedures, patients frequently complain of severe pain and bleeding from the donor site. To counter these symptoms, we routinely use a swab soaked in 10 ml bupivacaine (0.25%) and adrenaline (1/200,000) inserted into the bone following harvesting. This remains *in situ* from the time of harvesting until removal prior to closure. It is our anecdotal observation that bleeding and pain are markedly reduced using this logical method. Pain relief also appears superior when compared to infiltration of the wound after closure. We strongly recommend this method as an easy, cost-effective and successful step towards improving the patient's experience of surgery as well as the clinical outcome.

Correspondence to: Mr A Chojnowski, Department of Orthopaedics, Norfolk and Norwich University Hospital, Colney Lane, Norwich NR4 7DZ, UK
Tel: +44 1603 286286

Figure 1 Diagrammatic representation of tuberosity re-attachment with Shirodkar sutures. (A) implant; (B) greater tuberosity (supraspinatus, infraspinatus and teres minor); (C) lesser tuberosity (subscapularis); (D) suture.

An aid to external fixator application

ART McBride, PRM Black

Department of Trauma and Orthopaedics, Musgrove Park Hospital, Taunton, UK

Maintenance of reduction of long-bone fractures and application of external fixation devices can be frustrating, time consuming and necessitate the use of an assistant. Following reduction of the limb fracture under X-ray guidance, the Evacuated Bead Bag (trunk with cut-out model; Teasdale Hospital Equipment Ltd) is moulded firmly around the limb and a suction applied, forming an accurate mould of

Figure 1 The Evacuated Bead Bag in position.

the reduced limb shape (Fig. 1). The limb is then removed from the mould, prepped and draped in an appropriate fashion and replaced in the draped mould. The fracture is then screened to confirm maintenance of reduction. In our experience, the mould allows immediate re-alignment of the majority of fractures without further manipulation. External fixation can then be applied in a conventional fashion. With the fixator in position, final adjustments can be made with ease under X-ray guidance. This technique allows ease of application by a single surgeon and also prevents unnecessary soft tissue stresses during fixator application by immobilising the limb and reducing axial rotation, valgus and varus forces at the fracture site.

Correspondence to: Mr ART McBride, Department of Trauma and Orthopaedics, Musgrove Park Hospital, Taunton, UK
Tel: +44 1823 344028; Fax: +44 1823 342474; E-mail
andrewmcbride@hotmail.com

ERRATUM

Ann R Coll Surg Engl 2003; **85**: 207

A simple technique for removal of screws with damaged heads

MA Bhutta, PD Dunkow and ME Lovell
South Manchester University Hospitals NHS Trust

The correct authors and hospital for this Technical Tip are given above.