individually threaded through this loop and pulled through the canal using the primary suture. By the sequential use of the primary sutures in each bony canal, a single suture may be passed, in either direction, through as many bony tunnels as is required to complete the repair. Each primary suture may be used many times and to draw repair sutures in either direction. Once the repair is secure, the primary suture is removed.

DISCUSSION

Using this technique, the danger of crushing or cutting previously passed repair sutures is abolished, thereby preserving the strength of the repair suture. We have found this a simple technique to use, requiring no specialised equipment; it is particularly valuable in the repair of patellar tendons where multiple sutures must be woven through several bony canals.

References

1. Bucholz RW, Heckman JD. *Rockwood and Green's Fractures in Adults*, 5th edn. Philadelphia, PA: Lippincott, Williams and Wilkins, 2001.
2. Terry Canale S. *Campbell's Operative Orthopedics*, 9th edn. St Louis, MO: Mosby, 1998.

SQUASH – simple qualitative user-friendly amplitude sensor of hand-strength – a non-invasive monitor during carotid endarterectomy under local anaesthetic

R JOY, MA CHADWICK, JD ARTHUR, RJ MCCARTHY, IC CURRIE
Department of Vascular Surgery, South Devon Healthcare Trust, Torbay, Devon, UK

CORRESPONDENCE TO

R Joy, Department of Vascular Surgery, South Devon Healthcare Trust, Torbay, Devon
M: +44 (0)7968 619760; E: rajivejoy@yahoo.com

BACKGROUND

There is currently no objective, reliable and recordable method of measuring contralateral motor function during a carotid endarterectomy (CEA) under local anaesthesia. A major complication of CEA is cerebral hypoperfusion requiring shunting. We describe a novel technique to determine the need for shunting.

TECHNIQUE

SQUASH is a non-invasive technique, which allows assessment of contralateral upper limb grip strength during CEA under local anaesthesia without sedation (Fig. 1). The SQUASH system is shown in Figure 2. The voluntary grip strength is transduced to zero and the patient is instructed to grip the bag maximally and its trace calibrated for maximum 'squash' (Fig. 3). The patient is asked to squash (Fig. 4) during the procedure especially before

Figure 1 SQUASH bag.

Key points Figs 1 & 2:

- Fixed volume bag (100 ml normal saline), which fits the palm.
- Drawing up needle to which extension pressure tubing with Luer lock is attached.
- Reinforcing tape.
- The 'drawing up' needle, attached to the CVP tubing at the Luer lock, is inserted into saline bag via the rubber seal and reinforced (we use Elastoplast).
- Central venous pressure (CVP) transducer and non-expansile pressure tubing with Luer lock attachment.
- The CVP tubing is filled from the bag and flushed to ensure no air bubbles are present.
- * Monitor with flow module (standard anaesthetic monitor).

Figure 2 Schematic diagram of SQUASH system.

Figure 3 Maximum SQUASH.

Figure 4 Normal SQUASH.

Figure 5 Abnormal SQUASH.

Figure 6 Abnormal SQUASH.

and after the carotid cross clamping. Reduction in grip strength is used to determine the need for shunting. We noticed a decrease in the frequency and/or amplitude of the waveforms during episodes of hypoperfusion (Figs 5 and 6) in 2 patients out of 92 who were shunted and made a full recovery. There were no adverse outcomes intra-operatively using this technique.

DISCUSSION

CEA is potentially complicated by cerebral hypoperfusion resulting in irreversible neurological deficit. Intra-operative

indicators of cerebral hypoperfusion or hypofunction like transcranial Doppler, EEG, infrared spectroscopy, shunt pressures or awake-patient monitoring can be used to determine the need for selective shunting. Alternatively, the SQUASH provides a visual representation of grip strength in real time and is a direct measure of cortical sensorimotor integrity, which indirectly measures cerebral perfusion. It is easy to assemble, inexpensive, re-usable and non-invasive. Whether criteria to consider shunt placement can be derived from the waveform requires future study. Until then, we believe this will remain a promising means of non-invasive cerebral perfusion monitoring during CEA under local anaesthesia.

Protecting the distal anastomosis in diabetics undergoing dorsalis pedis bypass

HANY ZAYED, NICHOLAS FASSIADIS, HISHAM RASHID
Department of Vascular Surgery, King's College Hospital,
London, UK

CORRESPONDENCE TO

Hany Zayed, Senior Clinical Fellow, Department of Vascular Surgery, King's College Hospital, London SE5 9RS, UK
M: +44 (0)7899 790607; E: zayedha@yahoo.com

BACKGROUND

Peripheral arterial disease in diabetic patients is characterised predominantly by long segmental occlusion of the tibial arteries with patent segments of the dorsalis pedis artery (DPA). This allows distal bypass to the DPA using vein grafts.¹ Wound complications are frequent due to the presence of foot tissue loss and oedema.² Wound dehiscence with exposure of the DPA anastomosis is a serious postoperative complication which could lead to limb loss.³ This report describes specific measures that reduce the above risk.

TECHNIQUE

Strict leg elevation for 48 h prior to surgery is of paramount importance. Occasionally, epidural analgesia is required in patients with severe rest pain in order to achieve this. The position of the DPA is identified and marked using a hand-held Doppler, if a signal is obtainable. The incision is made 1.5 cm medial and parallel to the DPA (overlying the extensor hallucis longus tendon; Fig. 1). Dissection is continued down to the deep fascia prior to flap mobilisation to produce a lateral fascio-cutaneous flap. The DPA is dissected and the bypass is performed using a reversed autologous vein graft. The graft is tunnelled laterally through the interosseous membrane. Skin closure is performed with polyglactine 3/0 without fascial approximation. Leg elevation is maintained until skin healing is achieved.