Technical Section

TECHNICAL NOTES & TIPS

TECHNICAL NOTES

Using an extracorporeal suture loop to aid retroperitoneal pelvi-ureteric anastomotic suturing during laparoscopic pyeloplasty

RW RITCHIE, N THIRUCHELVAM, A ADAMSON
Department of Urology, Royal Hampshire County Hospital,
Winchester, UK

CORRESPONDENCE TO

RW Ritchie, Department of Urology, Royal Hampshire County
Hospital, Romsey Road, Winchester SO22 5DG, UK
E: robritchie@doctors.org.uk

BACKGROUND

Intracorporeal suturing during laparoscopic surgery is time-consuming but fundamental to performing a pyeloplasty. In addition, laparoscopic pyeloplasty is routinely performed via an extraperitoneal approach limiting available suturing space adding to potential technical difficulties.¹ We describe a technique that eliminates the need for an initial intracorporeal knot to begin the anastomosis.

TECHNIQUE

The method involves forming an initial suture loop outside the body. We recommend attaching an artery forceps to the operative

drape and tying an accurate reef knot around the points of the forceps (Fig. 1A). Removing the forceps leaves a small loop at the opposite end of the suture to the needle. The needle and suture are delivered into the retroperitoneal space. The first pass is taken of the renal pelvis and the needle subsequently passed through the ureter. We routinely use a Connel suture.² The needle is passed through the formed loop (Fig. 1B) and pulled tight, securing the beginning of the anastomosis. The formation of the anastomosis continues with tension on the running suture, knotting in a standard intracorporeal fashion at the end.

DISCUSSION

We have now used this technique in 15 consecutive procedures. Fourteen procedures were entirely uncomplicated; in 1 case, there was a prolonged urinary leak. Creating the initial knot extracorporeally ensures an accurate and safe initial hand tie under direct vision. This eliminates the need for an initial laparoscopic intracorporeal knot tie to begin the anastomosis, saving time and allowing rapid apposition of the renal pelvis and ureter. This, in turn, allows the rapid generation of 'momentum' to the suturing process that is important to the timely completion of an accurate anastomosis.

References

1. Eden C, Cahill D, Allen J. Laparoscopic dismembered pyeloplasty: 50 consecutive cases. *BJU Int* 2001; **88**: 526–31.
2. Cartledge J, Blacker T, Joyce A. Laparoscopic pyeloplasty – Lessons from the learning curve. *Urol News* 2004; **8**: 6–10.

Figure 1 Initial extracorporeal knot aids suturing. (A) Illustration shows artery forceps (F) grasping drape (D) as reef knot is tied (K). (B) Photograph shows extraperitoneal space, after bites through renal pelvis (P) and ureter (U); needle is passed through previously formed loop (L) and pulled tight.

Are tracheal dilators useful in stomal stenosis?

OWEN JUDD, COSTA REPANOS, MICHAEL W BRIDGER
Department of Otolaryngology, Derriford Hospital, Plymouth, UK

CORRESPONDENCE TO

Owen Judd, Senior House Officer in ENT, Derriford Hospital,
Derriford Road, Plymouth PL6 8DH, UK
E: owenjudd@doctors.org.uk

BACKGROUND

Tracheostomy has been performed since ancient times. In 1833, Armand Trousseau invented an instrument designed to dilate the incision into the airway, to allow easier passage of a tracheostomy

Figure 1 The Trousseau tracheal dilator.

Figure 2 Hole cut to simulate stenosed stoma.

Figure 3 Stenosis dilated 50%.

Figure 4 Stenosis dilated 100%.

Figure 5 Slit cut to simulate traditional tracheostomy incision.

Figure 6 Slit dilated 50%.

Figure 7 Slit dilated 100%.

tube (Fig. 1).¹ We have observed that tracheal dilators are always included in the pack when changing a tracheostomy tube in an established tracheostomy or when there is stenosis of the stoma.²⁻⁴ We believe, however, that, in these circumstances, the Trousseau dilator impedes the insertion of a tube and could be dangerous in an emergency.

TECHNIQUE

We used two sheets of rubber: one cut with a circular hole to represent a stenosed stoma and one with a vertical slit to represent a tracheostomy incision. We then measured the vertical and horizontal diameters of the holes before and after using the tracheal dilator to 50% and 100% of its capacity. In a circular stoma, the vertical height decreases from 5 mm to 4 mm (20%) when dilated (Figs 2-4). Dilating a vertical slit, however, only reduces the vertical diameter by 10% from 20 mm to 18 mm (Figs 5-7).

DISCUSSION

A slit incision, when dilated, becomes a diamond-shaped aperture through which a tube can easily be passed. In the circular hole, the instrument constricts the available aperture by 20%. Our simple experiment has confirmed our suspicions that using a tracheal dilator in cases of annular stomal stenosis can make the task of inserting a tracheostomy tube more difficult, by narrowing the aperture. Is it time for our wards and emergency departments to consign the Trousseau dilator to the historical archive?

References

1. Sebastian A. *A Dictionary of the History of Medicine*. London: Taylor & Francis, 1999
2. Bramwell KJ, Davis DP, Cardall TV, Yoshida E, Vilke GM, Rosen P. Use of the Trousseau dilator in cricothyrotomy. *J Emerg Med* 1999; **17**: 433-6.
3. Narrod JA, Moore EE, Rosen P. Emergency cricothyrotomy – technique and anatomical considerations. *J Emerg Med* 1985; **2**: 443-6.
4. Mirza S, Cameron DS. The tracheostomy tube change: a review of techniques. *Hosp Med* 2001; **62**: 158-63.

A technique to avoid leg-length discrepancy in total hip arthroplasty

M HOSSAIN, AK SINHA

Department of Trauma and Orthopaedic Surgery, Ysbyty Gwynedd, Bangor, UK

CORRESPONDENCE TO

M Hossain, Department of Trauma and Orthopaedic Surgery, Ysbyty Gwynedd, Bangor LL57 2PW, UK
T: +44 (0)1248 384384 bleep 241; F: +44 (0)2920 930 162;
E: munierh@doctors.org.uk

BACKGROUND

Leg-length discrepancy is common after hip arthroplasty.¹ Surgeons conventionally check leg-length by comparing the position of the two knees. But lateral positioning can rotate the pelvis and give a misleading impression. Calliper use is described to achieve leg-length equality.² We describe a simple technique.

TECHNIQUE

Pre-operatively, leg-length discrepancy is measured and a decision made on required length. Following exposure and before hip