glove for the safe endoscopic removal of a sharp gastric foreign body.

Gastrointest Endosc 2000; **52**: 127–9.

5. Bertoni G, Sassatelli R, Conigliaro R, Bedogni G. A simple latex protector hood for safe endoscopic removal of sharp-pointed gastroesophageal foreign bodies. *Gastrointest Endosc* 1996; **44**: 458–61.

TECHNICAL TIPS

k-Balls – a novel method of covering Kirschner wires

ANDREW SHEPHERD, PETER COX
Department of Trauma and Orthopaedics, Royal Devon & Exeter Hospital, Exeter, UK

CORRESPONDENCE TO

Andrew Shepherd, 5 Model Cottages, Clyst St George, Exeter EX3 ORD, UK
E: andrewshepherd@doctors.org.uk

We present a novel method for covering the ends of percutaneous Kirschner wires using Caprolactone (Orthoinvent, c/o Farlacombe House, Farlacombe, Bickington TQ12 6PD, UK: enquires@orthoinvent.co.uk). Caprolactone is a biodegradable thermoplastic. It is supplied in granular form and becomes mouldable and translucent when placed in hot water (58–60°C), again becoming solid at room temperature. We mould the Caprolactone into balls around the ends of the k-wires. The balls will not fall off the wire and can easily be removed in a clinic setting. Caprolactone is non-hazardous and can be re-used by submerging it in hot water.

Figure 1 Covering Kirschner wires with Caprolactone.

Use of a vaginal speculum to assist difficult urethral catheterisation

A NARAYANAN, M O'BRIEN, RS RAJPUT, S SUNDARARAMAN
South Warwickshire General Hospital, Warwick, UK

CORRESPONDENCE TO

A Narayanan, Department of Surgery, West Wales General Hospital, Glangwili, Carmarthen, Dyfed SA31 2AF, UK.
E: aravinn_72@yahoo.com

Visualisation of the male urethral meatus for catheterisation can be difficult in situations such as morbid obesity, gross penile oedema or huge inguinoscrotal herniation. A novel technique in our experience is the use of a vaginal speculum to overcome this. Two persons are ideally required for effective use – one to stabilise the device within the penis, to visualise the meatus whilst the other performs the catheterisation itself. This technique allows visualisation of the urethral meatus and hence enables catheterisation to be performed with relative ease in such patients.

A novel use of the Sommerlad pen: preparing the nail during nailed repair

AJK PATEL¹, MJD WAGTSTAFF²

¹Dept of Plastic Surgery, Leicester Royal Infirmary, Leicester, UK

²Dept of Plastic Surgery, Nottingham City Hospital, Nottingham, UK

CORRESPONDENCE TO

AJK Patel, Department of Plastic Surgery, Leicester Royal Infirmary, Infirmary Square, Leicester LE1 5WW, UK
T: +44 (0)1162 585786; F: +44 (0)1162 585852;
E: aj.patel@virgin.net

During surgical nailed repair, if available, the patient's nail plate is often re-sited in the nail fold. This prevents formation of adhesions of the eponychium and forms a biological dressing and splint over the repair. Prior to using the nail in this way, it needs to be trimmed and cleaned, as there is commonly a thin layer of innominate tissue that could adhere to the repaired nailed. We find using the handle of the Sommerlad marking pen a very effective method for this. The handle

Figure 1 (A) The Sommerlad pen; (B) preparing the nail.

of the Sommerlad pen has a surface similar to a rat-tail rasp. The convex surface of the handle fits snugly to the concave undersurface of the nail, which can then be adequately prepared.

Use of screen bar to assist in below-knee amputation

ABDOLALI ASSARIAN, RAYMOND CHUNG, PR THOMAS
Department of Surgery, St Helier Hospital, Carshalton, Surrey, UK

CORRESPONDENCE TO

A Assarian, 9 Fitzroy Close, Bassett, Southampton SO16 7LW, UK
F: +44 (0)23 8024 0320; E: aliassarian@yahoo.com

The routine technique for below-knee amputation involves laying the limb flat on the theatre table and manoeuvring it as required by a sec-

Figure 1 Screen bar in place.

Figure 2 Position of leg.

Figure 3 Surgeon is operating without assistance.

ond assistant. We use a screen bar, commonly used by the anaesthetic team, and hence freely available to all surgical theatres, to support the lower limb (Fig. 1). The lower limb is held in a stable position, with the hip and knee flexed and posterior part of knee lying over screen bar (Fig. 2). This provides an excellent support for the limb, allowing the surgeon to operate comfortably and independently, with greater circumferential view of the operating field (Fig. 3).

Forced expiration for chest tube removal – the other uses of a syringe

T SINGH¹, S RANDHAWA²

Departments of ¹Cardiothoracic Surgery and ²Accident and Emergency Medicine, New Cross Hospital, Wolverhampton, UK

CORRESPONDENCE TO

T Singh, Department of Cardiothoracic Surgery, New Cross Hospital, Wolverhampton WV10 0QP, UK. E: talvinder13@hotmail.com

The incidence of pneumothorax following chest tube removal can be high. The British Thoracic Society recommends that the chest tube should be removed briskly while the patient performs the Valsalva's manoeuvre or during expiration. Sometimes, it can be difficult for patients to comprehend the steps involved in producing this effect. We have devised a novel technique; after taking a deep breath, the patient is asked to blow through the tip of a syringe allowing the plunger to move backwards through the barrel. This technique produces enough positive intrathoracic pressure and decreases the chances of a pneumothorax forming significantly.

COMMENT ON

doi 10.1308/003588407X187603

P Marsh, JS Clark. The spouted colostomy. *Ann R Coll Surg Engl* 1997; **89**: 78

The spouted stoma

ANTHONY LANDER

Department of Paediatric Surgery, Diana Princess of Wales Children's Hospital, Birmingham, UK

CORRESPONDENCE TO

Anthony Lander, Consultant Paediatric Surgeon, Diana Princess of Wales Children's Hospital, Steelhouse Lane, Birmingham B4 6NH, UK
E: a.d.lander@googlegmail.com

I was surprised at the article and only comment because it was invited by the editor. In paediatric practice, colostomies are frequently spouted and have been for many years; this is not a new technique. Can I describe the same technique for the jejunum and put my name to it?

Interestingly, in sick premature babies, the bowel may not tolerate eversion at the time of surgery and one can leave the small or large bowel to evert itself to form a nice spout fixing only the bowel to the fascia a centimetre or so from its distal end.

An interesting question in relation to spouted urostomies is why the skin grows up to the top of the spout, a phenomenon not seen with the intestine.