THE EFFECT OF DIET ON THE PATHOLOGICAL CHANGES IN RATS WITH NEPHROTOXIC NEPHRITIS *

JOSEPH E. SMADEL, M.D., AND LEE E. FARR, M.D.

(From the Hospital of the Rockefeller Institute for Medical Research,
New York, N. Y.)

Nephrotoxic nephritis has been regarded by Masugi and the majority of recent workers as the experimental renal lesion which most closely simulates Bright's disease in man.^{1, 2} Previously we demonstrated that severe acute nephritis induced in rats by means of nephrotoxin generally progressed to a chronic disease with ultimate kidney failure.^{2, 3} An occasional rat, however, when fed a stock diet, recovered from the acute renal injury, while a number of animals with chronic nephritis of a year's duration still maintained a normal urea clearance value. This variation in the response of individual animals suggested that the course of the disease might be influenced by internal environmental factors, the "milieu intérieur" of Claude Bernard. Since diet has long been considered an important therapeutic measure in the treatment of Bright's disease, and has been shown to affect the clinical picture in one type of experimental renal injury,⁴ it seemed desirable to study its influence on the diffuse glomerulonephritis which results from injection of antikidney serum. Accordingly, severe nephrotoxic nephritis was induced in a large group of rats and the course of the renal disease was observed for 10½ months to determine the influence of different diets. Detailed clinical and chemical studies on 48 of the animals in this group are to be reported elsewhere. The present paper deals with the pathological changes found in these rats and in others followed for a shorter period; renal lesions observed in normal rats maintained on one of the experimental diets are also described.

MATERIALS AND METHODS

Nephrotoxic Serum: Antikidney serum was prepared in rabbits by immunization with sterile suspensions of perfused rat kidney. Throughout this study a single serum was used, viz. that from

* Received for publication November 19, 1938.

Rabbit 4557 which was always capable of inducing severe nephritis in rats receiving a total of 0.65 cc. per 100 gm. of body weight, in three divided doses given on consecutive days.⁵

Animals: Young hooded rats of the Whelan strain were injected when they weighed about 100 gm. Males and females were distributed equally in the respective diet groups.

Diet: Rats were fed on one of three isocaloric diets. Each ration contained 27 per cent fat, 4 per cent Osborne and Mendel salt mixture No. 1,* and vitamins plus the following constituents: Diet L — 5 per cent protein and 64 per cent carbohydrate; Diet B — 18 per cent protein and 51 per cent carbohydrate; Diet H — 40 per cent protein and 29 per cent carbohydrate. The protein consisted of Lactalbumin, the fat was chiefly Crisco, while the carbohydrate was composed of a mixture of 2 parts Karo powder and 1 part cane sugar. Cod liver oil U.S.P. which made up 5 per cent by weight of each diet contributed part of the dietary fat and vitamins. Brewers yeast 1 gm. (wet) given on alternate days provided further vitamins. Food and water were always available to the rats.

Technical Procedures: The methods used for collecting clinical and chemical data have been previously described.³ Moribund animals were generally etherized in order to obtain well preserved tissues. Organs were fixed in Zenker's solution and in 10 per cent neutral formalin. Paraffin sections were stained with eosin and methylene blue, and in addition, Mallory's aniline blue stain and McGregor's⁶ modification of the Mallory-Heidenhain method were applied to sections of all kidneys. The scharlach R method was used on frozen sections to show fatty changes. Other staining techniques were occasionally employed.

EXPERIMENTAL

Clinical Course of Nephrotoxic Nephritis in Rats Maintained on Different Diets

The course of the experimental nephritis was similar during the first month in all the rats irrespective of the type of diet that was fed.† Severe albuminuria with cylindruria appeared and per-

* Harris product.

† Data presented in abstract, Farr, L. E., and Smadel, J. E., *Proc. Soc. Exper. Biol. & Med.*, 1937, **36**, 472-473.

sisted. Anasarca was present for a variable number of days. Plasma protein values, which were low during the edematous phase, had generally returned to normal 1 month after the injections of nephrotoxin. On the other hand, blood urea nitrogen values and urea clearance determinations remained within normal limits during this early period, except in a few rats which are not included in this report because they succumbed to the acute disease.

The clinical findings in the group of nephritic rats fed Diet L (low protein-high carbohydrate) diverged markedly during the 2nd month from those of the animals fed the other two diets. Seventeen of the 19 animals that received this diet showed either a marked diminution, or, in a few instances, a complete disappearance of urinary abnormalities. Four of these rats were sacrificed at the end of the 3rd month for histological study. Two animals which had excreted urine of a consistently abnormal character died suddenly in the 5th month of the disease without antecedent renal failure. Eight and one-half months after nephritis had been induced, none of the 13 survivors had elevated blood urea or abnormal urea clearance values; moreover, only 1 showed moderate amounts of protein and casts in the urine, 4 had normal urines and the remainder occasionally put forth traces of urinary protein or a few casts. Two of the rats, which had apparently recovered, and 1 of those with slight evidences of nephritis were sacrificed at this time, while 5 of the 10 surviving rats were changed from Diet L to Diet H (high protein-low carbohydrate). During the final 2 months of observation it became apparent that the change to Diet H had had an adverse effect on the diseased kidneys. The single animal in this subgroup with normal urine at 8½ months did not suffer a relapse, but the other rats with low grade or latent nephritis showed an increase in albumin and casts sufficient to warrant a diagnosis of mild or moderate kidney irritation. Exacerbations of this sort did not occur in the rats maintained on the original Diet L.

Every animal fed Diet H (high protein-low carbohydrate) after receiving antikidney serum continued to have marked albuminuria and cylindruria until it died or was sacrificed. Four members of this group were sacrificed 3 months after injection. Only 2 of the 15 remaining rats survived until the experiment was terminated

10½ months after induction of nephritis; moreover, both of these animals were in a terminal phase of chronic progressive nephritis. Among 13 rats dying with renal failure, the average time of survival after injection was 6 months.

The course of the disease in rats fed on Diet B (basal) was similar to that observed in earlier experiments when the animals were maintained on a varied stock laboratory diet.³ One of the 15 rats in this group recovered clinically during the 2nd month, while 6 showed abnormal urinary contents throughout the experiment but did not develop renal failure. Eight of the 15 rats died of kidney insufficiency; their average time of survival was 5½ months.

Smaller groups of normal rats fed on Diets L and H for several months were sacrificed for histological study. These animals excreted normal urine throughout the period.

HISTOPATHOLOGICAL OBSERVATIONS

The characteristic lesions in rats with acute nephrotoxic nephritis and the chronic nephritis that follows the acute injury have been described.² Both glomeruli and tubules are affected in the acute disease. The tubular epithelium shows necrobiotic changes, principally hyaline droplet degeneration, while the outstanding glomerular lesion consists of swelling of the glomerular capillary basement membranes.

Renal Lesions in Rats Fed Diet B

The nephritic animals maintained on this normal basal diet not only followed the same general clinical course as those kept on a stock diet in previous experiments, but also had essentially the same type of kidney damage; hence, only a summary need be given here. Rats that died of progressive nephritis from 68 to 237 days after receiving nephrotoxin had enlarged kidneys with granular surfaces, and many cystic dilated tubules filled with coagulated material were visible macroscopically throughout the cut sections, especially in the corticomedullary region. These changes were most marked in animals with long-standing disease. Microscopic examination showed that practically all the glomeruli were abnormal: those least affected had distorted tufts with irregularly thickened glomerular capillary basement membranes;

in others the epithelium of Bowman's capsule was proliferated and the capsular membrane was thickened, and there were also glomeruli with varying amounts of connective tissue replacement. Glomerular changes of differing degree were always demonstrable in each section, although the number of severely scarred glomeruli was increased in rats with a more chronic disease. Severe damage was found in tubular structures on histological study; extensive dilatation with large hyaline casts, atrophy in areas of interstitial scarring, and epithelial hyperplasia of the remaining functioning units were all observed. The epithelium of the functioning tubules showed various grades of degeneration, including necrosis, in animals killed when moribund. Vascular lesions as well as perivascular and interstitial infiltrations of cells were characteristically present. Rats in this group, which survived the period of the experiment without developing renal failure, even though showing evidence of chronic nephritis until sacrificed, had the same types of kidney lesions, but many functioning renal units remained.

The single animal in this group in which the nephritic process diminished markedly during the 2nd month, and thereafter became negligible, had kidneys with surfaces that were essentially smooth. Moreover, on microscopic examination most of the structural units appeared normal. A rare dilated tubule filled with a hyaline cast, and an occasional small, completely scarred glomerulus could be found. However, fair numbers of glomeruli which appeared capable of functioning, had slight abnormalities in structure. Changes such as irregular thickening of the capillary basement membrane, local dilatation of capillary loops, thickening of capsular basement membrane and, rarely, small crescents were encountered.

The typical macroscopic appearance of a kidney from a rat with progressive nephritis in this diet group is illustrated in Figure 2. Figures 5 and 7 depict characteristic microscopic lesions seen in such kidneys.

Renal Lesions in Rats Fed Diet H

Young normal and nephritic rats thrived when fed the high protein-low carbohydrate diet H; they gained weight faster and grew larger than rats kept on the basal diet B. A normal male

animal weighing 90 to 100 gm. when placed on Diet H usually doubled its weight in 2½ months. The kidneys of such an un-injected control rat weighed 1.3 to 1.4 gm. each and, except for slight hypertrophy, were normal on inspection and on microscopic examination.

In general, similar kidney lesions were found in rats with chronic progressive nephritis maintained on either Diet B or H; certain differences, however, usually made it possible to distinguish between members of the two groups on macroscopic evidence alone. The kidneys of rats in the Diet H group were usually smaller. Thus, the average weight of the left kidneys of 7 rats dying in Group B was 1.6 gm., while the average of 13 rats dying in Group H was 1.2 gm. The cortical surface was more coarsely granular in Group H. Finally, the cut sections of kidneys from rats in Group B contained many more cystic tubules filled with hyaline material than did kidneys of animals in Group H. These differences were more pronounced in rats which succumbed 5 to 7 months after injection but are clearly depicted in the two kidneys obtained at the end of the experiment and illustrated in Figures 2 and 3.

All gradations of glomerular change described in rats of Group B were observed in animals fed the high protein-low carbohydrate diet; however, the number of completely or extensively scarred glomeruli was always greater in the latter group of animals for any given duration of the nephritis.

A more striking difference between the two groups was found in the varying proportions of the several types of tubular lesions. Cystic tubules filled with hyaline material, observed in such abundance throughout the cortex and medulla of rats with progressive nephritis in Group B, were less frequent and smaller in rats of Group H. On the other hand, necrobiotic changes in the tubular epithelium were more striking in sections from Group H; moreover, degeneration was observed in cells of the proximal segment of tubules in animals of Group H which were killed (3 months after nephritis had been induced) before renal failure was imminent. Widespread interstitial fibrosis encompassing destroyed tubules was marked in the kidneys of the rats in Group H and was observed as early as 3 months after injection of antikidney serum. In addition, hypertrophy and hyperplasia of the epithelium

of remaining functioning tubules were also significantly greater in rats of Group H. Urochrome pigment was conspicuous in the tubular epithelium of rats with terminal anemia irrespective of the diet group. Vascular lesions and perivascular cellular infiltrations occurred with equal frequency and intensity in kidneys of rats that died in the two groups.

In general, glomeruli and tubules which had been damaged by nephrotoxin apparently were unable to undergo repair when the rats were fed the high protein-low carbohydrate diet. Instead, progressive destruction of kidney substance, with connective tissue replacement, proceeded at a more relentless pace than in rats fed the basal diet. Macroscopic and microscopic changes characteristically found in nephritic rats which died while being fed the H Diet are represented in Figures 3, 6 and 8.

Renal Lesions in Rats Fed Diet L

Diet L was sufficiently well balanced to sustain life and to permit a retarded growth of rats, but various abnormalities were observed in animals maintained on it. For example, both untreated and nephritic rats fed Diet L failed to attain normal weight or stature; their fur continued to be soft and short even after the adult state had been reached; and finally, renal and hepatic lesions were consistently observed. It is necessary to present a description of the characteristic pathological changes found in young normal rats maintained on this diet for several months before attempting to evaluate the lesions attributable to nephrotoxin.

A young female rat (PN-13), weight 66 gm., was fed a stock diet for 2 weeks while repeated urine specimens were obtained; no urinary albumin, blood or casts were demonstrable. The animal was then transferred to Diet L. During the next 2½ months this animal gained from 100 to 130 gm. and continued to excrete normal urine.

The kidney, examined at the end of the period, weighed 0.6 gm. and had scattered shallow depressed scars on the cortical surface. On cut section, the corticomedullary region showed numerous, slightly raised yellow streaks arranged in parallel lines radial to the pelvis (see Fig. 11).

The principal changes found on microscopic examination of this kidney occurred in the areas represented macroscopically by

the yellow streaks. Groups of moderately dilated tubules were lined by low cuboidal epithelium, often so thinned as to resemble the connective tissue cells which lay outside the thickened and hyalinized tubular basement membrane. The lumens of dilated tubules generally contained refractile material, sometimes arranged in concentric rings, which stained very faintly with the acid dyes and failed to stain differentially with scharlach R or iodine. Remnants of a brush border could still be identified in certain of the affected proximal convoluted tubules. In the same area, atrophic tubules lined by epithelium with basophilic cytoplasm and with collapsed lumens were present. Their basement membranes were also hyalinized and surrounded by scar tissue. Less drastic changes were demonstrable in other proximal tubules of the corticomedullary area. In some, only a single epithelial cell was degenerated or necrotic, while in others many were affected and cast off cells lay loose in the lumen. Fat was often demonstrated in such damaged cells by scharlach R stains. Regeneration of epithelium was observed and occasionally resulted in an irregular stratification of cells protruding into the lumen. Thickening of the tubular basement membrane or increase in interstitial connective tissue was not conspicuous here. Figure 12 illustrates the microscopic findings in the corticomedullary region of this kidney.

Most of the glomeruli appeared normal but a few had changes in the tuft and occasionally extensive scarring was observed. Damaged glomeruli were usually found near diseased tubules and occasionally could be shown to connect with one of them. Vascular lesions were not found, and interstitial collections of lymphocytes, when present, were small and limited to the corticomedullary scars.

An extreme fatty change was present throughout the liver. The least affected parenchymatous cells, about the portal spaces, contained fat globules, while in the central portion of the lobules the cytoplasm of individual cells stained a homogeneous red with scharlach R.

The renal and hepatic injury which occurred in all young rats fed Diet L for several months varied in degree. Rat PN-13, described above, was one of the most severely affected of the group. On the other hand, Rat PN-7, which seemed to recover

from nephrotoxic nephritis, had a minimal amount of the kidney damage attributable to the low protein-high carbohydrate diet when sacrificed 3 months after injection of antikidney serum.

A young male rat (PN-7) weighed 55 gm. when nephrotoxin was administered and Diet L was started. The acute nephritis began to subside in several weeks and the urine was normal 4 weeks after nephritis had been induced. The animal weighed 94 gm. when sacrificed 3 months after injection. The kidneys, which weighed 0.5 gm. each, were apparently normal on macroscopic examination but showed glomerular lesions throughout the microscopic section. Practically all of the tuft capillaries contained blood but their walls were significantly thickened by an increased width of the basement membranes. About 20 per cent of the tufts were lobulated or otherwise distorted, and approximately 10 per cent of them were adherent to the capsular epithelium in one or more places; nevertheless, well developed crescents were not seen. A few tufts contained scattered cells, apparently epithelial in origin, with large, bright eosin-staining granules in their cytoplasm. Scattered small areas in the corticomedullary region contained dilated tubules, some of which were filled with laminated pale staining material similar to that seen in Rat PN-13; a few atrophic tubules were also present in these areas. An occasional atrophic tubular structure in the cortex was seemingly related to the pathological areas at the junction of cortex and medulla. In addition, tubules containing hyaline casts were occasionally encountered. Vascular lesions and interstitial cellular infiltrations were not observed.

The liver was the seat of mild fatty change. Microscopic sections of the heart were normal.

The kidneys of rats fed Diet L for 8 to 10½ months after nephritis had been induced with nephrotoxin were similar to those of members of the group sacrificed at 3 months. The lesions attributed to the diet were, however, less obvious at 10½ months. The kidneys of only a few of the animals had macroscopic yellow streaks in the corticomedullary region, which on microscopic examination appeared to be dilated tubules filled with poorly staining material. Narrow bands of old connective tissue arranged radially were present in this portion of all the kidneys; these were often conspicuous but sometimes appeared only once or twice in the

entire section. Five to 10 per cent of the glomeruli in the kidneys from older animals were represented by small contracted scars; in addition a few had moderate hyperplasia of the epithelium of Bowman's capsule. The majority of glomeruli appeared to be functioning but showed abnormalities of the tufts such as described in Rat PN-7. The capillaries of the tufts were often dilated in places and contained puddles of blood; these frequently occurred at the border of the tuft. Scattered atrophic tubules were present throughout the cortex. Vascular lesions and interstitial cellular infiltrations were not present in the kidneys of the group of older rats, nor were generalized vascular changes observed.

The livers of the animals sacrificed at the later date showed about as much fatty change as did those examined earlier; certain of the rats also had cirrhosis.

There was no consistent histological difference detected between the kidneys of rats maintained on Diet L throughout the experiments and those of rats transferred from Diet L to Diet H 8½ months after injection of antikidney serum. Fatty change in the livers of rats in this last group, however, was slight when present.

Two rats, in Diet Group L, failed to make a clinical recovery during the 2nd month and died 134 and 139 days, respectively, after nephritis had been induced. Neither of these animals had a significant reduction of kidney function at any time, but their urine contained 1 to 3 gm. of protein per 100 cc. and numerous casts until the end. Kidney lesions in these 2 rats were more extensive than in other members of the group but hardly severe enough to account for the early death. Changes in the tufts, crescent formation, complete scarring of glomeruli, numerous dilated tubules filled with hyaline casts and degenerative changes in tubular epithelium suggested the picture seen in nephritic rats fed Diet B. Both livers were heavily laden with fat. The death of these 2 animals probably depended on the combined effect of severe chronic nephritis and poor diet.

These histopathological observations agree with clinical data and indicate that rats with severe acute nephrotoxic nephritis tend to recover when maintained on a low protein-high carbohydrate diet. The tubular changes characteristic of the acute nephritis appear to be reversible for the most part when circumstances are propitious. Glomerular lesions, on the contrary, do not disappear;

but notwithstanding the presence of residual changes, most of the glomeruli continue to function.

Hematuria in Rats with Chronic and Latent Nephritis

Pearce⁷ held that hematuria was not a result of injury with nephrotoxin but depended on other factors in antikidney serum. We have also emphasized this point in reference to the acute syndrome induced by nephrotoxin.² Present observations offer no reason for changing our views on hematuria in the acute nephritis but do indicate that renal bleeding may occur in rats after the acute phase of renal injury has passed.

Hematuria, observed in 7 rats, was generally of the intermittent type and when it occurred was obvious macroscopically. Two rats in Group L showed red cells in the urine for the first time after they had been fed the low protein-high carbohydrate diet for 9 and 10 months, respectively; moreover, there had been neither casts nor more than a trace of albumin for months before the appearance of blood in the urine. Onset of hematuria in the 4 rats of Group B occurred in the 3rd (2 animals), 4th and 7th months and, in a single animal of Group H, hemorrhage was detected during the 7th month on one occasion.

Hematuria in these animals seemed to have no adverse prognostic significance for all of the rats lived throughout the experiment. No evidence of an infectious process which might have accounted for the bleeding was found in any of the kidneys, and cultures of renal tissue made at the time of autopsy were bacteriologically sterile. Dilated capillaries near the margins of diseased glomerular tufts were observed in rats of all three diet groups, and rupture of such diseased capillaries seems a likely explanation for the intermittent hematuria.

Generalized Vascular Lesions in Rats with Chronic Progressive Nephritis

Thickening and hyalinization of the walls of small arteries and of arterioles occurring in various organs of rats which succumbed to chronic progressive nephritis were described and illustrated in an earlier report.² Additional vascular changes, such as the presence of fat or of calcium in the media and a reduplication of the internal elastic membrane of coronary arteries, and perivascular

TABLE I
Extrarenal Vascular Lesions in Rats that Succumbed to Nephritis

Diet (protein)	Number of rats	Average time of survival	Heart			Pancreas			Brain			Intestine			Liver			Testicle		
			A	C	T	A	C	T	A	C	T	A	C	T	A	C	T	A	C	T
L (low)	2*	4½ mo.			0/2			0/0						0/0			0/2			0/0
B (basal)	7	5½ mo.	2/7	6/7	7/7	1/4	4/4	4/4	0/5	4/5	4/5	1/1	0/1	0/1			0/7			0/2
H (high)	13	6 mo.	3/13	7/13	9/13	0/9	2/9	2/9	0/12	2/12	2/12	1/1	0/1	1/1	0/11	6/11	6/11	0/7	3/7	3/7

A = Acute lesions
C = Chronic lesions
T = Total number of rats in each group showing lesions of acute or chronic variety or both
The numerator of each fraction indicates the number of rats with lesions; the denominator the number of organs examined.
* Rats did not have renal failure immediately before death but had severe chronic nephritis. Death probably resulted from a combination of nephritis and poor diet.

cellular collections in the pancreas were recorded. The occurrence of chronic generalized vascular lesions has also been noted in the present group of rats which died after long-standing nephritis. Their distribution is enumerated in Table I.

Fibrous myocarditis and encephalomalacia which apparently depended on these chronic vascular lesions were found in 13 of 22 rats with chronic nephritis terminating in death; myocardial scarring occurred 5 times in Group B and 3 times in Group H, while encephalomalacia was noted in 3 members of the former group and in 2 of the latter.

In addition to the chronic extrarenal vascular lesions, other changes of a more acute nature were observed, especially in the heart and intestine. These consisted of proliferation and swelling of endothelium, fibrinoid degeneration in the muscle coat, and pyknosis of muscle nuclei. Table I records the frequency of acute vascular lesions such as are illustrated in Figures 13 and 17.

Vessel changes in the heart, brain and pancreas, as well as myocarditis, were found in 1 of the animals of Group B which survived to the end of the experiment; however, its renal function had been depressed for a month before it was sacrificed. Renal insufficiency was present in both rats of Group H when they were sacrificed 10½ months after injection of nephrotoxin. One had vascular lesions in the heart, brain and testicle and also myocarditis, the other had only coronary abnormalities. None of the rats in Group L which were sacrificed during the experiment, or at its end, had generalized vascular lesions.

Acute Myocarditis and Enteritis in Rats with Renal Failure

An acute focal necrosis of cardiac muscle fibers was frequently seen in rats dying with chronic renal insufficiency. These lesions could usually be identified macroscopically as yellow flecks beneath the epicardium and endocardium. On microscopic examination some groups of degenerated cardiac muscle cells were represented by necrotic débris while others still retained their general outline. The cellular reaction in and about such areas was in the majority of cases mononuclear in type but occasionally consisted of polymorphonuclears or, rarely, was entirely absent. In certain areas cellular collections surrounded muscle fibers which retained their striations. Large mononuclear cells were the prin-

cial elements in these collections, but an occasional multinucleated cell was seen. Acute lesions, when present, were usually numerous; one section contained nine foci. The myocardium pictured in Figure 16 contained many affected areas. Acute myocarditis occurred in 5 of the 7 rats in Group B and in 9 of the 13 animals in Group H that died with progressive nephritis. Three of the former animals and 2 of the latter had, in addition, old scarring of the myocardium.

Fibrinoid swelling of the ground substance was found in 4 hearts which contained other acute lesions; 2 rats each were involved in Groups B and H. An area of intense fibrinoid degeneration is shown in the left lower portion of Figure 16, while a milder type of reaction is discernible in the area of focal myocarditis illustrated in Figure 15.

Acute myocardial lesions occurred in 14 of the 20 rats with progressive nephritis and protracted renal insufficiency terminating in death. On the contrary, this type of cardiac involvement was not observed in rats of Groups B and H that survived throughout the experiment even though some of these animals showed lowered urea clearance values during the later months of observation; moreover no such lesions were detected in any rat of Group L. McJunkin *et al.*⁸ described similar foci of acute myocardial necrosis in nephrectomized rats which were injected with large amounts of phosphate. Those authors do not, however, emphasize the occurrence of swelling and proliferation of endothelium in small coronary arteries and capillaries and these changes, which were usually prominent in our rats with acute myocarditis, may have played some part in the formation of foci of necrosis.

Typhlitis was a contributing cause of death in 2 animals with renal failure; 1 in Group B died on the 198th day and the other in Group H survived 136 days after nephritis was induced. In both instances the cecum showed macroscopic subserosal ecchymoses, edema and hemorrhage of the wall, pseudomembranous exudate covering the mucosa, and blood in the lumen. Figure 14 illustrates such a lesion. Vascular changes of the acute variety were conspicuous near the border of the involved areas (see Fig. 13) and probably were responsible for the intestinal lesion. Goldblatt⁹ has recently reported that intestinal bleeding is one

of the cardinal signs of the malignant phase of hypertension in dogs. While blood pressure readings were not made in the present experiment, hypertension was previously demonstrated in rats with chronic progressive nephritis of nephrotoxic origin.³

DISCUSSION

The experiments here presented demonstrate that the course of the nephritis in rats which follows injection of antikidney serum is significantly influenced by at least one internal environmental factor, namely, diet. Clinical evidence of renal disease rapidly subsided in almost all young rats placed on a low protein-high carbohydrate diet after severe nephrotoxic nephritis had been induced. Moreover, the acute tubular injury resolved without permanent damage and only moderate residual changes were present in the glomeruli. Renal failure generally occurred within a year, when nephritic rats were maintained on a high protein-low carbohydrate diet, and progressive destruction and scarring of both tubules and glomeruli was observed histologically. More than half the nephritic rats fed a basal diet developed kidney failure and these also had histological lesions indicative of progressive disease. The clinical and pathological features of the nephrotoxic disease were the same in rats nourished with this last diet as in rats fed a mixed stock diet.

While the glomerular lesions found in the fatal cases of nephritis in animals fed either the high protein or basal diets were essentially alike, the tubular changes differed in certain important respects. The marked dilatation of proximal tubular segments in kidneys of Group B rats suggests that the nephron had been rendered functionless through occlusion of the loop or of the distal tubule by scarring or plugging. Degenerative changes were found in the proximal tubular epithelium of kidneys in Group H even when the animals were sacrificed relatively early in the disease; and, in addition, obliterative scarring of proximal tubules was characteristically present in kidneys of the rats that died. It thus appears that one of the most significant effects of diet on the course of nephrotoxic nephritis involves the response of the acutely damaged tubular epithelium. There was a tendency towards recovery from tubular injury when a low protein-high carbohydrate diet was fed, while progressive involvement of the distal tubular

segment was characteristically found when a basal diet was given, and, finally, destruction of the proximal tubule was outstanding in nephritic rats maintained on a high protein-low carbohydrate diet. The studies of Richards and Walker¹⁰ on selective action of different portions of the kidney tubule in frogs and necturi suggest an approach to an explanation of our observations in rats, but the exact factors responsible for the recovery of the nephron or destruction of its proximal or distal segment are not clear. The response of the glomerular capsular epithelium was similar to that observed in the remainder of the nephron. Obliteration of the capsular space by crescent formation occurred relatively infrequently in rats of Group L notwithstanding the fact that residual changes in the glomerular capillary bed were common. In contrast to this, marked proliferation of the epithelium of Bowman's capsule with connective tissue ingrowth into the glomerular tuft was conspicuous in the kidneys of rats in both Groups B and H.

Medlar and Blatherwick¹¹ have commented on the similarity of the renal picture in chronic dietary nephritis observed in rats with unilateral nephrectomy which were maintained on rations rich in animal protein, and the terminal phase of nephrotoxic nephritis in our rats fed a mixed stock diet. The increased susceptibility of male rats to dietary nephritis noted by these workers was less clear-cut in the present experiments; nevertheless 5 of the 7 rats in Group B and both of the animals in Group H that survived the experiment were females.

Generalized vascular changes of both acute and chronic varieties and the visceral lesions that stem from them were encountered only in rats which developed chronic progressive nephritis and renal insufficiency. Diet apparently influenced production of the extrarenal disease only indirectly by way of its effect on the injured kidney. Differences in the frequency of vascular changes were observed in Diet groups B and H, but probably were not significant. Animals that lived throughout the experiment were approximately 380 days old when sacrificed. The failure to find chronic vascular disease or fibrous myocarditis in our surviving animals with normal urea clearance values agrees with the experience of Wilens and Sproul.^{12,13} These authors observed cardiovascular lesions in more than half the members of a large group of senile rats. However they state regarding the heart

that "almost all of the changes described make their appearance late in the second year of life and do not attain their maximum incidence until well into the third year." Furthermore, they observed few vascular lesions before the 700th day of life. It may well be that the process of aging and the concomitant development of chronic cardiovascular disease are merely accelerated in rats with severe nephritis.

It should be emphasized that rats used in these studies were all of a black and white hooded strain designated as Whelan stock. Experiments to be reported by Swift and Smadel have indicated that rats of the Wistar and Evans strains respond somewhat differently to the effect of nephrotoxin and diet.

CONCLUSION

In rats of the so-called Whelan strain the chronic nephritis which follows the administration of antikidney serum can be markedly influenced by isocaloric diets containing different proportions of protein and carbohydrate.

REFERENCES

1. Ehrlich, William E., Wolf, Richard E., and Bartol, George M. Acute experimental glomerular nephritis in rabbits, a correlation of morphological and functional changes. *J. Exper. Med.*, 1938, 67, 769-790.
2. Smadel, Joseph E. Experimental nephritis in rats induced by injection of anti-kidney serum. III. Pathological studies of the acute and chronic disease. *J. Exper. Med.*, 1937, 65, 541-555.
3. Smadel, Joseph E., and Farr, Lee E. Experimental nephritis in rats induced by injection of anti-kidney serum. II. Clinical and functional studies. *J. Exper. Med.*, 1937, 65, 527-540.
4. Chanutin, Alfred. Experimental renal insufficiency produced by partial nephrectomy. III. Diets containing whole dried liver, liver residue and liver extract. *Arch. Int. Med.*, 1934, 54, 720-745.
5. Swift, Homer F., and Smadel, Joseph E. Experimental nephritis in rats induced by injection of anti-kidney serum. IV. Prevention of the injurious effects of nephrotoxin *in vivo* by kidney extract. *J. Exper. Med.*, 1937, 65, 557-564.
6. McGregor, Leone. The finer histology of the normal glomerulus. *Am. J. Path.*, 1929, 5, 545-558.
7. Pearce, R. M. An experimental study of nephrotoxins. *Univ. Penn. M. Bull.*, 1903-04, 16, 217-235.
8. McJunkin, F. A., Tweedy, W. R., and Mencky, W. J. Necrosis of the myocardium induced by the orthophosphates. *Arch. Path.*, 1934, 18, 626-634.

9. Goldblatt, Harry. Studies on experimental hypertension. VII. The production of the malignant phase of hypertension. *J. Exper. Med.*, 1938, 67, 809-826.
10. Richards, A. N., and Walker, Arthur M. Urine formation in the amphibian kidney. *Am. J. M. Sc.*, 1935, 190, 727-746.
11. Medlar, E. M., and Blatherwick, N. R. The pathogenesis of dietary nephritis in the rat. *Am. J. Path.*, 1937, 13, 881-896.
12. Wilens, S. L., and Sproul, E. E. Spontaneous cardiovascular disease in the rat. I. Lesions in the heart. *Am. J. Path.*, 1938, 14, 177-200.
13. Wilens, S. L., and Sproul, E. E. Spontaneous cardiovascular disease in the rat. II. Lesions in the vascular system. *Am. J. Path.*, 1938, 14, 201-216.

DESCRIPTION OF PLATES

PLATE 38

- FIG. 1. Cortical and cut surfaces of kidney from Rat 4 L. Rapid recovery from acute nephrotoxic nephritis occurred on the low protein diet; albuminuria had diminished to a faint trace and casts were absent a month after injection of antikidney serum. Autopsied 10½ months after onset of acute nephritis. Left kidney, weight 1 gm. $\times 3$.
- FIG. 2. Kidney of Rat 8 B in Diet group B. Severe albuminuria and cylindruria continued from the time of induction of acute nephritis until the rat was sacrificed 10½ months later; final urea clearance value was within the lower limits of normal. Kidney weight 2.1 gm. $\times 3$.
- FIG. 3. Kidney of Rat 3 H fed the high protein diet for 10½ months after injection of nephrotoxic serum. Severe nephritis was evident throughout and a depressed urea clearance was observed during the 2 months preceding sacrifice. Kidney weight 1.7 gm. $\times 3$.
- FIG. 4. Rat 4 L. Few abnormalities are detectable in the section of kidney at this magnification. Mallory's aniline blue stain. $\times 35$.
- FIG. 5. Rat 11 B, basal diet group, died with renal failure 7 months after onset of nephritis. Extensive destruction of tubules and distortion of glomeruli are evident, but the most conspicuous abnormality is the markedly dilated tubules filled with hyaline material. Mallory's aniline blue stain. $\times 35$.
- FIG. 6. Rat 5 H, high protein diet, succumbed to kidney insufficiency 5½ months after receiving nephrotoxin. Glomerular damage is similar to that found in Fig. 5. Nests of hypertrophic tubules constitute the bulk of the substance. Extensive areas of interstitial scarring have replaced destroyed tubules. Mallory's aniline blue stain. $\times 35$.

PLATE 39

FIGS. 7 and 8. Higher magnification of the same sections illustrated in Figs. 5 and 6 respectively. $\times 150$.

FIG. 9. A higher magnification of the section shown in Fig. 4. The tubules are not significantly altered and the glomeruli appear capable of functioning. Moderate irregular thickening of the glomerular capillary basement membranes is apparent. Some distortion of the glomerular tuft on the right is to be seen, while slight proliferation of the capsular epithelium and a small capsular adhesion can be observed in the glomerulus on the left. $\times 175$.

FIG. 10. An area of encephalomalacia in the cortex of the parietal lobe of Rat 13 B, basal diet group, which died of renal failure $5\frac{1}{2}$ months after nephritis had been induced. Eosin-methylene blue stain. $\times 70$.

PLATE 40

- FIG. 11. Kidney of Rat PN 13, an uninoculated control animal which was fed the low protein diet for 3 months. The pale streaks in the cortico-medullary region were yellow in the fresh specimen. $\times 4$.
- FIG. 12. Higher magnification of Fig. 11, showing degenerated tubules with casts in the corticomedullary region. Eosin-methylene blue stain. $\times 250$.
- FIG. 13. Section of the cecum of Rat 8 H, which was fed the high protein diet and which died 7 months after the injection of nephrotoxin. An acute necrotizing process is present in all three vessels. These vessels can also be seen at the extreme right in Fig. 14. Eosin-methylene blue stain. $\times 210$.
- FIG. 14. Same as Fig. 13. Hemorrhage and edema in the submucosa with a fibrinous exudate covering the necrotic mucosa are shown. $\times 30$.
- FIG. 15. Section of the heart of Rat 15 B, basal diet group, which succumbed with nephritis in the 7th month. A small area of focal necrosis of muscle fibers and infiltration with mononuclear cells is seen. Eosin-methylene blue stain. $\times 450$.
- FIG. 16. Same as Fig. 15. Multiple focal lesions. An area of fibrinoid change surrounded by muscle fibers showing fresh necrosis is present at the lower left. Areas of necrosis with cellular infiltrations are shown at the lower right and upper portions of the microphotograph. Thickening of vessel walls was also present in other portions of the section. Eosin-methylene blue stain. $\times 175$.
- FIG. 17. Heart of Rat 10 B, basal diet group, which died with nephritis after 68 days. Swelling and vacuolation of cells of the media are prominent in the large vessel on the left. Hyperchromatic enlarged cells are seen in the wall of the small vessel in the lower center as well as swelling of the intimal cells. The vessel at the lower right is completely occluded by intimal proliferation. Eosin-methylene blue stain. $\times 450$.