marked retardation of the E.S.R. A six-hour period in which the blood can stand on the side-room or laboratory bench gives ample time for an accurate estimation of the E.S.R. Advantage can be taken of this by workers in haematology departments who have many samples delivered to them in a short space of time, and also by general practitioners who may wish to take blood during a busy surgery.

The hypothesis that because the E.S.R. at six hours is the same as the immediate E.S.R. of any sample, then at any intervening time the E.S.R. will be similar, is borne out by the small series of tests in experiment 4. The intermittent mixing of the same sample, as shown by experiment 5, has no apparent effect on the E.S.R., which probably slows after about eight hours' storage. It is probably safe to assume that such mixing as a sample might receive in a practitioner's case, car, or pocket will have no effect on the E.S.R. within the six-hour period following withdrawal of the blood. There must, of course, be adequate mixing of the blood sample at the time of setting up the E.S.R.

The use of sequestrene blood thus enables E.S.R.s to be estimated at any time up to six hours after withdrawal of the blood, and extends the usefulness of this clinical test.

Summary

The need to set up E.S.R. estimations immediately after withdrawal of blood samples is obviated by taking the blood into the anticoagulant sequestrene. When this is used there is no significant retardation of the E.S.R. up to six hours after the blood is taken. It does not matter whether the blood stands undisturbed or is mixed at intervals, provided that it is thoroughly mixed before the estimation is made. Storage for longer periods at room temperature is often accompanied by marked retardation of the E.S.R.

Storage of sequestrene blood in the refrigerator for 24 hours does not affect the E.S.R., but storage for 48 hours is usually attended by marked slowing of the E.S.R. It is recommended that sequestrene blood should not be used for E.S.R. estimations if it has been kept in the refrigerator beyond 24 hours.

We thank Dr. E. G. Oastler for his advice in the preparation of this paper.

REFERENCES

- Melville, I. D., and Rifkind, B. M. (1959). *J. clin. Path.*, **12**, 258.
Proescher, F. (1951). *Proc. Soc. exp. Biol. (N.Y.)*, **76**, 619.

A year ago the Council of Europe concluded an agreement whereby the signatory countries undertook to grant each other assistance with regard to the supply of blood and blood products, thereby taking the first steps towards setting up what virtually amounts to a European blood bank. The Council's Committee of Experts on Public Health, at a recent meeting in Strasbourg, examined the possibility of concluding another similar agreement providing for the pooling of test sera of human and other origin for purposes of blood-group determination. A preliminary study of this question was undertaken in July, 1959, in London by a group of doctors who will again meet this March. A draft European agreement on the exchange of test sera will be prepared for the consideration of the Council's Public Health Committee at its next session in June.

Preliminary Communications

Some Experimental and Clinical Data on the Sarcylisin Peptides

Since 1956 the synthesis and study of peptides and amides of sarcylisin have been carried out at the Institute of Experimental and Clinical Oncology, Academy of Medical Sciences of the U.S.S.R., in collaboration with the Institute of Element-organic Compounds, on the presumption that the use of the carrier of chloroethylamino groups may result in the production of compounds with a varying effect upon tumour spectra (Larionov, 1956, 1957, 1958b, 1959a, 1959b).

The first peptides of N-formyl-sarcylisin were synthesized by Knunianz, Kildisheva, and Golubeva (1956). Experimental study of these compounds carried out by Larionov and Sophina (1957) and Larionov (1958a) showed first of all that these are but slightly toxic. Thus the LD₁₀₀ for mice amounts to 750–1,000 mg./kg. Nevertheless, all these compounds showed anti-tumour activity in mice and rats at a dose of 25–60 mg./kg. daily by intraperitoneal injection and of 200–300 mg./kg. by mouth. It was also found that these peptides vary in their action upon the tumour spectrum in mice and rats. In addition, sarcylisin peptides were found to inhibit haemopoiesis to a lesser degree than does sarcylisin itself.

Subsequently, peptides of N-acetyl-sarcylisin have been synthesized (E. N. Shkodinskaya and others). Experiments on animals (Sophina, 1959) confirmed the data obtained with peptides of N-formyl-sarcylisin, with the difference that some of them, such as N-acetyl-sarcylisil-valine ethyl ester ("asaline"), proved effective at smaller doses. Their therapeutic intraperitoneal dose for rats is 7–25 mg./kg. daily.

Each of the N-acetyl peptides likewise has its own "action spectrum" with regard to animal tumours. Thus asaline cures rat sarcoma 45, inhibits growth of sarcoma M-I and 298 by 98%, but sarcoma 37 by only 50%. N-acetyl-sarcylisil-methionine ethyl ester likewise cures sarcoma 45 almost as effectively, but is much less effective with regard to sarcoma M-I and 298 (88% and 30% inhibition), though it is more effective than asaline with regard to sarcoma 37 (88% inhibition).

Clinical tests at the Research Hospital of the Institute of Experimental and Clinical Oncology, under the guidance of N. N. Blokhin, have as yet been carried out only with asaline (19 patients). Daily doses of 1 to 2 g. have been used, but it is not yet certain whether these are optimal doses. Total doses amounted usually to about 20 g.

The Table shows the number of patients, kind of tumour, and the effect of treatment.

The therapeutic effect manifested itself in partial, and less often in complete, regression of the tumours. In two patients with seminoma metastases and in three with myelomatosis relapses were recorded within one to three months. A positive effect was obtained after repeated courses of asaline treatment of myelomatosis. Both patients with far advanced

reticulosarcoma have succumbed despite regression of the tumours. In the patient affected with Ewing's tumour asaline gave a transient effect, and growth of the tumour has recurred even during the administration of the drug.

Asaline was well tolerated by the patients. No depressing effect on haemopoiesis was noted. The

Results of Treatment

Kind of Tumour	No. of Patients	Effect on Tumour	
		Yes	No
Metastases of seminoma of testis	4	3	1
Reticulosarcoma	2	2	—
Ewing bone tumour	2	1	1
Myelomatosis	4	4	—
Plasmocytoma	1	1	—
Cancer of stomach	2	—	2
Squamous-celled cancer of buccal mucosa	1	—	1
Cancer of uterine cervix	1	—	1
Metastases of osteosarcoma	1	—	1
Hodgkin's disease	1	—	1
Total	19	11	8

leucocyte count, though frequently remaining unchanged, was somewhat decreased in some patients. Side-effects have not been noted in the gastro-intestinal tract.

The effect of asaline on certain sarcolysin-sensitive tumours has thus been demonstrated, but clinical data are too scanty as yet to permit a complete comparison of the action spectra of each drug. Nor do they suffice to establish whether asaline has any advantage over sarcolysin. It will be noted, however, in favour of asaline, that this drug does not cause any apparent inhibition of haemopoiesis, so that it may be used in patients with leucopenia.

L. F. LARIONOV.

N. I. PEREVODCHIKOVA.

Institute of Experimental Pathology and Therapy of Cancer, Academy of Medical Sciences of the U.S.S.R., Moscow.

REFERENCES

- Knunianz, I. L., Kildisheva, O., and Golubeva, N. E. (1956). *Izv. Acad. Sci. U.S.S.R., Div. Chem. Sci.*, No. II, 1418.
 Larionov, L. F. (1956). *Brit. J. Cancer*, **10**, 26.
 — (1957). *Acta Un. int. Cancr.*, **13**, 393.
 — (1958a). *La chimiothérapie des cancers et des leucémies*, p. 181. Paris.
 — (1958b). *Vestn. Akad. med. Nauk*, No. 2, p. 8.
 — (1959a). *Acta Un. int. Cancr.*, **15**, 42.
 — (1959b). *Vestn. Akad. med. Nauk*, No. 6, p. 25.
 — and Sophina, Z. P. (1957). *Doklady Acad. Nauk*, **114**, No. 5, p. 1070.
 Sophina, Z. P. (1959). *Proceedings of the Second All-Union Oncological Conference*, p. 773. Leningrad.

"The hardest blows to marriages are given by 'personal defects,' a term which includes things like egoism, financial irresponsibility, immaturity, low intelligence, negligence, badness, etc. Nearly half the cases dealt with by the council come into this category. Then come sex difficulties (31%), health (22%), influence of in-laws (20%), and finally infidelity and incompatibility. Lack of money is one of the less important reasons, accounting for about 6%. At the Marriage Guidance Council they think marriage has a difficult future, especially in England. The independence of women has deprived the father of his authority, and the family has not yet adjusted itself to the new order. 'In Italy,' said Mr. Brayshaw, the general director, 'the throne is still occupied by the father; in the United States it is occupied by the mother, but here there are two thrones, one for the husband and one for the wife, and the equilibrium is very often unstable.'" (*Marriage Guidance*, December, 1959.)

Medical Memoranda

Case of Immune Aplastic Haemolytic Anaemia with Thrombocytopenia

The following case is worth recording in view of the importance of the recognition of the immune mechanism involved and its therapeutic implications.

For some time there has been speculation about the relationship between erythroblastic aplasia and antigen-antibody reactions (Smith, 1949, 1953; Rohr, 1952; Loeb, Moore, and Dubach, 1953; Gasser, 1957). Gasser summarizes the pathogenesis involved by saying that, whereas in the usual haemolytic anaemia the mature erythrocytes are destroyed and the reticulocytes in the blood as well as the erythroblasts in the bone marrow are in excess, on rare occasions the immune antibodies destroy the erythroid series in the bone marrow as well as the mature erythrocytes and reticulocytes in the peripheral blood. This is the so-called immune aplastic haemolytic anaemia, and was seen at certain times in the case described below. If the immune antibodies destroy erythrocytes and reticulocytes but leave the erythropoietic elements of the bone marrow intact, the peripheral blood findings suggest an aplastic anaemia but the marrow shows erythroid hyperplasia. Gasser calls this "immune pseudoaplastic haemolytic anaemia," and it was also seen at certain times in our case.

Bonham-Carter *et al.* (1954) reported the first case of an immune type of aplastic anaemia. This occurred in a 9-year-old boy whose anaemia was associated with severe thrombocytopenia. Splenectomy, by its beneficial effect on the platelet level, was life-saving. At present he is being maintained on a high critical dosage of prednisolone. He has iatrogenic Cushing's syndrome, but if the dose of prednisolone is reduced, haemolysis occurs and there is a rapid fall in haemoglobin. The infant in the following case had an immune type of aplastic anaemia associated with thrombocytopenia similar to that described by Bonham-Carter *et al.* The anaemia and thrombocytopenia could not be controlled by the use of prednisone; splenectomy, however, was of the very greatest value.

CASE REPORT

The patient, a female infant, was born on December 12, 1956, after a normal pregnancy. There was no evidence of ABO blood-group or rhesus incompatibility between mother and child, and the infant was not jaundiced in the neonatal period. There was no family history of anaemia, jaundice, or bleeding tendency; the only sibling is perfectly healthy.

The baby was thought by the mother to be pale in the neonatal period, and at 2 months of age some lesions resembling infantile eczema appeared behind the neck, on the ears, and behind the knees. Shortly afterwards she began having small, frequent nose bleeds; at 3 months purpura appeared for the first time, accompanied by increasing pallor, refusal of feeds, and listlessness.

She was admitted to hospital on June 24, 1957, looking very pale, covered with discrete pin-points of purpura, and with weeping areas of "infantile eczema." There was no enlargement of lymph nodes and the spleen was not felt. Investigations before treatment showed: Hb, 26% (3.85 g.); R.B.C., 1,000,000/c.mm.; W.B.C., 22,000/c.mm.; platelets, 30,000/c.mm., blood group O, Rh positive; bleeding time, more than 10 minutes; Schumm's test for methaemalbumin,