BRITISH MEDICAL JOURNAL

LONDON SATURDAY JANUARY 2 1960

CLINICAL AND METABOLIC STUDIES IN THYROID DISEASE*

BY

E. J. WAYNE, M.D., F.R.C.P.

Regius Professor of Practice of Medicine, University of Glasgow

I have chosen for these two lectures a title which will permit me to cover a very wide field, but which compels me, therefore, to be selective in my approach, and I should like first of all to explain what directed my choice of the topics on which I intend to speak. My interest in thyroid disease was first stimulated soon after I qualified, and since then I have seen and taken part in a steady advance in the diagnosis and treatment of thyroid dysfunction. The direction of my investigations has on several occasions been decided by events in my own particular career, and I hope I may be allowed the pleasure of a few reminiscences which will, I think, make clearer to you the shape of the lectures I am privileged to give to the College.

An important influence on my career was a period spent as an experimental pharmacologist, since this taught me to look statistically at medical problems. Though it is nearly thirty years ago, Professor A. J. Clark, of Edinburgh, was even at that time emphasizing the great variability of biological material, and in particular the variation which the individual animal may show in its response to drugs. He was a pioneer in demonstrating that statistical methods were necessary to deal with this phenomenon, and he and his successor, Professor J. H. Gaddum, extended the scope of these techniques and applied them to human pharmacology.

But my interests were not primarily in the laboratory, and I was soon fortunate enough to obtain an appointment to the Department of Clinical Research at University College Hospital, where I worked under Sir Thomas Lewis. At that time Lewis was formulating his ideas on clinical science. He thought it most profitable to study disease directly under controlled conditions, and to apply in the wards techniques devised by physiologists for the study of normal physiological processes. Though he thought mainly in terms of cardiovascular disease, he would, I am sure, have recognized in some of the studies I shall report the proper tradition of clinical science.

In these lectures I shall describe some new methods of handling clinical evidence. These techniques have a statistical basis, and I have little doubt that my approach to clinical problems stems from my training and experience as a pharmacologist. Some are modifications of the techniques for the clinical trials of drugs which Professor Bradford Hill has pioneered in this country with such striking success.

In my first lecture I shall describe certain aspects of thyrotoxicosis. A few of the ideas are an extension of

those put forward in the Bradshaw Lecture on "The Diagnosis of Thyrotoxicosis" (Wayne, 1954). I shall take the opportunity of indicating where further experience has reinforced or modified my views and shall also describe one or two new diagnostic techniques which have only recently become available. This lecture will also deal with the treatment of thyrotoxicosis, and especially with observations which were designed to make it possible to obtain statistically valid observations from a comparatively small number of cases. In this way we have been able to evaluate in a more precise way than has hitherto been possible the therapeutic effectiveness of various antithyroid drugs in man.

My second lecture deals mainly with the aetiology, diagnosis, and treatment of hypothyroidism and in particular with the relative incidence of the clinical features of the condition, but I shall also discuss the special problem of the diagnosis of Hashimoto's disease or auto-immunizing thyroiditis.

Profound alterations in body metabolism occur in both thyrotoxicosis and hypothyroidism. We have studied the effects of both conditions on total body water, total red-cell mass, total exchangeable electrolytes, and calcium metabolism, and I shall report our conclusions in brief and summary form.

In the five years since I went to Glasgow, where much of my time is occupied in teaching and administrative duties, it would not have been possible for me, without assistance, to carry out this work, and I am much indebted to all those who have collaborated with me, and especially to Dr. J. Crooks and Dr. I. P. C. Murray, who have taken a large and active share in these studies, and to Dr. R. A. Robb for his advice on the statistical aspects. I would also like to mention others who have contributed to the work I shall describe: Dr. M. M. Bluhm, Dr. James Buchanan, Dr. Watson Buchanan, Dr. F. P. Muldowney, and Dr. B. E. C. Nordin.

A Statistical Method Applied to the Clinical Diagnosis of Thyrotoxicosis

Though the diagnosis of thyrotoxicosis is often easy, there are many cases which give rise to uncertainty. This is especially so when the clinical picture is incomplete, or when atypical features are present, and in these circumstances even experienced clinicians may differ in their conclusions. Sometimes the same observer may alter his opinion on the same case on consecutive days. This state of affairs has always puzzled me, and the reason is not immediately obvious. It seems to depend on the nature of the mental processes involved in arriving at a diagnosis. In making his clinical

*The first of two Lumleian Lectures delivered before the Royal College of Physicians of London on April 14 and 16, 1959.

assessment a physician must first obtain a reliable history and elicit accurately the appropriate physical signs. At this stage we encounter "observer variation," since there is rarely complete agreement even among a group of experienced clinicians. Next, the physician must decide upon the relative importance he should attach to the diagnostically significant clinical features of the case. In so doing he falls back on his own experience or on that of his teachers. At this point further differences of opinion arise, due partly to the different degree of significance attached to certain features by different clinicians, and partly to the influence of a fortuitous run of positive or negative findings occurring towards the end of the clinical examination. It is because of these difficulties in making a diagnosis on clinical grounds alone that so many tests of thyroid function have been devised and are so widely used.

By using some of the principles of the statistical method known as discriminant analysis we have found it possible to increase the accuracy of the clinical diagnosis and also to study the effects of "observer error." The method consists in allocating to each clinical feature a positive or negative score, the value of which is based on analysis of the relative frequency of symptoms and signs in the disease. In this way a total score, or clinical diagnostic index, can be obtained in each case. There is a wide difference between these scores in frankly thyrotoxic patients and in normal persons, and in practice the index has proved helpful in distinguishing between toxic and non-toxic patients in cases presenting diagnostic difficulty. It also provides a numerical estimate of the degree of severity of the disease, which can be correlated with other indices of thyroid function.

We first studied 83 unquestionably thyrotoxic patients and 99 unquestionably non-toxic subjects, including in this group not only healthy persons but also patients with simple goitres, anxiety states, and post-menopausal symptoms. This group is termed "definite" in the tables and discussion. In each subject the presence or absence of the clinical features shown in Table I was recorded. These signs and symptoms were chosen because they had previously been shown by a clinical survey to differ in their incidence in thyrotoxic and in

normal subjects (Wayne, 1954). In order to reduce the effects of "observer variation" the method of history-taking and the procedure for the physical examination were rigid. We have described our criteria elsewhere (Crooks *et al.*, 1959).

The Clinical Index

The clinical features recorded in the cases which had given rise to no clinical diagnostic difficulty were weighted by allocating a score to each (Table I). The

TABLE I.—*Thyrotoxicosis—Clinical Diagnostic Index. Weighting Factors Allocated to the Symptoms and Signs of Thyrotoxicosis*

Symptoms of Recent Onset and/or Increased Severity	Present		Signs	Present Score	Absent Score
	Score	Score			
Dyspnoea on effort	+1		Palpable thyroid	+3	-3
Palpitations	+2		Bruit over thyroid	+2	-2
Tiredness	+2		Exophthalmos	+2	
Preference for heat (irrespective of duration)		-5	Lid retraction	+2	
Preference for cold	+5		Lid lag	+1	
Indifferent to temperature	0		Hyperkinetic movements	+4	-2
Excessive sweating	+3		Fine finger tremor	+1	
Nervousness	+2		Hands: Hot	+2	-2
Appetite increased	+3		Moist	+1	-1
Weight increased		-3	Casual pulse rate:		
Weight decreased		-3	Auricular fibrillation	+4	
	+3		Regular rates:		
			<80 per minute		-3
			80 to 90 per minute	0	
			>90 per minute	+3	

positive or negative values of these scores were initially allocated on the basis of the relative diagnostic significance of each symptom and sign, as found by Wayne (1954) or, in a few instances, by Lerman (1948). These scores were then modified so as to diminish the effects of "observer variation." The total scores, which we call clinical diagnostic indices, were then calculated. These indices produced a good separation between non-toxic and toxic subjects; but the weighting factors for the individual clinical features were further modified to produce the widest possible separation between the two groups without reintroducing excessive "observer variation" effects. The indices were then recalculated. Fig. 1 shows the range of values obtained and the sharp separation between the toxic and non-toxic groups, so

FIG. 1.—Clinical diagnostic index applied to cases in the definite group.

that an index of +20 or over indicated toxicity and one of +10 or lower that the patient was non-toxic.

We next studied a group of cases which were not obviously toxic or non-toxic. This group, which is subsequently called "doubtful," consisted of 118 cases, each of which had presented some diagnostic difficulty to one or more hospital physicians. All had been referred for radioactive iodine studies, and the final diagnosis was made only after prolonged observation, including the response to treatment. We were, therefore, confident of our final conclusions. At the patient's first visit the presence or absence of the symptoms and signs listed in Table I was recorded. Clinical diagnostic indices were calculated only when the final diagnosis was agreed after full investigation and the results of therapy were known. The distribution of the indices of the 118 patients in this group is shown in Fig. 2. A good division was still found between cases finally shown to be non-toxic and those shown to be toxic, 88% of their indices lying within the non-toxic and toxic ranges respectively. Thirteen subjects had indices lying between +11 and +19. We have called this the equivocal range. I shall mention these cases again later.

It thus seemed that in our hands the index was a reliable guide to the clinical diagnosis, but we could not be sure that it would be so when it was used by others. Fletcher (1952) has shown that competent physicians, investigating cases of emphysema, report widely different physical signs, and has emphasized the existence of "observer variation." We tried to limit the influence of this factor by formulating rigid definitions for the clinical features of thyrotoxicosis included in our index. But if it was to be generally applicable we knew we should have to investigate the results obtained by other clinicians. For this purpose we chose nine patients whose thyroid function had been difficult to assess clinically and nine observers co-operated with us. Analysis of variance of the results showed no significant difference in the score of seven of the nine observers. A medical student who had seen several cases of thyroid disease and who had received special instruction in the use of the scoring system was among the seven. The remaining two, who were both newcomers to the unit and had had no special experience of thyroid disease, recorded systematically lower scores. Nine further cases were used to make a comparison between the indices obtained by the physician who designed the score sheet and those allotted by a newcomer to the unit who had a wide experience of thyroid disease. There was no significant difference in their scores. These results suggest that no statistically significant difference will be obtained by observers using the diagnostic index provided they have some experience of thyroid disorders. Inexperienced observers tend to score low.

We have compared the relative diagnostic accuracy of the index with laboratory tests of thyroid function. Radioactive iodine studies were made on 202, and basal metabolic rate estimations were carried out on 188 of the 300 cases represented in Figs. 1 and 2. There was no statistically significant difference between the accuracy of the index and either radioactive iodine tests or basal

metabolic rate estimations. In making this comparison, all the indices in the equivocal range between +11 and +19 were regarded as incorrect. We next applied the index to a further 50 patients, referred to us because they had given rise to diagnostic difficulty. We also asked three colleagues at three other hospitals in Scotland to use the index at their routine clinics and to compare the results with their final diagnosis. The total number of correct diagnoses made in this way was 149 out of 171 cases investigated—that is, 85%. Of the remaining 22 cases which were not correctly classified as toxic or non-toxic it is important to note that the diagnoses were completely misleading in only five subjects, or 3%. The remaining 17 all gave indices lying in the equivocal range, thus indicating the need for further investigation.

The cases which fall in the equivocal range are of special interest. In this group, when the final diagnosis

FIG. 2.—Clinical diagnostic index applied to cases in the doubtful group.

is known, an examination of the individual features making up the index usually shows why difficulty has arisen and throws some light on the general problem of making a clinical diagnosis in thyrotoxicosis. Thus the non-toxic subjects in this group usually have severe anxiety states and have a high symptom score. In itself this is not usually sufficient to place them in the equivocal range unless they have in addition one or two heavily weighted signs such as goitre or tachycardia; several patients have had a previous thyroidectomy and the eye signs have persisted. The toxic subjects with scores falling in the equivocal range usually lack some diagnostic features which score highly; for example, a male patient who has neither goitre, bruit, nor hyperkinetic movements.

The clinical index can be used as a measure of the degree of severity of thyrotoxicosis, since it indicates the number of target-tissue effects, and, as might be expected, it correlates fairly well ($r=0.33$) with the basal metabolic rate (Fig. 3); it also correlates equally well ($r=0.36$) with both the four-hour uptake of radioactive iodine and the 48-hour protein-bound plasma radioactive iodine.

We believe that the clinical diagnostic index is a reliable and simple aid to diagnosis and is of value in day-to-day practice. It should be pointed out that our final decision on whether a patient was thyrotoxic or not was based on the response to therapy, and it follows that we define thyrotoxicosis as a condition in which antithyroid therapy produces a remission in the clinical features of the disease.

Lastly, I would emphasize that all clinical diagnosis involves the subconscious application of the principles underlying the method I have described. The clinician sifts and evaluates multiple symptoms and signs, and tries to select the clinical syndrome which best accounts for his findings. He attaches greater or less significance to the clinical features in the light of his past experience, and applies the technique of multiple correlation

much more consistent and reliable readings are obtained. We have modified their techniques and we now carry out duplicate readings on at least two consecutive days. The patient stays in hospital overnight and is given full doses of butobarbitone. The lowest of the four values is adopted. With this method we have compared the diagnostic accuracy of the Robertson and Reid and the Aub and Du Bois standards in 209 subjects (Crooks *et al.*, 1958).

There is no difference in clinically obvious cases, but in doubtful cases which were subsequently proved to be thyrotoxic three-quarters were correctly placed by the former standards, compared with only half (taking the upper limit as +10%) or a quarter (taking the upper limit as +20%) on the Aub and Du Bois scale. Ten definitely thyrotoxic patients who had "normal" B.M.R.s by both standards were reinvestigated after treatment. In all of them the rate fell. In some this was probably due to the occurrence of the disease in patients whose B.M.R.s in health were much lower than the established mean value (Robertson, 1934). In at least half of the cases it was more probably due to obesity. Depot fat has a very low rate of oxygen utilization (Krebs and Johnson, 1948), so that obesity produces an increase in surface area without a proportional change in oxygen consumption. The incorporation of lean body mass in the reference standard has been shown by Bernstein *et al.* (1956) to eliminate this fallacy. In practice this finding suggests that a normal B.M.R. in a patient in whom there are grounds for suspecting thyrotoxicosis should not preclude the diagnosis if the weight approximates to height and age and there is a history of loss of weight.

FIG. 3.—Clinical diagnostic index used as a measure of severity and correlated with the basal metabolic rate. The regression line of the index on the B.M.R. in toxic cases is shown.

subconsciously in a non-quantitative manner. The clinical index applies this principle at the conscious level and makes a diagnosis less individual and more objective.

Special Tests for Thyrotoxicosis

In my Bradshaw Lecture I reviewed the laboratory methods which have been used as diagnostic aids in thyrotoxicosis. Further experience has made me revise some of my views on their relative value and accuracy, and I shall give the reasons for this change of opinion and describe some new techniques which have been introduced.

Basal Metabolic Rate

This is the standard laboratory investigation used by most clinicians when they wish to confirm or refute a diagnosis of thyrotoxicosis. The basal metabolic rate (B.M.R.) is raised in a variety of other conditions—for example, in leukaemia, heart failure, and phaeochromocytoma. A complete list of diseases which may produce hypermetabolism and therefore misleading results is formidable (Bauer, 1956). In practice, however, the presence of these conditions is usually obvious and the estimation is not carried out. A thyrotoxic patient, however, may have a B.M.R. lying within the normal range, and Skanse (1949) refers to numerous studies which report the occurrence of normal rates in 10 to 20% of cases of thyrotoxicosis. At Sheffield we (Goodwin *et al.*, 1951) found that 35% of our cases had rates lying below +20% using the standards of Aub and Du Bois (1917). Further experience has modified my views on the degree of diagnostic inaccuracy in B.M.R. estimations. If the technique and standards advocated by Robertson and Reid (1952) are adopted,

Sleeping Pulse Rate

Though this is not strictly a laboratory test, I should like to discuss it here, since we always record the pulse rate while the patient is asleep under the influence of the hypnotic given before a B.M.R. estimation. In addition we have examined a larger number of readings taken by night nurses who had been carefully instructed on the technique they were to use. Details are given elsewhere (Crooks and Murray, 1958). The important findings are that sleeping pulse rates of more than 80 per minute make the diagnosis of thyrotoxicosis highly probable. There is, however, no significant correlation between the rate and the degree of clinical severity of the disease. Unfortunately, about a half of all the thyrotoxic patients studied had sleeping pulse rates of 80 per minute or lower, and this figure rises to two-thirds if diagnostically difficult cases only are considered. In 4 out of 138 cases of thyrotoxicosis sleeping pulse rates lower than 65 were recorded.

Chemical Estimation of Serum Protein-bound Iodine

The estimation of the amount of ^{127}I bound to protein is generally regarded as an accurate index of the amount of circulating thyroid hormone. Several workers in the United States of America have produced evidence that levels outside the normal range are the most reliable laboratory evidence of abnormal thyroid function. The subject is well reviewed by Rapport and Curtis (1950). We experienced serious technical difficulties with the method of Barker *et al.* (1951), but Macgregor and Farrell (1958) have introduced us to a modification of the method of Zak *et al.* (1952) which gives consistent and reliable results. It is, however, difficult to avoid

occasional contamination of glassware or chemicals by amounts of iodine greater than the extremely small quantities one is attempting to estimate. A room must be set aside for this determination, and all experiments must be performed in duplicate. The normal range is usually taken as 3.5 to 7 $\mu\text{g.}/100\text{ ml.}$ The level is raised in thyrotoxicosis, but mild cases usually show a lesser degree of abnormality than would be expected from the radioactive iodine criteria. The administration of either inorganic or organic iodine compounds affects the accuracy of the method. We have personally found it of value in assessing thyroid function in cases after thyroidectomy and when ^{131}I studies have given inconclusive results. This estimation, moreover, gives a more complete picture of iodine and thyroxine metabolism when the results of simultaneous studies with radioactive iodine are available (Riggs, 1952).

Radioactive Iodine

In my Bradshaw Lecture I described the use of ^{131}I as a diagnostic aid in thyrotoxicosis. It is now generally agreed that if the necessary equipment is available, tests using radioactive iodine cause the patient little inconvenience and give quick and relatively reliable answers. Macgregor and I (1958) have summarized our present views of the relative value of the different techniques using ^{131}I . It is clear that several techniques give results which correlate well with the final clinical diagnosis. The main recent advance in this field has been the introduction of the short-lived isotope ^{132}I , which delivers a much smaller amount of radiation to the gland. In addition, tests have been devised which distinguish between the high radioactive iodine uptake of thyrotoxicosis and of iodine-deficient glands.

At Glasgow we have used the same methods as at Sheffield and have studied two parameters of thyroid function. It is of some interest that the application of these techniques in a different department using different apparatus and controlled by different personnel and applied to a different population in a different country has given essentially the same results. We give 30 μC of the isotope by mouth and measure the four-hour uptake of ^{131}I by the gland and the 48-hour protein-bound plasma activity (P.B. ^{131}I). The measurement of plasma activity has been rendered technically much easier by using an ion-exchange resin (Zieve *et al.*, 1956). Gland uptakes of greater than 45% and plasma P.B. ^{131}I levels of more than 0.4% indicate thyrotoxicosis. When the results agree, and both indicate normality or abnormality, the correlation with the final diagnosis is of the order of 98%. About 10% of results are, however, discordant. When the gland uptake is raised and the P.B. ^{131}I is within normal limits the patient usually has a non-toxic goitre.

When the gland uptake is within normal limits and the P.B. ^{131}I is raised it is wise to make sure that the patient is not suffering from Hashimoto's disease or has not been treated either by ^{131}I or by partial thyroidectomy. In such cases the functioning part of the gland may be working to the maximum of a diminished capacity to synthesize the thyroid hormones. There is a diminished thyroid pool, and a relatively large proportion of the circulating hormone is labelled with the isotope, giving a high P.B. ^{131}I . In general, however, the 48-hour P.B. ^{131}I correlates a little better than the gland uptake with the final diagnosis and it can be used as the sole diagnostic criterion. This test has the

advantage that the isotope can be given to patients at hospitals far removed from the centre at which the estimations are carried out. By carrying out the thymol turbidity tests or colloidal gold reaction on blood showing significant plasma radioactivity it is sometimes possible to distinguish between samples from thyrotoxic patients and those from cases of Hashimoto's disease.

Our experience of the tests which use renal excretion of the isotope as an indirect measure of gland uptake suggest that the best is that of Fraser *et al.* (1953). Its only defect is the potential inaccuracy of urine collection. It has the merit that the patient need not come into contact with counting apparatus or technical personnel and that the dose of ^{131}I is relatively small. Two types of clearance test are used in this country. Pochin (1950) prefers the neck-thigh ratio at two hours and Coenegracht and Fraser (1955) have reported favourably on a method described by Berson *et al.* (1952). It gives an answer quickly, since it measures the ^{131}I clearance during the 35 minutes following the administration of the isotope, which must, however, be given intravenously. It is said to be the most reliable test in the assessment of cases which have been treated by thyroidectomy or ^{131}I (Goolden, 1958). We tend, however, to rely on the chemical estimation of P.B. ^{127}I in such cases. The general problem of the relative reliability of methods of assessing thyroid function after ^{131}I therapy is discussed by Schultz and Zieve (1956).

Tests using radioactive iodine yield fallacious results if the patient is receiving iodine in the form of a cough mixture or Lugol's solution, and it is wise to perform a qualitative test for urinary iodide such as that described by Fraser *et al.* (1953) if unexpectedly low uptakes are recorded. Radiological contrast media which contain iodine may affect the uptake for many months. In practice, the most important fallacy in uptake tests comes from relative iodine deficiency. This may be associated with goitre or it may have been induced by the previous administration of an anti-thyroid drug. Thyrotoxic patients often have high uptakes for many months after treatment by thyroidectomy or by radioactive iodine. Fraser (1956) has described modified uptake techniques which in most of the above circumstances enable satisfactory results to be obtained using ^{131}I . We use these special tests comparatively seldom, since we find that, provided it is known whether the patient is receiving or has received treatment for suspected thyrotoxicosis, the two tests we use in combination almost always guide us to a correct result. When they are in agreement they are very reliable, and when they are discordant they give a warning that some other approach to the problem must be made. For example, in these circumstances we usually determine the B.M.R. or the P.B. ^{127}I , or both. If there is still a doubt a therapeutic trial with an anti-thyroid drug should be carried out.

Use of ^{132}I

This isotope with a short half-life of 2.26 hours has recently become available. Its rapid decay enables studies to be carried out at frequent intervals, and its radiation effects are of such short duration that it can be used in children and during pregnancy. Compared with ^{131}I it gives only about one-fortieth of the radiation dose to the thyroid for an equivalent number of microcuries. At first, like Hanbury *et al.* (1954) we found it difficult to obtain consistent readings, but these

difficulties have now been overcome, and a new method of preparing the isotope from ^{132}Te , using an alumina column, obviates some of the hazards arising from the original technique, which involved distillation (Stang *et al.*, 1957).

^{132}I can be used in routine thyroid-function tests. Halnan and Pochin (1958) studied the neck-thigh ratio two hours after an oral dose taken fasting. They found a good separation between thyrotoxic and euthyroid subjects, and if there was doubt they repeated the test after the administration of thyroxine or triiodothyronine. Farran (1958) has shown that the rate of uptake of ^{132}I by the thyroid during the first 30 minutes after the intravenous injection of 3–5 μc . of the isotope is also a useful diagnostic test which gives a 90% separation between thyrotoxic and euthyroid cases. For detailed studies of thyroid function the short-lived isotope possesses great advantages over ^{131}I (Goolden and Mallard, 1958). It is, for example, specially suitable for use in investigations that require the repetition of tests, and can be used to study day-to-day variations in thyroid activity (Francois *et al.*, 1958; Halnan and Pochin, 1958) or to investigate the effect of antithyroid drugs on thyroid activity. This advantage over ^{131}I is well shown in Fig. 4, where the short-lived isotope

FIG. 4.—Effect of triiodothyronine on the thyroid uptake of ^{132}I in a normal subject, a thyrotoxic patient, and a patient with an "iodine deficiency" goitre.

has enabled daily estimations of the gland uptake to be made.

The suppression of the gland uptake of ^{132}I by triiodothyronine in the normal individual and in a case of non-toxic goitre is contrasted with the absence of effect in a thyrotoxic patient. The short-lived isotope is more difficult than ^{131}I to prepare and handle. Its chief defect from our point of view is that it cannot be used to estimate the plasma activity at 48 hours, which is the test of which we have most experience and on which we put most reliance. We agree with Goolden and Mallard (1958) that it is unlikely to replace ^{131}I completely in standard diagnostic work. We use it as they do in special investigations in which we wish to repeat our observations at frequent intervals. A more complete picture of iodine metabolism is obtained if estimations of P.B. ^{127}I are carried out at the same time.

Treatment of Thyrotoxicosis

Three rival and satisfactory forms of treatment are available for treating thyrotoxicosis, and in a given case it is not always easy to decide which is the most appropriate. Surgical treatment is probably the most widely used. It has been made extremely safe by modern anaesthesia and by the pre-operative use of antithyroid drugs. Operation may, however, occasionally give rise to unfortunate sequelae. Thus, though rare, tetany from parathyroid damage, damage to the recurrent laryngeal nerve, wound sepsis, secondary haemorrhage, and adherent scars, as well as the pulmonary complications secondary to any surgical procedure, are still seen. The main criticism of surgery is the relative frequency with which patients relapse—a specially unfortunate occurrence if radioactive iodine treatment is not available, since a second operation is technically difficult and is especially liable to lead to hypothyroidism (Bartels, 1953). It is of interest that 10% of our series of 150 thyrotoxic cases treated by ^{131}I , and 12% of the 500 cases reported from Sheffield by Blomfield *et al.* (1959), were examples of recurrence of the disease after operation. The incidence of post-operative hypothyroidism is also high, ranging in different series from 2.9% to 21%, with a mean rate of 5.4% (Bartels, 1953).

It seems clear, therefore, that partial thyroidectomy is by no means the ideal form of treatment. The alternative methods involve the use of antithyroid drugs or of radioactive iodine. Much controversy surrounds the comparative effectiveness of different antithyroid drugs, and we have devised a method of comparison based on the history and physical examination which can be used to investigate this problem.

The Therapy Index

The clinician concludes that a patient is showing improvement when he observes the gradual disappearance of a number of abnormal symptoms and signs. This he can do fairly satisfactorily, but he finds that the degree of improvement is much more difficult to assess. For example, different clinicians may attribute different significance to any given change, and even the same observer may find it difficult to decide on the relative importance to be attached to conflicting features—for example, to a persistence of tachycardia in a patient who has gained weight. The alternative method of assessing improvement by repeated estimations of the B.M.R. is time-consuming and inconvenient to the patient. The normal range of B.M.R.s is so wide that it is difficult to know when a given patient has reached the level which is normal for her, and a comparison between various forms of treatment must be based on statistical differences between groups of patients.

We have approached the problem of assessing the response to treatment in a new way. We have modified the clinical diagnostic index, which I have already discussed, so that it can be used as a therapy index. To do this we have removed from the list of the diagnostic features of thyrotoxicosis those which are unaffected by treatment. The scores used for this purpose are given in Table II. It will be seen that the figures differ in some respects from those given in Table I for the diagnostic index. For example, the combination of loss of weight with increased appetite is highly significant diagnostically, but in the assessment of therapeutic response an increase in weight is more

TABLE II.—Thyrotoxicosis—Therapy Index. Weighting Factors Allocated to the Symptoms and Signs of Thyrotoxicosis

Symptoms	Score	Signs	Score
Dyspnoea on effort ..	+1	Hyperkinetic movements ..	+4
Palpitation ..	+2	Fine finger tremor ..	+1
Tiredness ..	+2	Hands:	
Preference for cold ..	+5	Hot ..	+2
Excessive sweating ..	+3	Moist ..	+1
Nervousness ..	+2		
Appetite increased ..	+1	Resting pulse rate:	
Weight decreased* ..	+4	>85/minute ..	+3

* For each 4 lbs. weight gain deduct 1 point.

important than a diminution in appetite. Hence the rating for loss of weight was increased to +4 and that for increased appetite reduced to +1. Patients were examined weekly in the out-patient department by the same clinician, and a standard routine was used. The resting pulse rate was recorded after the patient had been lying undisturbed on a couch for 10 minutes. The total score or therapy index was derived at the first visit and was determined at each subsequent weekly visit. A symptom or sign had to disappear completely before its value was deducted from the index.

Application of the Therapy Index

We first studied 40 patients who were definitely but not severely thyrotoxic and who had diffuse goitres and exophthalmos; that is to say, they were examples of classical Graves's disease. They were allocated alternately to two groups. Those in the first group were given methylthiouracil in a dose of 600 mg. a day for two weeks, followed by 300 mg. a day; those in the second group had 600 mg. of potassium perchlorate a day.

Typical examples of the changes in the index during treatment are shown in Fig. 5 (A and B) along with the changes in the B.M.R. A response of this type with a linear regression on time was shown by 33 out of the 40 patients. The other patients were slower to respond

FIG. 5.—Changes in therapeutic indices and B.M.R.s during treatment with methylthiouracil and potassium perchlorate (600 mg. daily). A and B show linear changes of the therapeutic indices with time. C and D show the delayed type of therapeutic response.

to the drugs as judged by the changes both in the index and in the B.M.R. Typical examples are shown in Fig. 5 (C and D). This also shows that a failure to respond immediately to an antithyroid drug does not preclude ultimate control.

Average rates of change were not used to compare the drugs, since they do not represent accurately the influence of the cases showing the slow type of response. An alternative method of comparison was therefore devised, and we determined the time taken for all the patients in each group to reach a level which we decided would indicate complete remission or "cure." This was taken to have occurred when the therapy index reached and remained within the range zero to 5 for three consecutive weeks. The actual week of "cure" was chosen as the week with the lowest of the three values. The range of indices from zero to 5 was chosen because it allows only one feature of high diagnostic significance to be present. We found, moreover, that, in patients satisfactorily controlled the indices fluctuated for long periods between zero and 5. Further

FIG. 6.—Times taken to effect "cure" by potassium perchlorate (600 mg. daily) and methylthiouracil.

confirmation that this range indicated full control of the disease was afforded by estimations of the B.M.R. in the two groups. When examined statistically the mean rates at the time of "cure" did not differ significantly from zero.

Fig. 6 shows the number of patients reaching the level we have defined as "cure" at any given time. From it can be calculated that the mean time to "cure" for the group receiving methylthiouracil was 9.1 weeks, and for that receiving potassium perchlorate 13.1 weeks.

When these patients were euthyroid they were given maintenance therapy for three months, and then the administration of the drugs was stopped. Of the 40 cases, 14 relapsed and were regarded as a separate group. Eight subjects who had received methylthiouracil were now given potassium perchlorate and six subjects previously treated with perchlorate were re-treated with methylthiouracil. The design of this series of observations was similar to that of the "cross-over" method used by pharmacologists for the assay of substances such as insulin. In the group receiving potassium perchlorate the time taken to effect a "cure" was four weeks longer than when they had been given methylthiouracil. Similarly, the group receiving methylthiouracil reached the score indicating a "cure" four weeks earlier than when they had been given potassium perchlorate. In these observations each patient acted as her own control.

We also made a comparison between the initial values of the therapy index and the times taken to effect "cure." This showed no significant correlation between the two variables. In other words, there is no relationship between the initial severity of the disease and the rate of therapeutic response. This runs counter to the generally accepted but unproved view that severe cases of thyrotoxicosis take longer to become euthyroid than mild ones.

Comparison of Methylthiouracil, Carbimazole, and Potassium Perchlorate as Antithyroid Drugs

We were now satisfied that our clinical method of comparison gave valid results and that it was unnecessary to check the patient's progress by repeated estimations of the B.M.R. It was thought sufficient to carry out this test only at the start and at the end of the trial. Two further groups of patients with classical Graves's disease were treated, one with potassium perchlorate in a dose of 1,000 mg. a day and the other with carbimazole in a dose of 60 mg. a day for two weeks followed by 30 mg. a day—that is, in one-tenth of the dose received by the group which had received methylthiouracil. The cumulative distribution for both drugs is shown in Fig. 7. The mean time to

FIG. 7.—Times taken to effect "cure" by carbimazole and potassium perchlorate (1,000 mg. daily).

"cure" for the potassium perchlorate treated group was 9.4 weeks and for the carbimazole series 12.3 weeks. Variance analysis of the mean times to "cure" of the four series of 20 patients receiving different drugs or different doses of perchlorate showed that methylthiouracil (600 mg. daily for two weeks followed by 300 mg. daily) and potassium perchlorate (1,000 mg. daily) were equally effective; both were more effective than carbimazole in one-tenth of the dose of methylthiouracil and potassium perchlorate in a dose of 600 mg. a day.

When we first began to use potassium perchlorate we were impressed by its relative freedom from unwanted side-effects. It was first shown to be a useful antithyroid drug by Godley and Stanbury (1954). This was confirmed in a much larger series of cases by Morgans and Trotter (1954), who had the impression that the rate of response produced by 400 mg. daily was less than that produced by methylthiouracil in a dose of 200 mg. daily. Since potassium perchlorate is a simple inorganic compound it seemed to us *a priori* less likely to produce sensitization effects than the organic antithyroid substances and in particular to be less likely to damage the haemopoietic system. We decided, therefore, to try to assess the relative frequency of the side-effects of potassium perchlorate, methylthiouracil,

and carbimazole. No statistically significant difference was found between the liability of the organic substances to produce sensitization reactions such as skin rashes or blood dyscrasias (Table III). On the

TABLE III.—Toxic Effects in Cases Treated With Methylthiouracil, Carbimazole, and Potassium Perchlorate

Toxic Effect	Methylthiouracil 600 mg. Daily 2 Wks. then 300 mg. Daily	Carbimazole 60 mg. Daily 2 Wks. then 30 mg. Daily	Potassium Perchlorate	
			600 mg.* or 1,000 mg.	1,500 mg.+ or 2,000 mg.
No. of cases	151	85	200	50
Skin rashes	5	4	1	5
Nausea	—	1	3	2
Drug fever	2	—	—	—
Agranulocytosis	2	2	—	1
Thrombocytopenia	1	—	—	—

* 35 patients. + 10 patients.

other hand potassium perchlorate in doses of 1,000 mg. a day or under was less likely than the organic antithyroid compounds to produce adverse effects of this type. Higher doses of potassium perchlorate produced reactions as often as the organic compounds. Of the four patients who complained of nausea while taking perchlorate three were able to continue when an antacid was given, and the nausea disappeared within a week.

We draw two generalizations from our observations. Both the rate of response to antithyroid drugs and their liability to produce sensitization reactions seem to vary with the dose. We have examined the extensive literature on the antithyroid drugs and have found it extremely difficult to assess the significance of the results reported. In comparing the relative activity of different compounds few observers seem to have been aware of the complexity of the problem. Not only must the clinician know about and allow for the phenomenon common to all biological material of individual variation in response, but he must also standardize, so far as he can, the type of patient he treats and the doses he uses. We have found no published observations which cannot be explained on the view that the response of a patient to an antithyroid drug is a function of the dose. At a certain critical blood level, which is different for each drug, the synthesis of the thyroid hormone is completely blocked and all drugs are equally effective when given in a dosage which will produce this or a greater level.

This immediately raises the question of the relative liability of the different drugs to produce toxic side-effects at different dosage levels. We have also examined the extensive literature on this subject. It is impossible to draw definite conclusions. Few clinicians have compared properly matched groups of patients under their own observation on standard doses of different drugs, and when they have done so, statistical tests of significance have rarely been applied to the figures. Most of the conflicting conclusions can be traced to one or other of these defects. We do not think it justifiable, as many have done, to draw deductions from groups of cases collected from the literature.

Our own tentative conclusions are these. There is no convincing evidence that any of the organic antithyroid drugs is significantly more toxic than any other when given in therapeutically equivalent dosage. Our experience suggests that potassium perchlorate is less likely to give rise to skin rashes or blood dyscrasias in doses which act as quickly as the conventional therapeutic doses of methylthiouracil and carbimazole.

Use of Potassium Perchlorate as an Antithyroid Drug

Since our investigations suggested that potassium perchlorate was the antithyroid drug of choice I should like to deal briefly with one or two practical aspects of its use. It affects the iodide-trapping mechanism of the thyroid, and there is evidence that the action of substances of this type can be overcome if sufficient iodide is given (Franklin *et al.*, 1944). We therefore made observations on groups of patients with relatively high and relatively low intakes of iodine in the diet. There was no statistical difference in the rate of response to perchlorate. There is thus no need to advise dietetic restriction of iodide intake, though we warn the patient against taking certain cough medicines which contain iodide in relatively large amounts.

We have used potassium perchlorate pre-operatively in 20 patients, of whom 18 had nodular goitres of moderate size, and our surgical colleague Mr. D. H. Clark did not encounter technical difficulties due to increased vascularity in this type of case. It may well be that the contradictory views which have been expressed on the effects of antithyroid drugs on the vascularity of the gland are due to the lack of an investigation designed to eliminate the effects of biological variation and of bias on the part of the surgeon.

Potassium perchlorate has been given to nine pregnant thyrotoxic patients and in each case satisfactory control of the disease was achieved. One of the infants had a very slight enlargement of the thyroid gland which disappeared within six weeks.

We now use potassium perchlorate in a dose of 250 mg. four times a day as our standard antithyroid drug, and so far we have had no reason to regret our choice.

Reserpine in Treatment of Thyrotoxicosis

Canary *et al.* (1957) have claimed striking improvement in thyrotoxic patients treated with reserpine. Ten patients who were given 0.25 to 1 mg. of the drug orally every six to eight hours improved considerably in three weeks. Palpitation, excessive sweating, heat intolerance, dyspnoea on exertion, the skin manifestations, and tremor were completely relieved in all the members of the group, and nervousness improved in all but one. Eye signs disappeared in three patients and improved in two. The gland's uptake of radioactive iodine, however, was not affected. Others (Moncke, 1955; Lövei and Bona, 1956; Chowdhury, 1958) have also claimed that reserpine is of value in the treatment of thyrotoxicosis, and Gordan (1958) states that since the publication of the paper by Canary *et al.* almost all the thyrotoxic patients he has seen have been treated with this drug.

Reserpine diminishes the results of sympathetic overactivity partly through central and partly through peripheral mechanisms. There are thus theoretical reasons why it might diminish the effects of an excess of thyroid hormone, since this may produce some of its peripheral actions by sensitizing the tissues to adrenaline and noradrenaline.

We did not feel justified in using this potentially toxic substance as an alternative to a standard antithyroid drug. We decided, however, that it might be given with advantage to patients immediately after the administration of radioactive iodine. It would then have its effect in the latent period before the isotope had begun to depress thyroid function. This is a period in which in

our view the administration of drugs which affect thyroid function is undesirable.

A double-blind trial was used, but, since the side-effects of reserpine might have enabled the observers to differentiate between the groups receiving the reserpine and the inert tablets, a second observer was interposed. Part of the interest of this trial lies in the use of this method, which could be applied to the evaluation of any toxic drug. That this precaution was necessary is demonstrated by the high incidence of severe toxic effects which occurred in the group receiving reserpine.

Twenty patients who had received ^{131}I therapy took part in the trial. They were allocated by random selection to two groups, one of which received reserpine by mouth in a dose of 0.5 mg. four times a day and the other inert tablets of identical appearance. They were interviewed by a physician who recorded the blood-pressure and supplied the tablets, and who noted any symptoms or signs which either he or the patients attributed to the tablets. The patient was then told to withhold this information from the second physician, who assessed the degree of thyrotoxicosis.

Progress was judged by the weekly assessment of those symptoms and signs which improved in the cases treated by Canary *et al.* (1957). Neither at four weeks, the time at which Canary *et al.* made their observations, nor at six weeks was there any difference between the two groups in the number of clinical features recorded except in the case of the pulse rate and weight. There was a significant fall in the pulse rate ($P < 0.01$) and a significant gain in weight ($P < 0.05$) in the reserpine-treated series. Full details of this work have been published elsewhere (Buchanan *et al.*, 1959).

It is a remarkable and rather disturbing fact that so toxic a substance can have been widely used in the treatment of a disease for which much safer and more specific remedies are available. It is all the more remarkable when a controlled trial shows that reserpine administered orally in doses apt to give severe side-effects produces no obvious benefit in a group of thyrotoxic patients. Indeed, the only alterations in our series which could be attributed to reserpine were due to a non-specific action of the drug. Clinicians have no doubt been unduly impressed by the increase in weight and the slowing of the pulse rate which reserpine produces in thyrotoxic patients. When these are secondary to a diminished amount of circulating thyroid hormone they undoubtedly indicate improvement in the disease. But reserpine can produce these effects in euthyroid individuals, and in themselves they confer no special benefit on the patient. The thyrotoxic patient given deoxycortone and salt would put on weight, but this could hardly be regarded as evidence that it was a valuable form of treatment.

We conclude that reserpine has no place in the treatment of thyrotoxicosis except possibly in a thyrotoxic crisis, where different considerations apply.

Radioactive Iodine in Treatment of Thyrotoxicosis

It is universally agreed that ^{131}I is effective in the treatment of thyrotoxicosis, and there are some who believe that it is the best and safest form of treatment for every case of the disease. The majority of centres, however, use this form of therapy only in selected cases. There are several excellent recent reviews of this subject (Myant, 1956; Pochin, 1958; Blomfield *et al.*, 1959), and I do not propose to cover the same ground. I shall outline our own criteria for the selection of cases and

the simplified method we use for deciding on the dose of the isotope. We have compared the results of our technique with those of more complex methods, and in addition have made observations on the influence of pre-treatment with methylthiouracil on the radio-resistance of the thyroid gland.

Selection of Cases

The chief consideration which limits the use of ^{131}I in the treatment of thyrotoxicosis is fear of the late development of malignant changes in the thyroid gland. Many thousands of patients have been treated in the past 15 years and so far there are no reports of cases of carcinoma which could be attributed to this form of treatment. There is, however, evidence that ionizing radiations can lead to the development of malignant disease of the gland in children (Wilson *et al.*, 1958). It seems improbable that the incidence of post-radiation carcinoma in the adult, if it occurs at all, will be high. Most clinicians, however, prefer to treat patients with an expectation of life of 20 to 25 years. Younger patients may be treated if surgery is contraindicated. In women before the menopause and in comparatively young men the possibility of genetic effects arising from irradiation of the gonads should be taken into account (Blomfield *et al.*, 1959).

These general considerations lead us to select the following groups of patients as suitable for treatment with ^{131}I . (1) Patients over the age of 45 in whom it is the treatment of choice. (2) Any patient who, because of associated disease, has a life expectancy of less than 20 years. (3) All patients, irrespective of age, who have relapsed following thyroidectomy. A second operation is specially apt to produce myxoedema or damage structures adjacent to the gland. (4) Young patients who have relapsed following antithyroid drugs or who have shown sensitization reactions to them but only if they have refused operation.

The only absolute contraindication to ^{131}I is pregnancy. A few workers are unwilling to treat a case with a multinodular gland, especially if there is tracheal compression. As a rule, however, such cases fulfil our criteria of suitability, and in our experience ^{131}I therapy gives satisfactory results.

Methods of Deciding on Dosage of ^{131}I

The selection of the appropriate dose of ^{131}I is important. It is the aim of the physician to find a dose which will cure the patient without producing hypothyroidism. A few centres give small and repeated doses of the isotope, hoping thus to minimize the possibility of overtreatment of the case (Williams *et al.*, 1949). The majority, however, emulate the surgeon and try to give a single dose which will destroy the optimum amount of functioning tissue. In general they are no more and no less successful than the surgeon, and the overall incidence of myxoedema in 1,720 cases from different centres was 9% (Seed and Jaffé, 1953). In the investigation carried out in Sheffield we attempted to deliver to the thyroid gland a dose of between 6,000 and 8,000 rads (Blomfield *et al.*, 1955). We used a formula which involved a knowledge of the 48-hour uptake of a tracer dose of ^{131}I and of the weight of the gland. A review of 140 cases showed that 62% of patients were made euthyroid with one therapeutic drink and ultimately 84% were cured; 12% showed evidence of hypothyroidism. One member of this team went to Edinburgh and reported the results of treating

150 cases of thyrotoxicosis using a greatly simplified dosage scheme (Macgregor, 1957). The results did not differ significantly from those obtained by using the more elaborate and quantitative technique described above.

In Glasgow we have used a similar dosage scheme. To patients with glands which could not be palpated we have given 5 mc., to those with definite diffusely enlarged but small glands 6 to 7 mc., and to those with larger glands doses up to 25 mc. Factors other than gland size have influenced our decisions on dosage; for example, patients who had a previous thyroidectomy received relatively small doses, usually between 5 and 6 mc., while those with nodular glands and patients with cardiac failure received relatively large doses. All patients were seen at monthly intervals and were retreated three to four months after the initial dose if they were still thyrotoxic. After they had become euthyroid they were seen at intervals of from two to six months. Hypothyroidism was treated only after it had persisted for from three to four months.

The results of treating by this method a series of 100 cases followed for a minimum period of 12 months is shown in Table IV. For comparison we give the figures

TABLE IV.—Clinical Results of Radioactive Iodine Therapy (Minimum Follow-up, One Year)

	Present	Macgregor (1957)	Blomfield <i>et al.</i> (1959)
No. of cases	100	150	500
Euthyroid with one dose	67%	59%	59%
Myxoedema	6%	9%	12%

for a much larger series of 500 cases reported from Sheffield by a different group of workers (Blomfield *et al.*, 1959) and for the 150 cases to which we have already referred, treated at Edinburgh by Macgregor (1957). There is no significant difference in the figures for the one-dose cure-rate or for the incidence of hypothyroidism. It should, however, be remembered that our minimum follow-up period of one year underestimates the incidence of hypothyroidism which may develop several years after treatment.

We, however, had also determined the 48-hour uptake of ^{131}I and estimated the size of the gland in grammes, so that we were able to calculate the dose we would have given if we had been using the more precise Sheffield technique. This alternative dose was not calculated until after treatment so as to avoid any bias in the original dose prescription. We have plotted the doses we actually used against those which would have been given if we had used the Sheffield procedure. Fig. 8 shows that we have tended to give larger doses to those in the small-dose range and smaller doses to those in the large-dose range. This phenomenon is probably due to the reluctance of those using the clinical method of dose prescription to prescribe very large doses in the absence of the moral support of a formula. The difference is, however, not reflected in the one-dose cure-rate probably because of the relatively small numbers of patients who required large doses, and also because these patients tend to be under-treated even when the more precise method is used (Blomfield *et al.*, 1959).

We agree with Macgregor that the prescription of an entirely accurate radiation dose is not possible partly because of errors inherent in the technique for the estimation of the gland mass, partly because of differences in the behaviour of the tracer and therapeutic

doses, and partly because of the biological variability in the radiosensitivity of different glands. With his method thyrotoxic patients can be treated safely and effectively without elaborate techniques. Though we do not allow for the percentage uptake of ^{131}I by the gland in deciding on the dose to be given, we consider preliminary tracer investigations desirable to confirm the diagnosis.

Pre-treatment of thyrotoxic cases with methylthiouracil and thyroxine before giving ^{131}I has been used by Fraser *et al.* (1954) to produce a more uniform uptake of the isotope by the gland. We investigated the effects of pre-treatment with methylthiouracil on the results of ^{131}I therapy. A series of 28 cases treated with methylthiouracil until one week before the ^{131}I was administered provided a significantly smaller number of one-dose cures (28%) than a comparable series which received no premedication (75%). Further investigations have shown that this effect is probably due to an increased degree of resistance to radiation conferred on the gland by the pre-treatment. This factor must be taken into account whatever scheme of dosage is used, and further observations suggest that a dose at least 25% larger must be given to those who have recently received methylthiouracil.

Fig. 8.—Therapeutic doses of ^{131}I , prescribed in the present series, plotted against doses calculated from the formula of Blomfield *et al.* (1955). The oblique line has been drawn at 45 degrees to the ordinates.

Choice of Treatment

No hard-and-fast rules can be laid down, but the clinician will show wisdom if he takes advantage of ^{131}I treatment when it is available and uses it to treat patients whose disease has recurred after operation and those with heart disease. It is of course much the most pleasant form of treatment, and if I were to become thyrotoxic it is the form I would choose. Younger patients are usually best treated in the first instance

with a course of an antithyroid drug such as potassium perchlorate. Those who relapse should be advised to have a thyroidectomy after they have been rendered euthyroid by a further course of an antithyroid drug.

Figs. 1, 2, and 3 are reproduced from the *Quarterly Journal of Medicine* by permission.

[The Second Lecture will be published next week, with the list of references.]

REVERSAL OF ANTIBIOTIC RESISTANCE IN HOSPITAL STAPHYLOCOCCAL INFECTION

BY

MARY BARBER, M.D.

Reader in Clinical Bacteriology

A. A. C. DUTTON, M.A., B.M., B.Ch.

Assistant Lecturer

M. A. BEARD, BSc.*

Research Assistant

Department of Bacteriology

P. C. ELMES, M.A., M.D., M.R.C.P.

Late Senior Registrar

ROGER WILLIAMS, M.B., M.R.C.P.

Late Registrar

Department of Medicine

Postgraduate Medical School of Medicine, Hammersmith
Hospital, London

There is much evidence to suggest that some at any rate of the multiple-resistant strains of *Staph. pyogenes* so prevalent in our hospitals to-day are of enhanced virulence, both in their capacity to initiate infection and in the severity of the infection produced. Thus, although such strains are relatively uncommon among staff carriers, they are responsible for the majority of cases of post-operative infection (Alder *et al.*, 1955; Barber and Burston, 1955; Shooter *et al.*, 1958), and some investigators have reported that infection with "hospital staphylococci" may be of unusual severity (Rountree and Freeman, 1955; Beaven and Burry, 1956; Ede *et al.*, 1959). While it cannot be assumed from this that multiple resistance and enhanced virulence are necessarily associated, the process of selection of drug-resistant staphylococci in hospitals may have tended to select strains of high virulence. In fact, nearly all the multiple resistant staphylococcal infections in hospitals to-day are due to a relatively few strains of *Staph. pyogenes* most of which belong to phage group III or type 80.

The principles for preventing hospital cross-infection enunciated long before the introduction of antibiotics remain the same to-day, but the ubiquity of the staphylococcus makes their strict application difficult in relation to this microbe. It is possible that pathogenic staphylococci could be kept at bay if we had sufficient isolation facilities, but at present this is not the case in most hospitals in Britain.

It seemed reasonable, therefore, to attempt to break the vicious circle by introducing a controlled antibiotic policy, designed to inhibit the emergence of drug-resistant staphylococci and to favour penicillin-sensitive strains. The present investigation was undertaken to

*Working with a grant from the Medical Research Council.