of medical knowledge on the part of the patient would seem to be a prerequisite for development, even though Barker and Grygier (1958), among others, have encountered the condition in a subject with mental deficiency. In addition, it is likely that other factors such as illegitimacy (Grant, 1952) and drug addiction (Todd, 1951) play a part in some cases.

The reasons for the development of the syndrome in the case described here remain uncertain, though it is reasonable to suppose that the professional apprehension shown after the initial electrocardiogram was duly recognized by the patient and taken advantage of on subsequent occasions. Unfortunately, it has not been possible to discover whether electrocardiography was performed before December, 1953 (the first available record). It is conceivable, of course, that electrocardiography was originally carried out on account of genuine cardiac pain associated with a small coronary occlusion or acute pericarditis. Whether or not this was so, the possession of a sickle-shaped R-T electrocardiographic pattern has enabled the patient to simulate organic heart disease on numerous occasions during the past five years. It is certainly regrettable that the psychopathic origin of such behaviour is likely to prove a stumbling-block in his management and treatment in the future.

ADDENDUM.—After exercise the S-T segment of the sickle-shaped R-T plateau pattern returns to the isoelectric line. For this reason, if this patient attends other hospitals in the future it would be of interest to perform electrocardiography before and after exercise.

I thank those cardiologists who have kindly given their valuable advice regarding the interpretation of the electrocardiograms of this patient. I would also like to acknowledge the help given by the medical staff of the numerous hospitals which this patient has attended.

REFERENCES

- Barker, J. C., and Grygier, T. G. (1958). *Brit. med. J.*, 2, 1274.
Bedford, D. E., and Thomas, G. (1954). *Brit. Heart J.*, 16, 469.
Clyne, M. B. (1955). *Brit. med. J.*, 2, 1207.
Davidson, C. (1953). *Lancet*, 2, 621.
Grant, A. P. (1952). *J. Irish med. Ass.*, 31, 299.
Heathfield, K. W. G. (1957). *Lancet*, 1, 533.
Todd, J. (1951). *Ibid.*, 1, 527.
Todd, J. W. (1959). *Ibid.*, 1, 845.

Steady progress is being made in the campaign to rid Scotland of smoke, and clean-air exhibitions have recently been opened in Glasgow and Edinburgh. The Clean Air Council for Scotland—of which Sir DAVID ANDERSON is chairman—has been told that 26 town and county councils have already made smoke control orders or are considering making them. Altogether 90 local authorities have been urged to set up smoke-controlled areas. Arrangements are being made for appeals to householders in smog-prone areas to burn smokeless fuel or to use gas or electricity when conditions become bad. It is known that about half the smoke-polluting air comes from household fires. Some of this comes from the lighting of fires, and the council is having a new smokeless firelighter tested by its technical advisers. The possibility of an electrical firelighter for use in houses where there is no gas is also being studied. Another problem being examined is the effects of fumes from buses and other motor vehicles. The Clean Air Council for Scotland is in touch with the Clean Air Council for England and Wales in the whole field of smoke control.

INFANTILE SCURVY

BY

M. K. STRELLING, M.B., D.C.H., D.Obst.R.C.O.G.

Late Paediatric Senior House Officer, Morriston
Hospital, Swansea

That dietary deficiency of vitamin C remains a possibility is emphasized by instances of overt scurvy which, exceptionally, do still occur. The following examples in infants and children have been reported from England and Wales in the last decade.

Six out of 10 cases seen at the Hospital for Sick Children, Great Ormond Street, in 1947–9 were in mental defectives aged 23 months to 4½ years who were cared for at home by their parents. Their diet, perforce liquid or semi-solid, was mainly milk, custard, cereals, milk puddings, cakes, biscuits, and tea. Vegetables were refused, and orange juice was either refused or vomited and discontinued, or never offered (Beynon and Evans, 1951).

Derham (1950) describes the case of a 10-year-old girl who, in addition to a history of tuberculous peritonitis with prolonged diarrhoea, had perverse eating habits to which much attention had been paid, refusing fruit, green vegetables, milk, and potatoes.

Baber (1953) reports two cases. One patient at 9 months was still on cow's milk and cereal. Orange juice had caused vomiting and had been discontinued early. The other developed scurvy at 4 months while in hospital with gastro-enteritis, having had orange juice for two weeks only.

From a population of 250,000 in Cornwall, seven further cases were seen by Howells *et al.* (1954). These babies of 7 to 12 months had been fed on boiled cow's or National dried milk since birth; none had had solids, and six had not had orange juice.

Reports of adult scurvy appearing in the same period (Riches, 1954; Dewhurst, 1954; Hughes and MacLennan, 1954) confirm Cutforth's (1958) finding that they fall predominantly into two groups: (1) old men living alone, failing to feed themselves properly; and (2) those whose rigid diets, usually for peptic ulcer, exclude fruit and vegetables (save mashed potato) for periods of a year or more. One woman of 64 had had little but tea, sugar, beef extract, biscuits, bread, butter, and paste for 26 years (Glyn and Campbell, 1957).

Case Report

Scurvy in this child is reported because he was receiving a normal diet in an institution in which regular vitamin C supplements were given.

A boy of 15 months was admitted to hospital in 1958 with a painful swelling of one leg; diagnosis? osteomyelitis. Since he was 2 weeks old he had been in a pleasant home for deprived children run by nuns under medical supervision. Details of family and birth are unknown.

Previously normal, he had been listless for a fortnight with a swelling of his right thigh, and for five days he screamed on being handled. The sister, noting his gums to be swollen and to bleed, had in fact given him extra rose-hip syrup for two weeks. His appetite, usually good, had been poor for two days.

On examination he was fairly well covered, pale, and fretful, with a temperature of 101.8° F. (38.8° C.). He kept his right leg still in the classical "frog position"—that is, externally rotated and moderately flexed at hip and knee.

FIG. 1.—Radiograph showing characteristic changes in lower ends of femurs.

There was a diffuse, firm, warm swelling with crepitus above the knee, and the overlying skin, though shiny, was not reddened. He screamed when this was touched. Marked angular rib-beading at the costo-chondral junctions and a purplish-red swelling of the gums around erupted teeth were also noted. A positive Hess test demonstrated increased capillary fragility. There was no other physical abnormality.

Investigations.—A microcytic hypochromic anaemia: Hb, 56%; colour index 0.65. W.B.C. 9,000/c.mm. (polymorphs 50%, lymphocytes 44%). Urine—no excess of R.B.C.s. Serum calcium, phosphorus, alkaline phosphatase, platelet count, bleeding-time, and clotting-time were all normal. W.R. and Kahn test negative. The urine ascorbic acid content two and five days after starting treatment was 5 mg. and 3 mg./100 ml. respectively.

FIG. 2.—Radiograph showing marked calcification in subperiosteal haematoma at 18 days.

haematoma just starting to calcify; and (4) displacement of the right lower femoral epiphysis. Bone rarefaction and thinning of the cortex were not visible, possibly because extra rose-hip syrup had already been given.

He was given 500 mg. of ascorbic acid daily for three weeks, reducing to 200 mg. a day, coupled with oral iron and galls traction to his legs.

After a week he was happier, was eating better, and his leg was less tender. He ran a temperature of 99–101° F. (37.2–38.3° C.) for 18 days, but no infection was found. By one month his gums were normal; haemoglobin 78%. At two months the rib-beading had disappeared. The Hess test remained positive until the sixth week. Radiographs

showed marked calcification in the subperiosteal haematoma at 18 days (Fig. 2), and at seven weeks (Fig. 3) healing was advanced (left leg).

Discussion

The cause of this child's scurvy was not at first obvious. He had been given 50 mg. of ascorbic acid daily until he was 9 months old, and then, for six months, one teaspoon of rose-hip syrup a day. At 57 mg./fl. oz. (200 mg./100 ml.) this roughly equals 7 mg.

His diet for six months had been "farex," milk (about a pint (570 ml.), pasteurized and boiled), bread-and-butter, rusks, old potatoes (mashed), meat, cabbage or carrots, milk pudding, and plain cake. Banana, stewed apple, or orange juice was given no more than once a week. Uncooked, these vegetables could provide anything between 10 mg. and 130 mg. of ascorbic acid a day.

The minimum vitamin C intake which will prevent scurvy is not accurately known. 10 mg. daily is usually quoted, this amount having protected adult volunteers from scurvy for 424 days, and also cured six in whom the disease had been induced (Krebs, 1953). Lower figures have been suggested. Cruickshank (1959), for example, regards 7 mg. a day as protective for a baby, while Mottram and Graham (1956) suggest 4 mg.

This child developed scurvy while receiving something over 7 mg. a day for six months, having had an optimum dose previously, and there could therefore have been very little in his food.

Losses could have occurred in several ways, and are perhaps particularly apt to occur in institutions. For example: (1) In storage. Vegetables, as opposed to fruit, rapidly lose their vitamin C at room temperature. (2) In cooking. Maximum losses occur if vegetables pass through the 65°–85° C. range, when plant oxidases are most active. The potatoes were in fact put in cold water and boiled, while the cabbage and carrots, dropped in large amounts into boiling water, almost certainly lowered the temperature below 85° C. (3) In keeping food hot, and in mashing potatoes. (4) In boiling milk after it has been pasteurized, which destroys most of the vitamin C. Boiling fresh milk for five minutes, however, does not (Woodruff, 1956).

The above case therefore emphasizes how a diet which is relatively fixed, as in an institution, and relies on every-day methods of preparing vegetables may be grossly deficient in vitamin C. Ebbs (1951) stated: "Gingivitis, a common finding in institutional groups of children and young people, if cleared by local treatment,

FIG. 3.—Radiograph taken at 7 weeks, showing advanced healing.

can be prevented from recurring by the administration of adequate amounts of vitamin C."

Clinically, the subperiosteal haematoma, the tenderness and "frog position" of his leg, plus marked radiological changes in rapidly growing bones were characteristic of infantile scurvy. Skin and subcutaneous haemorrhages, and hyperkeratotic hair follicles, seen in older patients, were absent. Anorexia, swollen gums, anaemia, and low-grade fever (probably due to blood-absorption) may be features of scurvy at any age.

Summary

Reference is made to 16 examples of scurvy in infants and children reported from England and Wales in the past 10 years. All but one failed to take a mixed diet containing fruit and vegetables due to (a) lateness in weaning (eight infants aged 7 to 12 months), (b) mental defectiveness (six children aged 23 months to 4½ years), or (c) food faddism (one girl of 10 years), and all were combined with failure to offer orange juice, or a substitute if refused or vomited. Malnutrition due to gastro-enteritis accounted for one at 4 months.

Of 17 adult cases reported over the same period, most were either old men living alone or people on restricted diets chiefly for peptic ulcers.

Scurvy occurring in a 15-months-old boy in an institution is described. The accepted minimum daily vitamin C requirement of 10 mg. had not been met by a mixed diet, including vegetables and a teaspoonful of rose-hip syrup containing about 7 mg. of ascorbic acid.

Reasons for the low dietary content are suggested, and the liability of this to occur in institutions is emphasized.

I am grateful to Dr. R. T. Jenkins, Dr. K. Mendl, and Mr. Mervyn Evans for their help and encouragement.

REFERENCES

- Baber, M. D. (1953). *Med. Ill. (Lond.)*, 7, 539.
 Beynon, D. W., and Evans, P. R. (1951). *Gt Ormond St J.*, No. 2, p. 90.
 Cruickshank, E. W. H. (1959). *Practitioner*, 182, 29.
 Cutforth, R. H. (1958). *Lancet*, 1, 454.
 Derham, R. (1950). *Arch. Dis. Childh.*, 25, 159.
 Dewhurst, K. (1954). *Brit. med. J.*, 2, 1148.
 Ebbs, J. H. (1951). In *Modern Trends in Paediatrics*, edited by L. Parsons, p. 211. Butterworth, London.
 Glyn, J. H., and Campbell, E. J. M. (1957). *Ann. phys. Med.*, 4, 137.
 Howells, G., Palmer, P. E. S., and John-Brooks, W. H. (1954). *Brit. med. J.*, 2, 1143.
 Hughes, W., and MacLennan, D. C. (1954). *Ibid.*, 2, 1149.
 Krebs, H. A. (1953). *Proc. Nutr. Soc.*, 12, 237.
 Mottram, V. H., and Graham, G. (1956). *Hutchinson's Food and the Principles of Dietetics*, 11th ed., p. 141. Arnold, London.
 Riches, H. R. C. (1954). *Brit. med. J.*, 2, 286.
 Woodruff, C. (1956). *J. Amer. med. Ass.*, 161, 448.

"The increasing clarity of the atmosphere has been noted on many occasions during the year; the effects of the steady improvement in the condition of the atmosphere on health will take longer to become evident. Custom and a mistaken conception of comfort are difficult hurdles to overcome, but the response of the public to the measure for clearing the atmosphere has been very encouraging. Privy middens were once usual but are now unacceptable—in twenty years' time people will recount with amused censure and an air of superiority. 'Fancy! they had open coal fires.'" (*Annual Report on the Health of the City of Sheffield*, 1958.)

OBESITY IN CHILDHOOD

A CLINICAL TRIAL OF PHENMETRAZINE

BY

JOHN RENDLE-SHORT, M.D., M.R.C.P., D.C.H.

Lecturer in Child Health, University of Sheffield

Phenmetrazine ("preludin") is a synthetic derivative of ephedrine. It is not pharmacologically related to amphetamine. It has been much used in the treatment of obesity owing to its effect on the appetite. According to the manufacturers the primary action is on "the satiety centre in the hypothalamus. Satisfaction of appetite occurs sooner than normal. On the other hand, the urge to eat remains unaffected." The value of phenmetrazine in the control of obesity in adults has been demonstrated by several controlled investigations (Fazekas, Ehrmantraut, and Kleh, 1958; Leith and Beck, 1958; Patterson, 1959). There are also references in the literature to the value of phenmetrazine in the control of obesity in children (Bleckmann and Salus, 1955), but no clinical trial of the use of the drug in children has so far been published. For this reason the present study was undertaken.

Present Investigations

The obese children included in this investigation were treated as out-patients. They all weighed over the ninetieth percentile above the average weight for their age, and all were gaining weight at the beginning of the trial. During the investigation they were on their normal diet.

The children received no treatment during a preliminary period of 14 days. They were weighed at the beginning and at the end of this period. They were then sent to the pharmacist, who selected at random a paper from an envelope. This stated whether the child was initially to be given phenmetrazine or dummy tablets.* Sufficient tablets for 28 days' use were dispensed. The dose of phenmetrazine was 25 mg. in the morning and midday for children aged 8 years and over, and 12.5 at the same times for children under 8. At the end of the 28 days the child was weighed and a further supply of tablets was prescribed. The second set of tablets was the opposite—active or dummy—to that which the child had received previously, so that each child acted as his own control. At the end of the second period of 28 days the child was again weighed and then took no further part in the investigation. The children and their parents did not know that the tablets were different, and only the pharmacist knew which tablets the child received until after the investigation was completed and the results had been tabulated.

Eleven children did not complete the full course and are not included in the series. This was because of two main reasons. (1) Some children failed to return for weighing at exactly the right time because of holidays or for other domestic reasons. Many of these children attended a week later, but as they had been without tablets for a week they could not be included in the trial. (2) Some children had an illness, such as measles or tonsillitis, which probably affected their appetite.

*The phenmetrazine and dummy tablets were kindly supplied by Boehringer Products division of Pfizer Ltd.