Drug Treatment of Disease**THYROTOXICOSIS**

BY

OLIVER GARROD, M.B.E., M.D., M.R.C.P.*Physician, Barnet General Hospital*

The treatment of thyrotoxicosis requires both radical and palliative measures. Radical treatment aims at a partial suppression or ablation of the overactive thyroid gland. Suppression can be achieved with antithyroid drugs, and ablation either with radio-iodine or by subtotal thyroidectomy. Surgery will be mentioned only in connexion with pre-operative therapy, and to indicate where it is the treatment of choice. Some form of palliative or symptomatic treatment is usually required, at least until radical measures have taken effect. It may also be needed to control the ophthalmic, cardiac, or neurological complications of the disease.

RADICAL TREATMENT

Choice of treatment is governed by many factors. Among these are the age of the patient; the severity and duration of the disease; the size, type, and situation of the goitre; the scope and nature of any previous treatment; the kind and severity of the complications; the facilities available; and to some extent the wishes and fears of the patient after fair presentation of the various pros and cons.

ANTITHYROID DRUGS**Iodide**

Until 1944 iodide was the only safe and effective therapeutic agent for blocking the release of thyroid hormone. Iodide, by ways that are still obscure, causes involution of a toxic goitre when given in relatively large amounts. Its action is rapid, increasing the storage and diminishing the release of thyroid hormone, and so lessening toxicity. Also, it facilitates the surgeon's task by reducing the vascularity and increasing the firmness of the gland. Remissions were not infrequent from its use in mild cases of diffuse toxic goitre. But in severer cases usually the effects were transient, lasting only a month or two. In toxic goitre iodide is now used only as a preparation for subtotal thyroidectomy. It is prescribed traditionally as Lugol's iodine—an aqueous solution of iodine and potassium iodide—0.3 ml. three times a day. The operation must be timed to coincide with the optimal effect of iodide on the gland, usually 14–21 days after starting treatment. Mild cases can safely be prepared for surgery with iodide alone, without risk of post-operative thyroid crisis. Severer cases should first be rendered less toxic by a 6–8-weeks course of carbimazole ("neomercazole") (see below), followed or overlapped by the standard course of Lugol's iodine.

Thiouracil and Related Compounds

Since the advent of methyl thiouracil in 1944, a number of drugs have become available for the control of thyrotoxicosis. The four compounds most widely used have been methyl thiouracil, propyl thiouracil, and more recently methimazole and carbimazole. All are reducing agents and act by interfering with the iodination of tyrosine, so interrupting the synthesis of thyroid

hormone. They cause hyperplasia and increased vascularity of the gland. They are all potentially toxic.

More recently a simpler and probably much safer compound, potassium perchlorate, has been introduced for this purpose. It acts differently, by interfering with the trapping of iodide by the thyroid gland. It is now used routinely in at least one centre, and more generally as alternative treatment for those who cannot tolerate the other drugs.

Antithyroid drugs are used either in the short-term preparation of more toxic cases for surgery, or as a long-term measure. Pre-operatively, they are given for 6–8 weeks, followed or overlapped by the standard course of Lugol's iodine. Long-term treatment in adults is seldom to be recommended except in mild cases in young women without large or nodular goitres, whose disease is often transient like the emotional crisis which may precipitate it. In pregnancy it is the treatment of choice, though surgery between the third and sixth month still has its advocates. Overdosage should be avoided, for it may cause transient goitre and hypothyroidism in the baby. Thyrotoxicosis in men tends to be severe and protracted and should be tackled more radically. In children, subtotal thyroidectomy carries an increased risk of hypothyroidism later in life. For this reason treatment for at least two years with antithyroid drugs is to be recommended in the first instance, provided the goitre is not very large or does not become so during the course of therapy, in which case surgery will be required.

In general, treatment should be continued for at least 12 months. Unless there is careful selection of cases there will be a high rate of relapse on stopping the drugs, which will necessitate treatment by surgery or radio-iodine.

Carbimazole

In Great Britain carbimazole has largely replaced the thiouracil derivatives on account of its lower toxicity and a clinical impression that it causes less hyperplasia and vascularity of the gland. It is approximately 10 times as potent as methylthiouracil. The initial dose is 10–15 mg. 8-hourly, the larger dose being given only for large, nodular, or severely toxic goitres. The dose is halved after 4–6 weeks, depending on the response, and thereafter brought down at monthly intervals to the least amount which will control the disease. In mild cases this may be as little as 2.5 mg. twice a day, the more usual dose being 5 mg. twice a day, or 2.5 mg. three times a day.

Toxic reactions from carbimazole, though less frequent than from the thiouracils, are of similar nature and occur most often within 3–6 weeks of starting treatment. As with other antithyroid drugs, their incidence increases with the dose. The commonest reactions are itching, erythema, and urticaria of the skin, sometimes combined with conjunctivitis, lymphadenopathy, and fleeting joint pains simulating rheumatic fever. Though the milder reactions may be

amenable to control by antihistamines, such as promethazine ("phenegan") 25 mg. nocte, it is nevertheless wise to switch over to potassium perchlorate for suppressing the thyrotoxicosis (see below). More dangerous toxic reactions fortunately are very rare: these include agranulocytosis, thrombocytopenic purpura, and aplastic anaemia. The drug must be withdrawn at once, and appropriate protective measures applied such as antibiotics, blood transfusion, and corticosteroids. Examination of the sternal marrow may show early regeneration before signs of this appear in the peripheral blood. The prognosis for recovery is much more favourable with agranulocytosis than the other blood dyscrasias.

Potassium Perchlorate

The therapeutic status of potassium perchlorate as an antithyroid drug is still under review. A great advantage is its low toxicity, though this increases with larger doses. Toxic symptoms consist of nausea and a maculopapular rash. Only two cases of agranulocytosis have so far been reported, but more may follow. Minor disadvantages of the drug are a tendency to cause undue vascularity of the gland, a somewhat slower action than the thiouracils, and possible interference with its action from a high intake of iodide. The recommended initial dose is 600–1,000 mg. a day in 4–5 divided doses, this being halved after 2–4 weeks.

RADIO-IODINE

Radio-iodine, first used in 1942, offers the safest treatment for thyrotoxicosis in the majority of older patients. It avoids the toxic reactions of antithyroid drugs and the disastrous surgical hazards of laryngeal palsy and hypoparathyroidism, not to mention the acute risks of laryngeal oedema and reactionary haemorrhage causing tracheal compression.

Unfortunately its use has to be restricted because of certain genetic and carcinogenic risks, the true extent of which is not yet known. The first risk applies to pregnant women and concerns the foetus and its progeny. The second risk, to the remaining thyroid tissue, seems to be minimal, at least over an 18-year period.

Current policy in Great Britain is that therapeutic doses of radio-iodine should be given only exceptionally to patients under 45 years of age whose mean life expectancy would exceed 20 years. In those over 45 it is the treatment of choice for: (1) diffuse toxic goitre; (2) toxicity recurring after previous subtotal thyroidectomy; (3) very nervous patients who fear surgery; (4) patients with severe thyrotoxic heart disease or other debilitating conditions, after adequate preparation with antithyroid drugs.

With toxic nodular goitres the rate of recurrence after radio-iodine is higher than with diffuse toxic goitres, and several courses of treatment may be required. For this reason, and because malignancy cannot always be excluded in these patients, there is still a case in older patients for subtotal thyroidectomy by a surgeon experienced in this operation, especially if the goitre is large or unsightly. Solitary toxic nodules and toxic intrathoracic or sublingual goitres are best treated surgically; the last two lesions may swell during exposure to radio-iodine.

There are several relatively minor disadvantages of treatment with radio-iodine. First, the patient must be

admitted to a radiotherapeutic unit for about one week. Secondly, it is hard to estimate the dose precisely. This is attempted by formulae based on (a) the amount of a preliminary tracer dose of radio-iodine taken up by the gland, and (b) the weight of the gland, which cannot be determined with any accuracy. The ideal dose should result in neither myxoedema nor recurrence of thyrotoxicosis; yet in practice the incidence of these sequelae is quite high.

The outcome depends partly on how evenly the dose is taken up by the gland, for areas of thyroid tissue which are inactive at the time of treatment may later hypertrophy and cause toxicity. This is especially so with toxic nodular goitres, and these may need repeated treatments with radio-iodine. Thirdly, radio-iodine corrects toxicity more slowly than thyroidectomy or antithyroid drugs: at least three months must elapse before the patient becomes euthyroid. Severely toxic patients, and those with advanced thyrocardiac disease, cannot safely wait that long and will need preliminary treatment with antithyroid drugs.

Treatment with radio-iodine is justified only exceptionally in patients under 45, the main indications being recurrence of thyrotoxicosis after subtotal thyroidectomy, and severe cardiac or other associated diseases.

PALLIATIVE AND SYMPTOMATIC TREATMENT

Sedation

Until thyrotoxicosis has been corrected, most patients will benefit from sedation and a hypnotic at night. In doubtful or borderline cases it is wise to defer definitive treatment until the outcome is clearer; little is to be lost from a period of observation and sedation with amylobarbitone mg. 15, two to three times a day, combined with a sympathetic and reassuring psychological approach.

Malignant Exophthalmos

Several hormones or their synthetic analogues have been used in attempts to suppress the pituitary, the *vis a tergo* believed to be responsible for this dreaded complication. Whether the pituitary factor is overproduction of thyrotrophic hormone, or, as seems more likely, of an associated exophthalmic substance (E.P.S.), is not yet clear. The unpredictable natural history of the complication makes the task of therapeutic assessment very difficult. Testosterone and oestrogens have been abandoned, and thyroxine in large doses now has few advocates, in the treatment of the condition. It is, however, generally agreed that, when severe exophthalmos threatens, any sudden reduction or overcorrection of thyrotoxicosis is dangerous, presumably because it stimulates further secretion of thyrotrophic hormone or E.P.S. The patient needs repeated reassurance and should be kept at a mildly hyperthyroid level by careful undertreatment with antithyroid drugs for at least six months, or until the outcome is clearer. Some consider the addition to the regimen of laevo-thyroxine 0.4 mg. a day to be a prudent safeguard against the risk of therapeutic hypothyroidism. If the patient has in fact been allowed to become hypothyroid, then the rapidly acting thyroid hormone, tri-iodo-thyronine, should be prescribed in doses of 0.04 mg. twice a day for two to three weeks until laevo-thyroxine has had time to take effect.

If exophthalmos rapidly worsens, despite careful management, there should be no delay in applying such measures as lateral tarsorrhaphy, radiotherapy to the orbits or pituitary gland, and, as a desperate sight-saving measure, orbital decompression (Naffziger's operation). Corticosteroids equivalent to a daily dose of 100 mg. cortisone may assist in reducing the orbital oedema. The eyes should be protected with oily drops and shielded by dark glasses.

Cardiac Complications

The most common cardiac complications of hyperthyroidism are auricular fibrillation or flutter and congestive failure; these are the reactions of an ageing heart to the long-sustained demands of increased cardiac output. Other lesions of the heart may be present coincidentally. Thyrotoxic heart disease steadily increases in incidence from middle age onwards. Recognition of it is a signal for early treatment, for its reversibility depends to some extent on the duration of symptoms.

Until their thyrotoxicosis has been corrected, patients in congestive heart failure should be treated by bed rest and the appropriate cardiac therapy, depending on the severity of the condition. Those with uncomplicated auricular fibrillation will require digitalization only if the heart rate is rapid. If the cardiac rhythm does not revert to normal very shortly after hyperthyroidism has been corrected, the use of quinidine should be considered. Severely toxic patients may respond poorly to digitalization, but overdosage must be avoided.

Muscle Wasting

In the older age-group thyrotoxicosis occasionally presents with severe wasting of muscles, simulating myopathy or myositis. In this event, one of the new anabolic steroids, such as "durabolin" (19-nortestosterone phenyl propionate), 25 mg. once a week by intramuscular injection, may assist return of strength, alongside radical treatment.

Glycosuria

Transient glycosuria is not uncommon in thyrotoxic patients. It may be due to a lag in storage of glucose consequent on rapid absorption from the gut, or to unmasking of a mild latent diabetes by increased metabolism. The condition usually responds to treatment of the underlying disease. If diabetes is severe, or persists after treatment of the thyrotoxicosis, it should be controlled by the usual measures.

Thyroid Crisis

This dreaded and often fatal complication is rarely seen these days. It erupts when the circulating thyroid hormone reaches a critically high level in patients already exhausted from their disease. Most often it is precipitated by surgical manipulation of a toxic goitre inadequately prepared for operation. There is no way, unfortunately, of neutralizing thyroid hormone once it has escaped into the blood-stream, and its degradation is very slow. All that can be done is to prevent further release of the hormone by promptly giving iodide in large doses, and protecting against its effects by digitalization, sedation, and corticosteroid therapy; for there is some evidence that adrenocortical exhaustion may be an important factor in these cases.

ASSESSMENT OF APPLICATIONS FOR REHOUSING ON MEDICAL GROUNDS

BY

H. D. CHALKE, O.B.E., T.D., M.A., M.R.C.P., D.P.H.

R. W. WATTON, M.B., B.S., D.P.H., D.C.H.

Public Health Department, Borough of Camberwell, London

AND

B. H. PENTNEY, M.R.C.S., L.R.C.P.

General Practitioner, Herne Hill, London

The housing problem in certain areas is still a long way from solution. It has considerable public-health implications, and it places heavy responsibilities on the medical officer of health, who must assess the degree of priority to be awarded to applicants furnishing medical certificates in support of their rehousing claims. The number of these certificates has increased steadily in the London area since the war. Many patients believed that the greater the number of certificates the more priority was obtained and the quicker the rehousing. Though this was not so, it was difficult to explain the real situation to them. The variety of certificates submitted was considerable and their value and accuracy were sometimes obviously doubtful. It was clearly necessary for some regulation to be introduced to produce conformity and a substantial measure of fairness in allocating priority, and also to allow the family doctor a chance to give his objective recommendation confidentially. Argles (1950) drew attention to the increase in the number of rehousing applications; he showed that, whereas in 1938 the number received by the L.C.C. was 1,000, by 1949 (first nine months) it had risen to nearly 23,000. He observed that some applicants used every means in their power to secure accommodation and that the medical certificate was the most favoured method of gaining some preference.

Between 1955 and 1958 the Borough of Camberwell received 3,235 applications for rehousing; during the same period 1,623 housing applicants claimed medical priority. The applications were accompanied by certificates from the family doctor and to a lesser extent from hospital doctors and almoners.

In a review of 1,028 applications for rehousing, supported by medical certificates, submitted to Hampstead Metropolitan Borough, Warren (1954) found that 25% related to respiratory infections, 15% to rheumatic disorders, and 14% to psychological disorders. He stressed the need of the medical officer of health, in awarding "medical points," to have detailed information about the housing and medical conditions before he can allocate points.

After consultations with general practitioners and hospital staffs it was decided to introduce in Camberwell a new system that should prove more equitable. It was agreed that the form of medical certificate for housing should be the same for all cases and that only one certificate should be submitted for each patient unless the medical condition changed.

The Certificate

In 1955 the accompanying housing form was devised and brought into use.