movements depends less upon the active efforts of the patient, and it is therefore more suitable in the elderly. The results of the operation might be better if it were not so commonly applied to the grossly disabled patient.

None of these operations will enable a patient to escape from hospital much before three months have elapsed. In the elderly, time is of the greatest importance, and any procedure which entails a shorter stay in hospital demands consideration. This is the merit of the replacement arthroplasty. It must be accepted, however, that no prosthetic replacement of the femoral head in the arthritic hip has yet been shown to give a durable, stable, painless joint. To some extent this must be due to a failure of materials and design. Mere replacement of the femoral head, however, cannot be expected to relieve all pain and restore all movement. It may, however, provide an acceptable compromise between the disadvantages of the stiff hip and the perils of the formal arthroplasty.

These patients were under the care of Mr. J. S. Batchelor and Mr. T. T. Stamm. It is a pleasure to express my gratitude to them.

REFERENCES

- Adam, A., and Spence, A. J. (1958). *J. Bone Jt Surg.*, **40B**, 219.
 Adams, J. C. (1953). *Ibid.*, **35B**, 199.
 Apley, A. G., and Denham, R. A. (1955). *Ibid.*, **37B**, 195.
 Batchelor, J. S. (1948). *Postgrad. med. J.*, **24**, 241.
 Brown, A. (1954). *Guy's Hosp. Rep.*, **103**, 13.
 Gade, H. G. (1947). *Acta chir. scand.*, **95**, Suppl. 120.
 Judet, J., and Judet, R. (1950). *J. Bone Jt Surg.*, **32B**, 166.
 McMurray, T. P. (1939). *Ibid.*, **21**, 1.
 Osborne, G. V., and Fahrni, W. H. (1950). *Ibid.*, **32B**, 148.
 Pridie, K. H. (1955). *Ibid.*, **37B**, 224.
 Shepherd, M. M. (1954a). *Ibid.*, **36B**, 355.
 — (1954b). *Ibid.*, **36B**, 567.
 Smith-Petersen, M. N. (1939). *Ibid.*, **21**, 269.
 — (1948). *Ibid.*, **30B**, 59.
 Stinchfield, F. E., and Cavallaro, W. U. (1950). *Ibid.*, **32A**, 48.
 Wardle, E. N. (1955). *Ibid.*, **37B**, 568.
 Watson-Jones, R., and Robinson, W. C. (1956). *Ibid.*, **38B**, 353.

GERIATRIC SURGICAL EMERGENCY

BY

D. E. BOLT, M.B., F.R.C.S.

*The West Middlesex Hospital, Isleworth**

In general, geriatric medicine may be regarded as embracing all patients over 50 years (Thewlis, 1946), but the major problems of surgical management in the geriatric patient are not seen with any frequency until a much greater age. By the age of 70, however, few patients fail to show unmistakable signs of physical or mental degeneration. It was therefore decided to review all the patients of 70 years and over admitted as emergency cases to a single surgical unit over a period of four years. It is not suggested that there is any essential pathological difference between the surgical diseases of patients under and over 70, but the treatment of these diseases in the aged often calls for some modification of standard practice, and geriatric surgery resolves itself into a series of compromises between the optimum surgical treatment required by the disease and the needs of a patient whose capacity to withstand treatment is rapidly diminishing. To emphasize this aspect of the management of the geriatric surgical emergency, a system of classification has been used which compares the programme of treatment provided for the patient with that which would have been offered to a fit man of 50 years with comparable pathological condition.

*Now at St. Mary's Hospital, London, W.2.

Classification

Group 1: Optimum Management.—*a*, No operative treatment required. *b*, Full optimum operative treatment provided.

Group 2: Compromise Management.—*a*, Emergency surgery not required. Unfit for desirable elective surgery. *b*, Emergency or palliative surgery provided. Unfit for desirable elective surgery. *c*, Emergency or palliative surgery provided. Lesion otherwise inoperable by ordinary surgical standards.

Group 3: Unsatisfactory Management.—Unfit for emergency or palliative surgery.

Present Series

During the four-year period from August, 1954, to August, 1958, 408 patients of 70 years or over were admitted as emergency cases to a unit of 70 beds, or were transferred to it as emergencies from medical beds. The details are summarized in Table I. It will be seen

TABLE I.—Summary of Whole Series of 408 Cases

Group	No. of Cases	Male	Female	Deaths		Average Age in Years	Average Days in Hospital	No of Cases with Medical Complications
				No.	%			
Misdiagnoses ..	35	24	11	7	20	77	12	—
Strangulated hernia ..	73	38	35	9	12	79	12	28
Malignant obstruction ..	23	13	10	11	48	73	28	8
Simple ..	20	10	10	6	30	78	21	7
Appendicitis ..	29	12	17	6	21	77	23	11
Diverticulitis ..	16	10	6	1	6	72	19	4
Biliary disease ..	37	17	20	5	14	77	13	15
Perforated peptic ulcer ..	8	6	2	2	25	86	27	5
Other gastric lesions ..	16	7	9	4	25	77	26	5
Pancreatitis ..	6	3	3	2	33	73	10	1
Retention of urine ..	118	118	0	34	29	77	32	70
Vascular lesions ..	10	6	4	8	80	77	13	5
Miscellaneous ..	17	10	7	9	53	75	14	7
Total (excl. 35 misdiagnoses) ..	373	250	123	97	25	77	22	166 (45%)

Total number of operations = 185. Operative deaths = 54 (29%).

that a group of 35 cases is described as misdiagnoses and is excluded from the total analysis. These cases were admitted with a surgical diagnosis but subsequently proved to have no surgical disease and were either discharged or transferred to medical units. Of these 35 patients, 8 were admitted for intestinal obstruction but were proved to have constipation or faecal impaction, 3 were sent in for retention of urine because of the oliguria of congestive cardiac failure, 12 diagnosed as cases of perforated peptic ulcers had either coronary occlusions or exacerbations of peptic ulcers, while the remainder included such diverse conditions as retention of urine in a woman with diabetic polyneuritis, a myeloid leukaemia with abdominal pain, a case of Paget's disease of bone, and a patient with an aneurysm of the thoracic aorta. The mortality of this group during an average stay of 12 days was 20%, making an interesting comparison with the surgical conditions summarized in the rest of Table I.

A few other general comments are necessary. The ratio of 250 male cases to 123 female almost exactly reflects the proportion of 2 male beds to 1 female bed which exists in this unit. The male predominance was due almost entirely to cases of urinary retention, so that in other sections of the table the sex incidence is probably a reasonably accurate reflection of the natural ratio for the disease concerned. The deaths were those occurring in hospital from any cause, and, while it was

not thought worth while to complicate the table with separate operative and non-operative mortalities in each condition, the total number of operations and the operative mortality are included in the final summary.

Turville and Dripps (1948) record 1,064 anaesthetics given to patients of over 70 with 4.9% mortality. Against this a mortality of 29% among patients operated upon may sound a discouraging figure, but the present series, exclusively composed of emergency cases, includes a high proportion of major procedures, and takes account of all deaths in hospital whatever the cause. Moreover, it implies that 70% of patients aged 70 to 90 years subjected to major operations, often in the presence of important medical complications and under emergency conditions, survived to leave hospital. The average stay in hospital was 22 days. This figure includes those who died, as well as those discharged, so that the time in hospital must be considered in conjunction with the mortality for the corresponding condition, as a high mortality tends to shorten the average length of stay.

The medical complications referred to in Table I are those present on admission and not any developing in hospital. Minor disorders were disregarded and medical complications were included only if their severity would significantly influence the approach to the case. The chief conditions found were hypertension (with pressures above 180/100), chronic bronchitis and emphysema (with notable dyspnoea and productive cough), congestive cardiac failure, fibrillation, aortic or mitral valvular disease, previous coronary occlusion or effort pain, previous cerebrovascular accident, paralysis agitans, and major arthritis, especially if treated with steroids. In all, such complications were present in 45% of the cases, confirming the view of Cates (1959) in haematemesis, that in elderly patients associated conditions are as great a hazard as the primary disease.

In the general surgical management of these patients the investigation and treatment have been restricted to the essential minimum. Major dehydration or electrolyte disturbance may make intravenous correction unavoidable, but "it is better for intravenous injections to err on the side of giving less than giving more" (Elman, 1952). While the risks of dehydration are obvious in the elderly, the effect which drips produce—of tethering in immobility a patient already reluctant to move and of depressing his spirits with discomfort and anxiety—is insufficiently regarded. Rectal or subcutaneous routes for fluid administration are often effective, and at times a compromise with ideal fluid-balance requirements in the interest of an active patient may be well worth while. The amount of good which can be done for most of the medical complications during the brief period available before an emergency operation is small, but rapid digitalization of a fibrillating patient in early congestive failure may make an impossible risk eminently possible. Where more time can be gained, chronic bronchitis and congestive failure may be greatly improved. The place of antibiotics in the treatment of chest complications before and after operation is obvious, but of equal importance is a nursing staff trained to insist on frequent breathing exercises and coughing. Finally, in maintaining the will to live, which is so important in the elderly, there is no substitute for the mixture of sympathy and cajolery provided by an experienced ward sister.

C

The need for skilled anaesthesia by staff experienced in managing elderly patients cannot be overemphasized. Such a service has been constantly available to the cases in this series, and those rejected for operation have been rejected by the surgeon rather than by the anaesthetist. Criteria for assessing suitability for anaesthesia broaden rapidly when working among this type of patient. Where advantage can be taken of local anaesthesia, it is often wise to do so, strangulated hernias, in particular, lending themselves to surgery under a local block. The mortality does not seem to be much affected, but the convalescence after local anaesthesia appears to be smoother and quicker, possibly because local anaesthetics encourage speed and gentleness in the surgeon, qualities of particular importance to elderly patients. These patients do not stand prolonged operations well, especially under emergency conditions, and there is no substitute in their management for expedition on the operating-table.

Strangulated Hernias

The 73 cases (38 male, 35 female) had an overall mortality of 12%. There were 29 inguinal, 38 femoral, 4 umbilical, 1 incisional, and 1 obturator among them. In eight no operation was performed; of these, one with sloughing colon in an umbilical hernia was moribund on admission, one difficult incisional hernia was effectively reduced, and six inguinal hernias in unfit patients were successfully treated by taxis and a truss. For patients with inguinal hernias undergoing operation it was a frequent practice to obtain consent for orchidectomy in advance, as the speed and security of surgical treatment was increased by removal of the testis, especially in fat patients or in those with hernias with large sacs.

The advantages of a virtually subcutaneous operation easily performed under local anaesthesia with minimum trauma to muscle appear to outweigh the more direct access to the neck of the sac provided by the Lotheissen operation, especially in elderly patients. When resection was essential, and general anaesthesia was strongly contraindicated, it was quicker to supplement the low approach by a small paramedian incision rather than to disrupt the inguinal canal, even this being possible, with great gentleness, under local anaesthesia. Patients needing operation for their hernias under local anaesthesia were seen in advance by the anaesthetist and premedicated and prepared for operation as if for a general anaesthetic, so that in cases needing resection a change of method could safely be introduced during the operation. Among the 65 cases operated upon there were eight bowel resections with two deaths (25%), while 26 cases were done wholly under local anaesthesia and 37 had all or a major part of their operation under general anaesthesia. All the femoral hernias were explored initially by the low route, and the approach was modified only if bowel required resection.

Malignant Intestinal Obstructions

All 23 patients in this group were suffering from carcinoma of the colon or rectum, and only one, who was moribund on admission, was refused any surgery. In 15 patients only palliative surgery by colostomy or ileotransverse anastomosis was performed, but the remaining seven had initial decompression, subsequent resection, and anastomosis, and, where appropriate, closure of colostomy. Of the 22 patients operated upon, 10 died (45%), of whom seven were among the palliative

operations (48%) and three among the radical procedures (43%), a noticeably similar result with each form of treatment. The summary of the form of treatment compared with the optimum appears in Table II,

TABLE II.—*Comparison of Treatment Given with Theoretical Optimum*

Group:	1a	1b	2a	2b	2c	3
Strangulated hernia ..	—	65	7	—	—	1
Malignant obstruction ..	—	7	—	—	6	1
Simple ..	—	14	3	2	—	1
Appendicitis ..	—	24	4	—	—	1
Diverticulitis ..	7	—	1	7	—	1
Biliary disease ..	—	6	25	4	—	2
Perforated peptic ulcer ..	3	4	—	—	—	1
Other gastric lesions ..	—	10	—	4	—	2
Pancreatitis ..	2	4	—	—	—	—
Retention of urine ..	13	75	—	14	—	16
Vascular lesions ..	—	5	—	2	—	3
Miscellaneous ..	1	7	2	1	1	5
Total ..	26	221	42	43	7	34
Percentage of whole series (373) ..	7	59	11	11.5	1.9	9

and it should be noticed that in six patients given only palliative treatment the tumour would have been regarded as inoperable in a 50-year-old man, but that in nine cases only palliative surgery was offered to patients with resectable growths because of age or infirmity.

Simple Intestinal Obstruction

The results in this group are particularly disappointing. Of 20 patients (10 male, 10 female), one was moribund on admission, but 16 of the remainder were operated upon, with five deaths, a mortality of 31%. The cases undergoing operation included eight adhesion obstructions, two of which required gut resections, three internal hernias, one intussusception of ileum due to a lipoma, and one gall-stone ileus. The remaining six cases had volvulus of the sigmoid colon, a condition treated conservatively by tube decompression when diagnosed (Bolt, 1956). In three of these, however, laparotomy was performed, one having a Mikulicz operation and the others a simple untwisting of the volvulus with a temporary sigmoid colostomy.

Though the overall mortality of 30% was well below that for malignant obstruction, deaths in the simple cases are clearly of even more moment than in the malignant cases. These patients suffer from the general reluctance to diagnose emergency abdominal conditions in the elderly, consequently they come in too late, with a degree of distension and dehydration which makes subsequent management very difficult. When patients present early and can be operated upon promptly without long periods of drip and suction, even very elderly patients do remarkably well. The mortality quoted by Cooke (1958) for simple intestinal obstruction at any age was 21%, with which a mortality of 30% in patients over 70 years of age does not compare too unfavourably, but both figures suggest the paramount need in these cases for early surgical treatment.

Appendicitis

Of the 29 patients in this group (12 male, 17 female), six died (21%). One of these was considered unfit for any surgery, one died while undergoing conservative treatment for an appendix abscess, and the remaining four died after operation, an operative mortality of 17%. Three patients who recovered from appendix abscesses under conservative management were not advised to have elective surgery later, as the risk of operation was thought to exceed the risk of

recurrent appendicitis during their life expectancy, and so far none has returned with a recurrence. Of the 29 cases, 20 (69%) were perforated upon admission: the perforation rate among unselected appendicitis admitted to the unit is 17%. The high rate of perforation agrees with the experience of Christensen (1958) and reflects both the trivial signs and symptoms produced in the elderly by major abdominal pathology and the general reluctance to diagnose appendicitis in old people. The teaching that acute appendicitis is rare in the aged is inaccurate and should be abandoned.

Diverticulitis

There were only 16 cases of diverticulitis in the series, a number too small to be significant. This may explain the low mortality of 6%, which does not correspond with the general experience that diverticulitis provides some of the most difficult problems of management encountered in emergency surgery. Of the 16 cases, seven had local inflammatory masses in the left iliac fossa or pelvis resolving on conservative management, five had peritonitis or an abscess, two had vesico-colic fistulae, and two had intestinal obstruction. The analysis of the type of treatment offered, compared with the optimum, is to be seen in Table II, where the emphasis on palliation at this age is evident. Seven patients treated conservatively and offered no definitive surgery had lesions which might reasonably have been treated on similar lines in young patients, but one patient in group 2a who had no emergency treatment should optimally have had a cold resection subsequently, while the seven in group 2b would not have been left with permanent colostomies at 50. Probably no two surgeons would make quite the same choice of patients for resection in diverticulitis, and the classification consequently reflects the practice of the unit, but very few patients over 70 have a physique which justifies elective resection of the colon for a non-malignant condition, especially as the stronger the indication for resection the greater the technical difficulty.

Biliary Disease

Most acute biliary conditions may be regarded as urgencies rather than emergencies, but they are commonly admitted as emergencies, as were the 37 cases considered here, of which 29 had acute cholecystitis or biliary colic, and eight had obstructive jaundice. It has not been the practice of the unit to indulge in emergency cholecystectomy, certainly not at this advanced age, and emergency operation was performed in only four cases, three because of deterioration on conservative management, and one because of a diagnostic error. In each case cholecystotomy, with removal of stones, was performed, and one patient died (25%). The majority of patients treated conservatively for acute cholecystitis had subsequent cholecystography, but it has not been the practice to offer elective cholecystectomy for proved stones in this age group. Successful cholecystectomy has occasionally been performed for a patient with severe recurrent symptoms, but the frequency of serious recurrence has not been sufficient to suggest that the need for surgery usually outweighs the high operative risk. Lee (1958) suggests that gall-stones *per se* are not a danger to life nor an indication for cholecystectomy, and in the higher age groups this appears to be true, as the chance of further major complications is not great within these patients' expectation of life.

Of the eight patients with obstructive jaundice an exploratory operation was performed on five—four while jaundiced and one subsequently—and all had stones removed from the common bile duct, with or without cholecystectomy; two died, giving an operative mortality of 40%. The remaining three included one patient with carcinoma of the pancreas dying of liver failure without surgery, and two patients with unmistakable obstructive jaundice, resolving spontaneously, who were not regarded as fit subjects for elective surgery. The mortality over the whole biliary group was 13.5%, but the operative mortality was 30%, reflecting the fact that the least favourable cases eventually demand surgery.

Perforated Peptic Ulcer

There seems at the moment to be a noticeable decline in the incidence of perforation of peptic ulcers, and the condition is becoming a relative rarity among abdominal emergencies. Consequently there were only eight cases in this group—three gastric ulcers, four duodenal ulcers, and one an anastomotic ulcer following gastro-enterostomy. The average age of this group was the unusually high one of 86 years, and two patients died, a mortality of 25%. Of the eight cases, one was moribund and beyond help, while four were operated upon and had simple closures of their perforations with one death. The remaining three were treated conservatively despite clear evidence of perforation, including free gas beneath the diaphragm in x-ray films, and all these did well. There seems to be a particularly useful place for conservative management of elderly patients with perforated peptic ulcers who are bad operative risks. In cases otherwise suitable for non-operative treatment liberal free gas has not appeared a contra-indication: a woman of 92 years, with each half of her diaphragm harbouring large amounts of gas, did particularly well on a medical regime.

Other Gastro-duodenal Emergencies

There were 16 cases in this group, nine with haemorrhage, and seven with pyloric obstruction. Of the nine with haemorrhage, seven had emergency partial gastrectomy, one who was bleeding from an anastomotic ulcer after gastro-enterostomy and whose duodenum appeared normal had his ulcer excised and the stoma closed, while the other died without becoming fit for surgery. Of the seven patients with pyloric stenosis, four were due to carcinoma and three to simple ulcers; three of the carcinoma patients had partial gastrectomy, and the three with simple stenoses had gastro-enterostomy, the remaining patient dying without surgery. The total mortality for the whole group was four (25%), but there were only two deaths after 14 operations, a mortality of 14%.

Pancreatitis

There were only six cases of acute pancreatitis, with a mortality of 33%. In two cases in which a firm diagnosis was made without operation the serum amylase was convincingly high, but as both patients recovered satisfactorily the diagnosis was not absolute. The remaining four were diagnosed at laparotomy and no further manoeuvre was performed; but, despite this, two died—an operative mortality of 50%.

Retention of Urine

This constitutes the largest single group of cases, 118 in all, and presents a wide variety of problems in management. Many were admitted in a hopeless condition, the retention arising as a terminal event in a serious medical illness and presenting problems of management insuperable at home. They were then admitted to a surgical ward as cases of retention of urine, and help to explain the 60% incidence of medical complications in this group on admission, and a mortality of 29% for the group as a whole compared with 15% among those subjected to major surgery.

Of the 118 cases, 88 had benign prostatic enlargement, 25 had carcinoma of the prostate, four had strictures, and one had carcinoma of the bladder. Sixteen were treated by catheterization only and either succumbed rapidly to associated medical disease or, on becoming ambulant again, after recovery from medical conditions were able to pass urine effectively. None of these was suitable for elective prostatectomy, and when able to pass urine satisfactorily they were discharged. In 13 of the cases with carcinoma of the prostate, catheterization followed by stilboestrol and eventual removal of the catheter proved adequate, and they were discharged with adequate micturition for subsequent follow-up; in two of them stilboestrol was supplemented by castration. In nine cases transurethral resection followed by stilboestrol re-established micturition, but in three others who failed to respond to hormone therapy, even transurethral surgery was considered too hazardous and permanent suprapubic drainage was performed.

Apart from those treated solely by catheter drainage, 11 benign obstructions were regarded as fit only for permanent suprapubic cystostomy. This decision was taken with reluctance, and only after a considerable period of catheter drainage and energetic treatment of associated medical disease. Consequently it is unlikely that any of them will subsequently be considered fit for major surgery, and, in fact, no second-stage operation has been done in the whole series. Major prostatic surgery has been performed on 64 of the benign cases, 52 having retropubic prostatectomies, two having transvesical operations because of associated bladder disease, and 10 transurethral resections, using the McCarthy resectoscope. In the early cases of the series, low spinal anaesthesia was supplemented with light general anaesthesia, but, more recently, epidural anaesthesia similarly supplemented has been used consistently and found highly satisfactory. The details have been discussed elsewhere by Morris and Candy (1957). The operative mortality in these cases, together with the nine carcinoma patients having transurethral resections, was 11 in 73 cases, or 15%.

It has not been the practice to do emergency prostatectomies. The opposing views on this subject are well known, but we have been influenced both by the high rate of associated disease in these patients, demanding pre-operative treatment, and by the belief that a few days in the ward under supervision, growing accustomed to staff, surroundings, and routine, benefit these elderly and often feeble patients more than does the avoidance of pre-operative catheterization. More leisurely pre-operative assessment and corrective measures greatly benefit them, and it is not apparent that catheter drainage makes an important contribution to mortality or morbidity in this series, or that major

surgery, precipitously arranged at inconvenient hours, would add much to the patients' prospect of recovery.

Vascular Lesions

These cases are included in the interest of completeness. Of the 10 patients, four had major limb gangrene on admission, and all died, with or without amputation, one had mesenteric occlusion, and three had leaking aortic aneurysms, and all died. The only survivors were two patients who underwent local amputation of gangrenous toes.

Miscellaneous

This group allows no useful analysis, as it contains such varied conditions as bleeding colonic neoplasms, polyposis of the ileum, and a wide variety of abscesses, the total of 17 cases including nine operations with four deaths and an overall mortality of 53%.

Adequacy of Treatment

The comparison of treatment provided with the theoretical optimum is summarized in Table II. It will be seen that 26 patients (7%) needed no operative treatment, and that a further 221 (59%) received operative treatment which could be regarded as optimum for their condition, so that a total of 247 cases (66%) received fully adequate treatment. It would perhaps be legitimate to add to these the seven cases of group 2c who were inoperable on pathological grounds, so as to complete the group in whom age and associated disease did not prevent an adequate programme of treatment—68% of the total. In 85 patients (23%) palliation only was possible, and elective surgery was considered to be unjustifiable on grounds of age or general health, while a further 34 (9%) had to be rejected even for emergency measures. Thus in 119 cases (32%) the programme of treatment was notably inadequate to deal with the known pathology.

Clearly, the decision on whether to accept or reject a particular patient for radical surgery is a very difficult one, and is often best taken after assessment of the patient by physician and anaesthetist, as well as surgeon, and a joint discussion of the findings. The surgeon will be influenced by his personal experience of the disease in question, the facilities at his disposal, the views expressed by his colleagues on the general health of the patient, and by his own assessment of the patient's morale. Consequently, different surgeons would select differently, and a ratio of one patient in three rejected for radical surgery may be thought to err towards the conservative, but it is also an illustration of the increasing place of the elderly and unfit among the general surgeon's patients and of the difficult decisions involved in giving them an adequate surgical service.

Summary

A study is presented of 408 emergency surgical cases in persons over 70 years of age admitted to one surgical unit during a four-year period.

The cases are reviewed in disease groups, and the management and outcome are discussed.

A system of classification comparing treatment provided with a theoretical optimum is described and applied.

My thanks are due to Mr. W. J. Ferguson, under whose care all these patients were admitted, for permission to make and publish this study, and to Dr. M. W. Warren,

consultant geriatrician at the West Middlesex Hospital, for much helpful advice. I am also indebted to the records officer and his staff at the West Middlesex Hospital for their great help in tracing records.

REFERENCES

- Bolt, D. E. (1956). *Brit. J. Surg.*, **44**, 172.
 Cates, J. E. (1959). *Brit. med. J.*, **1**, 206.
 Christensen, E. (1958). *Ibid.*, **2**, 832.
 Cooke, R. V. (1958). *Proc. roy. Soc. Med.*, **51**, 503.
 Elman, R. (1952). In *Cowdry's Problems of Ageing*, 3rd ed., edited by A. I. Lansing, p. 868. Williams and Wilkins, Baltimore.
 Lee, A. E. (1958). *Brit. J. Surg.*, **45**, 523.
 Morris, D. B., and Candy, J. (1957). *Brit. J. Anaesth.*, **29**, 376.
 Thewlis, M. W. (1946). *The Care of the Aged*, 5th ed., p. 34. Mosby, St. Louis.
 Turville, C. S., and Dripps, R. D. (1948). *Penns. med. J.*, **51**, 434.

HYPOGLYCAEMIA IN METASTATIC FIBROSARCOMA OF THE LIVER

BY

G. A. BOUSVAROS,* M.D.

From the Department of Clinical Therapeutics, University of Athens Medical School, Athens, Greece

The association of primary carcinoma of the liver with hypoglycaemia was originally observed by Nadler and Wolfer (1929). The subject has since been carefully reviewed by McFadzean and Yeung Tse Tse (1956), who reported nine cases of this disorder in a personal series of 25 patients with malignant hepatoma. Reports on the association of hypoglycaemia with secondary tumours of the liver are rare. Wilder (1927) mentioned two cases, without giving details. Bielschowsky (1932) observed hypoglycaemia in a patient with hepatic metastases from carcinoma of the stomach, and Irsigler (1934) found increased insulin sensitivity, but no frank hypoglycaemia, in patients with liver metastases from cancer of the rectum. Himsworth (1943), in a review of hypoglycaemia, mentioned secondary carcinoma of the liver as one of the causes without giving other details. Hines (1943) reported on a patient with a retroperitoneal liposarcoma who had hypoglycaemic attacks which ceased when the sarcoma was removed and recurred a few years later when hepatic metastases developed. Among Mellinkoff and Tumulty's (1952) series of 20 patients with hepatogenic hypoglycaemia were two with metastatic liver cancer and biliary cirrhosis due to obstruction by the primary pancreatic tumours. The metastases were small and the diffuse cirrhotic process was obviously responsible for hypoglycaemia. Holten (1957) described the case of a patient with a metastatic spindle-cell sarcoma of the liver who had hypoglycaemic attacks five years after removal of the primary growth.

The following case was one in which extensive metastatic involvement of the liver from a gastric fibrosarcoma gave rise to recurrent hypoglycaemic attacks.

Case Report

This 70-year-old man was admitted to hospital in coma on November 23, 1957. He was a known diabetic and had been treated with daily insulin for six years. For four months he had been showing increasing anorexia, weakness, tiredness, and loss of weight, and his insulin requirements decreased, 20 units of protamin insulin being given on alternate days instead of every day. His last injection had

*British Council Scholar, Postgraduate Medical School, London, W.12. Now at the Cardiac Department, Guy's Hospital, London.