REFERENCES

- Altschule, M. D. (1949). *Physiology in Diseases of the Heart and Lungs*. Harvard Univ. Press, Cambridge, Mass.
- Barach, A. L., and Richards, D. W., jun. (1931). *Arch. intern. Med.*, **48**, 325.
- Berger, E. Y., Galdston, M., and Horwitz, S. A. (1949). *J. clin. Invest.*, **28**, 648.
- Case, R. B., Berglund, E., and Sarnoff, S. J. (1955). *Amer. J. Med.*, **18**, 397.
- Chart, J. J., Gordon, E. S., Helmer, P., and Le Sher, M. (1956). *J. clin. Invest.*, **35**, 254.
- Drury, D. R., Henry, J. P., and Goodman, J. (1947). *Ibid.*, **26**, 945.
- Fishman, A. P., Maxwell, M. H., Crowder, C. H., and Morales, P. (1951). *Circulation*, **3**, 703.
- Love, A. H. G., Roddie, R. A., Rosensweig, J., and Shanks, R. G. (1957). *Clin. Sci.*, **16**, 281.
- Moltke, E., and Worning, H. (1958). *Acta med. scand.*, **160**, 397.
- Motley, H. L., Cournand, A., Werkö, L., Dresdale, D. T., Himmelstein, A., and Richards, D. W., jun. (1948). *J. Amer. med. Ass.*, **137**, 370.
- Richards, D. W., jun., and Barach, A. L. (1934). *Quart. J. Med.*, **27**, 437.
- Werkö, L. (1947). *Acta med. scand.*, suppl. 193.
- Whittenberger, J. L., and Sarnoff, S. J. (1950). *Med. Clin. N. Amer.*, **34**, 1335.
- Wilson, R. H., Hoseth, W., and Dempsey, M. E. (1955). *Ann. intern. Med.*, **42**, 629.

POLYMYOSITIS

REPORT ON THREE CASES FROM WEST AFRICA

BY

B. G. BARNARD, M.R.C.P.

A. M. RANKIN, M.R.C.P., M.R.A.C.P., D.T.M.&H.

Senior Registrars, Department of Medicine, University
College Hospital, Ibadan

AND

J. H. ROBERTSON, M.D.

Registrar, Department of Pathology, University
College Hospital, Ibadan

The term "polymyositis" was first introduced by Wagner (1863) when he reported a case of acute generalized muscular affection with skin involvement which ended fatally within six days. Potain (1875) described a similar case in which the illness was less acute but which terminated fatally as a result of bronchopneumonia four to five months after the onset of the disease. Hepp (1887) gave a detailed account of the clinical features of a similar disorder, which he called pseudotrichinosis, and, while there were no skin lesions, the picture was that of a subacute polymyositis. Unverricht (1877) stressed the almost invariable occurrence of skin lesions in these cases, and therefore applied the name dermatomyositis to the condition. Subsequently he described the pattern of muscular involvement and pointed out that not all cases were fatal (Unverricht, 1891).

The first reported cases of this disease from Great Britain came from Gowers (1899) and Batten (1899), and though changes occurred in both skin and muscle in their patients they still used the term polymyositis to describe the condition. Oppenheim (1899, 1903) drew attention to the occasional involvement of mucous membrane and both ocular and cardiac muscle. He pointed out that the skin lesions could be sclero-dermatous in character. Petges and Cléjat (1906) also stressed the extreme degree of atrophy and sclerosis of the skin (poikiloderma) that could occur, and described a chronic form of the disease.

It would appear from the above that much uncertainty has existed in the past about the relationship between dermatomyositis and other diseases of muscle without skin lesions, and only recently has it become established that dermatomyositis can occur without recognizable skin lesions. It is the purpose of this paper to describe three such cases, which represented clinically the acute, the subacute, and the chronic stages of the disease. All three cases occurred in Africans and are thought to be unique, as a review of the literature has shown that, apart from a single case in a Eurasian from South Africa (Sament and Klugman, 1957), the condition has so far not been reported in an African from this continent.

Case 1

A girl aged 16 was admitted to hospital with a five-days history of swelling and slight weakness of her arms and legs associated with pain on movement. The day after the onset of these symptoms she experienced some difficulty in swallowing, and this was followed the next day by the development of a nasal intonation to her voice.

At the time of examination her temperature was 100° F. (37.8° C.), and a striking clinical feature was swelling which involved particularly the arms and to a less extent the legs. The swelling in the upper limbs was confined to the forearms, while, in the lower limbs, the calves were mainly affected and the thighs only slightly so. The soft palate moved poorly but was otherwise normal, the gag reflex was elicited with difficulty, and water dripped from the patient's nose when drinking. A thorough and systematic examination of the central nervous system did not reveal the presence of any abnormalities, and the other systems were also normal.

In view of the findings on admission a tentative diagnosis of polymyositis was made, and this was later confirmed histologically.

Routine investigations were all normal, except Hb, 8.6 g./100 ml.

A biopsy specimen taken from the extensor muscles of the left forearm showed groups of muscle cells to be swollen, with intensely eosinophilic cytoplasm and a loss of their transverse striations and sarcolemma nuclei. The degeneration was patchy and was associated with a slight inflammatory infiltration composed mainly of lymphocytes and plasma cells.

As the patient was not acutely ill on admission, no specific therapy was given, pending the muscle-biopsy report. A week after admission, however, her temperature began to rise, the swelling of the arms increased, and the difficulty experienced in swallowing was greater. Assuming that the diagnosis was correct, prednisolone, 20 mg. eight-hourly, along with tetracycline, 250 mg. six-hourly, was given, the latter drug for controlling the bronchopneumonia which had developed. In addition, a Ryle tube was passed and intragastric feeding begun. In spite of these measures she steadily deteriorated, and finally died from extensive bronchopneumonia 16 days after admission.

Post-mortem Findings

The body was that of a well-nourished young female. The subcutaneous tissues of the arms and legs were the seat of a gelatinous oedema which was especially marked in the forearms. The underlying muscle was pale. Recent adhesions were present in both pleural cavities, and on the left side there was about half a pint (280 ml.) of straw-coloured effusion. Examination of the viscera showed extensive bronchopneumonic consolidation of the lower lobes of both lungs. The liver was the seat of a moderate degree of fatty infiltration and the spleen was enlarged, weighing 215 g. No other abnormal findings were observed macroscopically, and urine taken from the bladder showed no evidence of myoglobin on spectroscopic examination.

Sections were taken from the intercostal muscles, vastus lateralis, supinator of the forearm, diaphragm, soft palate, and oesophagus. While these varied in appearance, degeneration of the striated muscle fibres was prominent and affected small groups or even single fibres scattered unevenly throughout the sections. The cells were swollen and eosinophilic, they had lost their striations, and their sarcolemma nuclei were either shrunken or had disappeared. Elsewhere the fibres had become fragmented and were being phagocytosed by histiocytes. In other parts lesser degrees of damage were indicated by vacuolation and variation in the size of the muscle fibres. These changes were prominent in all the sections, except in those from the soft palate, where fibrosis was pronounced but where the remaining muscle appeared healthy. Inflammatory infiltration varied in degree in the muscles examined, and while virtually absent in the supinator muscle (Fig. 1) it was quite

FIG. 1.—Case 1. Supinator muscle. Groups of swollen, deeply staining muscle fibres are present. There is no inflammatory reaction. (H. and E. $\times 96$.)

pronounced in the striated muscle of the upper part of the oesophagus, the features here being very similar to those in Fig. 4. The inflammatory cells comprised plasma cells, lymphocytes, and histiocytes, with only occasional polymorphonuclear cells. In none of the sections did the small blood-vessels show any abnormalities.

Sections of the skin from forearm and thigh showed thickening of the dermis, both by oedema and by an increase of its collagen. A mild plasma-cell and lymphocytic infiltration was present and the hair follicles and sebaceous glands were atrophic. There was no evidence of fibrinoid degeneration of collagen, and the dermal vessels showed no abnormal features.

In the heart, the muscle bundles were separated by oedema fluid and there was a diffuse interstitial infiltration by mononuclear cells with some polymorphs. Changes in the cardiac muscle, however, were inconspicuous.

The liver showed periportal fatty infiltration, and deposits of malarial pigment were present in the spleen.

Case 2

A man aged 50, a Government official, was admitted to hospital for the investigation of symptoms of breathlessness with undue fatigue on exertion and loss of weight, these symptoms having begun three months earlier. In addition, he also complained of a slight cough which on occasion proved troublesome at night. His previous health had been good and his family and social history revealed nothing of relevance.

On examination he was pyrexial, with an increased respiratory rate, poor chest expansion, and bilateral basal crepitations in the lungs. No abnormal signs were found in any of the other systems. His blood-pressure was 120/70.

Routine investigations revealed nothing abnormal except (1) the E.C.G. showed some flattening of the T waves, and (2) chest x-ray examination showed bilateral basal bronchopneumonic changes.

Over the next few weeks he complained of increasing weakness and fatigue. His temperature varied from 98 to 100° F. (36.7 to 37.8° C.), rising on occasion to 101 to 102° F. (38.3 to 38.9° C.). Addison's disease was considered in view of an equivocal Kepler water-excretion test. However, treatment with cortisone, 12.5 mg. twice a day, produced no improvement.

One month after admission, when a firm diagnosis had still not been reached, the patient volunteered the information that he had been unable to raise his arms above his head while having his chest radiographed. In view of this a more thorough examination was made of the patient's musculo-skeletal system along with a further examination of the central nervous system; the following were the results.

There was an obvious degree of wasting and atrophy of the muscles of the shoulder-girdle which extended downwards to involve the proximal portions of the arms, these changes being more pronounced on the left side. The wasted muscles were weak, and in the upper limbs the movements most affected were abduction of the shoulder and extension of the forearm. In the lower limbs obvious wasting was not seen, but there was a marked diminution of power in the muscles of the pelvic girdle and thigh. In addition the muscles of the trunk were very weak, great difficulty being experienced by the patient in raising himself up in bed from the supine position. The neck muscles were also weak, the head tending to be held in the flexed position owing to the weakness of the extensors. Other muscles similarly affected were the intercostals, this resulting in respirations that were shallow and rapid and in the main diaphragmatic. Chest expansion was less than $\frac{1}{2}$ in. (2 cm.). No fasciculation was observed in any of the muscles, and neither myotonic nor myasthenic reactions were elicited. Muscle-swelling and tenderness were absent. The tendon jerks, though present, were diminished. Both plantars were flexor. Sensation, both superficial and deep, was intact throughout.

A few days later the patient noticed that if he drank fluids too quickly they trickled back through his nose. A re-examination of the cranial nerves at this stage showed that these were all normal and that both the soft palate and the pharynx showed no evidence of paralysis and appeared to move well. The cough reflex, on the other hand, though normal-sounding in character, was definitely diminished in intensity and ineffectual.

In view of the above and the finding of a urinary creatine excretion of 590 mg. in 24 hours, it was thought that he was

FIG. 2.—Case 2. Deltoid muscle. At the top left and extreme bottom right, degenerate muscle fibres are being invaded and phagocytosed by histiocytes. Also, towards the right, hyperplastic sarcolemma cells are arranged in short columns. (H. and E. $\times 111$.)

probably suffering from a disease which involved muscle both primarily and extensively, and on these grounds a tentative diagnosis of polymyositis was made.

A biopsy specimen taken from the deltoid muscle showed the features of a subacute polymyositis (Fig. 2). There was a variation in size of the fibres, some of which were swollen and eosinophilic, with loss of their striations and sarcolemma nuclei. In several places it was seen that the change affected only a short segment of the fibre. Elsewhere necrotic fibres were being invaded by plasma cells and histiocytes, and undergoing phagocytosis and replacement by fibrous tissue. In some areas small basophilic fibres were prominent. These contained columns of large proliferated sarcolemma cells, the appearance being that which has been ascribed to muscle-cell regeneration.

Case 3

A woman aged 55 was admitted to hospital with a five-years history of increasing weakness of her arms and to a less degree of her legs, resulting in some difficulty in walking.

She also complained of symptoms suggestive of a peptic ulcer. Her health prior to the onset of the original illness had been good. Her family history and obstetric history were also of no significance. She had had five pregnancies, all of which were normal, and the children were alive and well.

On examination she presented as an elderly adult in whom the most striking clinical feature was a marked degree of wasting of the muscles of the shoulder-girdle (Fig. 3). The quadriceps femoris of both lower limbs was also atrophied. The atrophy was symmetrical in distribution, muscle tenderness was absent, and no fasciculation was observed in the muscles involved, or elsewhere. Power was greatly diminished in the deltoids and triceps on both sides, the patient being unable to abduct the shoulder or extend the elbow. It was also diminished, but to a less degree, in the biceps and the quadriceps femoris, these muscles appearing to be wasted

FIG. 3.—Case 3. Showing wasting of muscles of the shoulder-girdle.

rather than weak. A moderate degree of weakness in the trunk muscles was also noted. The tendon reflexes were absent in the arms and decreased in the legs. Both plantars were flexor and the abdominal reflexes were present. Sensation was intact throughout. Passive movements in all joints were painless and full in range.

Routine investigations revealed nothing abnormal. A barium-meal examination showed the presence of a large duodenal ulcer. The E.C.G. was normal. Lumbar puncture yielded normal fluid.

A biopsy specimen taken from the deltoid muscle (Fig. 4) showed degeneration of the muscle fibres with phagocytosis and replacement fibrosis. Extensive infiltration by plasma cells and lymphocytes was present, the appearances being similar to those seen in the oesophagus of Case 1. A skin biopsy of the V area of the chest showed no abnormal features.

Discussion

While it is now generally accepted that an acute type of polymyositis does occur, it is comparatively rare, and the disease more commonly presents in either a subacute or a chronic form. In the three cases

described here, we were fortunate enough to see what we considered to be, on clinical grounds, all three types of the disease.

Case 1, though terminating fatally from an aspiration bronchopneumonia within three weeks of the onset of symptoms, offered little difficulty in diagnosis, for the patient presented with the characteristic features of the acute form of the disease—namely, fever, generalized muscle pain, and weakness with subcutaneous oedema. Dysphagia and regurgitation of fluids down the nose were also prominent symptoms, and sections taken from the oesophagus and the palate at necropsy showed both these structures to be involved in the disease process. No obvious skin lesions were seen at any stage of her illness, yet sections of the skin examined after death showed thickening of the dermis by oedema and an increase of its collagen content in addition to a mild round-cell and plasma-cell infiltration.

In retrospect it is felt that this patient's life might well have been prolonged, and perhaps a remission obtained, if a tracheotomy had been done soon after admission, as bronchopneumonia was undoubtedly the main cause of death in her case.

The clinical presentations of Cases 2 and 3 were in sharp contrast to that of Case 1, and they illustrate the wide range of clinical picture that may occur. It is in these subacute and chronic forms of the disease that difficulties in diagnosis may arise unless the protean nature of the symptomatology is recognized. Thus in the second patient exertional dyspnoea, cough, and basal crepitations in the lungs at first suggested a primary cardiac or respiratory disorder, though no evidence of cardiac enlargement was present radiologically. On recognition of the muscular weakness, however, it became clear that his symptoms were due to weakness of the respiratory muscles, resulting in an increase in respiratory rate and an accumulation of secretions in the lung bases, and also that his asthenia was similarly due to more generalized muscular involvement.

Case 3 presented with yet a different clinical picture, the salient features being chronic wasting and weakness of the proximal limb-girdle type, in which the arms were affected more severely than the legs. The diagnosis in this case was more difficult, and polyneuritis due to nutritional deficiency, which is common in Nigeria, as

FIG. 4.—Case 3. Deltoid muscle. An area of fibrosis heavily infiltrated by plasma cells and lymphocytes is shown. Many of the remaining muscle fibres are degenerate. (H. and E. $\times 152$.)

well as progressive muscular atrophy and degenerative lesions of the cervical spine, had to be considered.

Whereas the acute form of polymyositis presents with constitutional symptoms, muscle pain, and tenderness, these are all strikingly minimal or absent in the subacute and chronic forms of the disease (Eaton, 1954). Others have observed that, though striated muscle anywhere in the body could be involved, characteristically the proximal limb muscles and trunk muscles are chiefly affected (Biernard, 1948; Walton and Adams, 1958). It is this distribution of muscular wasting which has led to confusion with muscular dystrophy, so that muscle biopsy is essential for the diagnosis of most cases of subacute and chronic polymyositis. As the disease involves necrosis of muscle, creatinuria has been found to be a constant feature of it (O'Leary and Waisman, 1940; Jager and Grossman, 1944; Cumings, 1953), and persists as long as the disease remains active.

With regard to the prognosis of polymyositis and dermatomyositis, it is now well recognized that up to 50% of the florid cases terminate fatally, usually from pharyngeal or respiratory paralysis or from associated bronchopneumonia. In many of the others the disease becomes arrested, with varying amounts of residual disability (Oppenheim, 1903; Brock, 1934; O'Leary and Waisman, 1940; Sheard, 1951).

In conclusion, it is felt that rare and fatal conditions which have previously been reported from European communities in the temperate zones can also occur in tropical countries among the indigenous population, and that more and more of these conditions will doubtless be discovered as medical services, with adequate laboratory facilities to establish accurate diagnosis, become more readily available.

Summary

Three cases of polymyositis occurring in West Africans are described and the pleomorphic character of the clinical picture is discussed.

The literature is reviewed. Constitutional symptoms, it is generally agreed, are absent in the subacute and chronic forms of polymyositis and the presence of skin lesions is variable. Spontaneous remissions often occur.

The existence of the disease in West Africa is shown.

REFERENCES

- Batten, F. E. (1899). In *Allbutt's System of Medicine*, 7, 461. Macmillan, London.
- Brock, W. G. (1934). *Arch. Derm. Syph. (Chicago)*, 30, 227.
- Biernard, A. (1948). *Ned. T. Geneesk.*, 92, 3486.
- Cumings, J. N. (1953). *Brain*, 76, 299.
- Eaton, L. M. (1954). *Neurology (Minneapolis)*, 4, 245.
- Gowers, W. R. (1899). *Brit. med. J.*, 1, 65.
- Hepp, P. (1887). *Berl. klin. Wschr.*, 24, 389.
- Jager, B. V., and Grossman, L. A. (1944). *Arch. intern. Med.*, 73, 271.
- O'Leary, P. A., and Waisman, M. (1940). *Arch. Derm. Syph. (Chicago)*, 41, 1001.
- Oppenheim, H. (1899). *Berl. klin. Wschr.*, 36, 805.
- (1903). *Ibid.*, 40, 381.
- Petges, G., and Cléjat, C. (1906). *Arch. derm.-syph. (Paris)*, 7, 553.
- Potain (1875). *Bull. Soc. méd. Hôp. Paris*, 12, 314.
- Sament, S., and Klugman, L. H. (1957). *S. Afr. med. J.*, 31, 430.
- Sheard, C. (1951). *A.M.A. Arch. intern. Med.*, 88, 640.
- Unverricht, H. (1887). *Z. klin. Med.*, 12, 533.
- (1891). *Disch. med. Wschr.*, 17, 41.
- Wagner, E. (1863). *Arch. Heilk.*, 4, 282.
- Walton, J. N., and Adams, R. D. (1958). *Polymyositis*, p. 6. Livingstone, Edinburgh.

A CASE OF PROGRESSIVE HEMIATROPHY PRESENTING WITH SPONTANEOUS FRACTURES OF THE LOWER JAW

BY

PAUL BRAMLEY, M.B., B.D.S., F.D.S.R.C.S.Eng.

AND

ALEC. FORBES, D.M., M.R.C.P.

From the South Devon and East Cornwall Hospital,
Plymouth

Hemiatrophy of the body is a rare recessive hereditary degeneration, which can affect the skin, connective tissue, muscle, bone, and peripheral nerves, and involve many organs and systems. It presents to specialists of widely separated interests, so that none may have a true picture of the disorder. To the neurologist it is progressive facial hemiatrophy, its usual textbook designation, and a rare cause of trigeminal neuralgia, epilepsy, and hemiplegia. To the dermatologist it is morphoea, scleroderma en coup de sabre, and various minor paracentral abnormalities of pigmentation of the hair and skin. To the dental surgeon it is a cause of unilateral malocclusion and dental deformity. To otolaryngology it contributes unilateral deafness and malformation of the sinuses. For gastro-enterologists it may induce dysphagia from unilateral atrophy of the oesophagus. It has also caused hypertension from unilateral kidney disease and unilateral diaphragmatic paralysis. To those interested in medicine as a whole its widespread modes of presentation will be of interest. To these is added another, of interest to surgeons, particularly those specializing in facio-maxillary work. Spontaneous fractures of the lower jaw are usually caused by weakening of the bone from infection or neoplasia. The following case is believed to be the first recorded one where the cause of the fractures was progressive facial hemiatrophy.

Case History

The patient is a widow aged 53, with one son. There has been no left-handedness, tic douloureux, or anything else of note in the family history. There were 11 siblings, three of whom have died of unknown causes.

At the age of 13 she had skin changes which began below the right breast and which slowly spread. Since this time the right arm and breast have always been smaller than the left.

At 33 exophthalmic goitre and pre-eclamptic toxæmia occurred, from which she recovered with termination of the pregnancy. She has had no injury, miscarriages, or blood transfusions. At 43 about four attacks of unconsciousness occurred without warning, in which she fell to the ground. She has bruised herself, but not been incontinent or bitten her tongue in these attacks. There was no observer. About this time Raynaud's phenomenon began, and persisted for some years, but she does not get this now.

At the age of 49, frontal and vertical headaches began, yielding to aspirin and usually associated with nausea and vomiting. These occurred several times a week, usually on waking. They have been less severe and without nausea for about a year. A few months after the headaches began, momentary jabs of severe pain in the right infra-auricular region developed. These radiated up into the temporal scalp