were increased in the first week of illness, were usually maximal during the second week, and returned to normal about five weeks from its commencement. The S.G.O.T. and S.G.P.T. levels ran roughly parallel, the S.G.P.T. exceeding the S.G.O.T. in about two-thirds of the cases.

Transaminase values were below 500 units per ml. in all but two patients; high levels were not necessarily associated with severe illness. Variations in individual patients' clinical state were reflected in alterations of their transaminase levels.

Conventional liver-function tests were less often abnormal and were not affected so early. The zinc sulphate turbidity and serum γ -globulin were sometimes raised some months after recovery, when other tests were normal; this was thought to be related to persisting antibody.

Transaminase determination was helpful in early diagnosis and differential diagnosis, and may assist case management. Our findings confirm that liver involvement is a usual feature of the disease.

We thank Dr. J. Pryse-Davies and Dr. J. Henry Wilkinson for permission to publish transaminase data of Cases 1, 2, and 3 from their paper; Dr. R. I. S. Bayliss, who supervised the care of the nursing staff; the practitioners who referred cases to us; the technical staff of the chemical pathology and haematology departments; and the patients themselves for their willing co-operation.

The Unicam S.P. 500 spectrophotometer used for the transaminase determinations was purchased with the aid of a grant to Dr. J. Henry Wilkinson from the Central Research Fund of the University of London.

REFERENCES

- Bennett, H. D., Frankel, J. J., Bedinger, P., and Baker, L. A. (1950). *Arch. intern. Med.*, **86**, 391.
- Berk, J. E., Shay, H., Ritter, J. A., and Siple, H. (1948). *Gastroenterology*, **11**, 658.
- Cohn, C., and Lidman, B. I. (1946). *J. clin. Invest.*, **25**, 145.
- Davidsohn, I. (1937). *J. Amer. med. Ass.*, **108**, 289.
- Downey, H., and McKinlay, C. A. (1923). *Arch. intern. Med.*, **32**, 82.
- Hsia, D. Y., and Gellis, S. S. (1952). *Amer. J. Dis. Child.*, **84**, 175.
- Jordan, W. S., and Albright, R. W. (1950). *J. Lab. clin. Med.*, **35**, 688.
- Karmen, A., Wróblewski, F., and LaDue, J. S. (1955). *J. clin. Invest.*, **34**, 126.
- King, E. J., and Armstrong, A. R. (1934). *Canad. med. Ass. J.*, **31**, 376.
- Kunkel, H. G. (1947). *Proc. Soc. exp. Biol. (N.Y.)*, **66**, 217.
- Leibowitz, S. (1953). *Infectious Mononucleosis*. Grune and Stratton, New York.
- and Brody, H. (1950). *Amer. J. Med.*, **8**, 675.
- MacLagan, N. F. (1944). *Brit. J. exp. Path.*, **25**, 234.
- (1946). *Ibid.*, **27**, 369.
- Bendandi, A., and Cooke, K. B. (1957). *Clin. chim. Acta*, **2**, 49.
- Malloy, H. T., and Evelyn, K. A. (1937). *J. biol. Chem.*, **119**, 481.
- Mason, W. R., jun., and Adams, E. K. (1958). *Amer. J. med. Sci.*, **236**, 447.
- Nelson, R. S., and Darragh, J. H. (1956). *Amer. J. Med.*, **21**, 26.
- Pryse-Davies, J., and Wilkinson, J. H. (1958). *Lancet*, **1**, 1249.
- Rennie, L. E., and Wróblewski, F. (1957). *New Engl. J. Med.*, **257**, 547.
- Sullivan, B. H., jun., Irely, N. S., Pileggi, V. J., Crone, R. I., and Gibson, J. R. (1957). *Amer. J. digest. Dis.*, **2**, 210.
- Van Slyke, D. D., Hiller, A., Phillips, R. A., Hamilton, P. B., Dole, V. P., Archibald, R. M., and Eder, H. A. (1950). *J. biol. Chem.*, **183**, 331.
- Wróblewski, F. (1958). In *Advances in Clinical Chemistry*, edited by H. Sobotta and C. P. Stewart, **1**, 335. Academic Press, New York and London.
- and LaDue, J. S. (1956). *Proc. Soc. exp. Biol. (N.Y.)*, **91**, 569.

GASTRO-DUODENAL HAEMORRHAGE AS A SURGICAL EMERGENCY

BY

JOHN M. LARGE, F.R.C.S.Ed.

From the Royal Berkshire Hospital, Reading

From this hospital, Bohn (1949) reported his experience of 150 cases of gastro-duodenal haemorrhage treated under his personal care in surgical wards. 135 were of acute or chronic peptic ulcers, 33 (25%) underwent emergency surgery, and the overall mortality was 7.4%.

Since that time changes in policy and organization have taken place, and it seemed timely to survey the situation again. The five years 1953 to 1957 are reviewed in this report. Over this period all cases of gastro-duodenal haemorrhage were admitted as routine surgical emergencies under the care of the surgeon on duty. As a result four consultant surgeons and various registrars have been responsible for the treatment of the patients, in contrast to the earlier series, in which all the cases received the individual attention of a single surgeon specially interested in the problem.

In these five years 551 patients were admitted with gastro-duodenal haemorrhage, but 11 are now excluded because the evidence concerning the haemorrhage was doubtful. Of the remaining 540 cases, 488 were due to acute or chronic ulceration; 97 of these underwent emergency surgery, and the overall mortality was 5.9%. This paper is concerned with an analysis of these statistics, and reviews some aspects of the whole period of 11 years from 1946, during which time 861 cases of haemorrhage from peptic ulcers have been treated.

Statistical Analyses

Of the 540 cases of definite haemorrhage, 488 were diagnosed as due to ulceration. The other cases were diagnosed as follows: hiatus hernia 30, oesophageal varices 10, gastric carcinoma 7, purpura 2, and ulcerated gastric polyp, leiomyoma of stomach, and haemophilia 1 each. The diagnosis of hiatus hernia needs some comment. The high incidence in this compared with earlier series is probably due to increased knowledge of the lesion, and in consequence a routine radiological search is made for it. These cases, like nearly all the others, received medical treatment, and later, when bleeding had ceased, a barium-meal examination was made. The hiatus hernia was not firmly diagnosed, therefore, until some days after bleeding had ceased. In many cases oesophagoscopy was done at this stage, but few were found to have oesophagitis. Whether this indicated that oesophagitis had settled on medical treatment, or that the hernia was a coincidental accompaniment of an acute ulcer, it is not possible to say.

The major problem in all series of this sort is, however, the haemorrhage from peptic ulceration (including both acute and chronic ulcers), and the following tables give data concerning the 488 patients in this group.

Table I shows how the group is divided according to diagnosis and sex, and gives numbers treated surgically, with the mortality rates. The "peptic ulcer" group, in which the site of the ulcer was not known, is a rather unsatisfactory one, especially so because it had a high mortality rate. In all there was a history of chronic ulceration, but the site remained in doubt

TABLE I.—*Diagnosis, Sex, Mortality, and Operative Statistics*

Diagnosis	Total	Sex		Deaths		Operations No.	%	Sex		Operation Mortality
		M	F	No.	%			M	F	
Acute ulcers	155	89	66	6	3.9	24	15.5	10	14	4
Duodenal ..	190	142	48	10	5.3	34	17.9	29	5	7
Gastric ..	95	59	36	5	5.3	34	35.8	23	11	2
Peptic ..	35	29	6	8	22.8	0	0	0	0	0
Post-gastrectomy ..	13	10	3	0	0	5	38.5	4	1	0
Total ..	488	329	159	29	5.9	97	19.9	66	31	13

because the patient was either too old and frail to warrant investigation or failed to attend for radiography after discharge. In those who died, post-mortem examination was not carried out. The experience of others, and our own in previous years, is that haemorrhage from a gastric ulcer carries a higher mortality risk than that from a duodenal ulcer, but in the present series the rates are equal. It may well be, therefore, that the peptic ulcer group were in fact mostly cases of gastric ulceration.

In the post-gastrectomy group no chronic ulcers were found in the gastric remnant, the jejunum, or at the anastomosis, by radiography, gastroscopy, or at operation, and in all cases, therefore, acute ulceration must be presumed. These patients are always referred to hospital with a supposed diagnosis of anastomotic ulceration. It does not seem to be widely known that acute ulceration is quite common in the gastric remnant—perhaps even more common than in the normal stomach.

Table II shows what the possibilities are as regards diagnosis, emergency operation, and death in any individual patient, male or female, admitted with gastro-duodenal haemorrhage. The operative liability is approximately equal between the sexes, but the male mortality risk is higher than the female. This agrees with the findings in other series. Fraenkel and Truelove (1955) found the male : female mortality ratio for chronic ulcers to be 6.1% : 4.3%. Cates (1959) found a ratio for all ulcers of 6.6% : 3.2%. In this series exclusion of acute ulcer cases still leaves the males at a disadvantage, and an analysis by age does not explain the disparity.

TABLE II.—*Diagnosis, Operation Liability, Mortality, by Sexes*

Diagnosis	Males (329)			Females (159)		
	Percentage	Operative Liability	Mortality	Percentage	Operative Liability	Mortality
Duodenal ulcer ..	43.2	} 20-1%	9.4%	30.0	} 19.5%	2.7%
Acute ..	27.1			35.5		
Gastric ..	18.0			22.5		
Others ..	11.7			12.0		

TABLE III.—*Reasons for Surgery*

	No. of Ulcer Cases	Percentage of All Ulcer Cases	Percentage of Operations for Ulcer
To control severe bleeding before or within 12 hours of admission	23	4.7	23.7
Continued to bleed or bled again while on medical treatment	68	13.9	68.9
Moderate bleeding but operation within 12 hours. Much previous trouble with ulcer	6	1.2	7.4

TABLE IV.—*Age-grouping of Emergency Operation Patients*

	<29	30-	40-	50-	60-	70-	80+
Total No. ..	2	5	24	23	25	15	3
Percentage of age group	7.1	10.9	24.5	24.2	20.8	12.8	7.5

In Table III are given the reasons for performing emergency operations, and Table IV shows the age-grouping of the patients so operated upon. It is impossible, in retrospect, to define a "severe" bleed, which is purely a clinical decision based on the history, clinical appearance, blood-pressure, and pulse rate. Continued or recurrent bleeding is more easily defined, though not always made manifest by haematemesis or melaena. A rising or persistently high pulse rate, often associated with a fall in blood-pressure, or a persistently low haemoglobin level in spite of transfusions of several litres of blood, has been regarded as an indication of further bleeding.

Age-grouping of the surgically treated cases shows that the peak is reached in the 40-59 age range. This fits the expected pattern, since younger patients rarely bleed to death from their ulcers and consequently a more strict interpretation of the indications for surgery is applied, while the older age-groups increasingly suffer from cardiac, pulmonary, or other disorders which discourage surgery.

Types of Operation Performed

In all cases of chronic ulceration a partial gastrectomy of either a Polya or a Billroth I type was performed. Where possible the ulcer was resected, but in one patient with a very inflamed and adherent duodenal ulcer a prepyloric section gastrectomy was performed. Subsequently, bleeding continued and could not be balanced by transfusions, with a fatal result. The danger of continued bleeding if the ulcer is left *in situ*, even when the intestinal stream is diverted from it, must be weighed when considering how to deal with a difficult ulcer.

Of the cases classified as acute ulcers 14 were treated by local excision of the ulcer, 6 by partial gastrectomy, and 4 by gastrotomy and, in the absence of any specific bleeding-point, simply closure.

Four of the five post-gastrectomy cases were found to have multiple acute ulcers, and a further resection was performed. In the other case gross obesity and adhesions around the site of operation made full exploration impossible without extensive dissection, and a vagotomy was performed.

Causes of Death

In the series of 488 ulcer cases there were 30 deaths, but, following the example of other authors, one death is excluded because the haemorrhage was a minor incident in an otherwise fatal condition. This patient was already seriously ill with carcinoma of the gall-bladder with gross hepatic invasion. The haemorrhage was from acute gastric ulcers, and was not severe. The remaining 29 died from the following causes:

No Surgery.—(16 patients.) 11 bled to death (3 transfusions unable to restore blood volume; 5 too old and frail to warrant surgery; 3 unfit for surgery owing to obesity and bronchopneumonia); 3 cardiac failure—possibly anaemic in origin; 1 ulcer perforated while on medical treatment—unfit for surgery at any time; 1 died 10 minutes after admission.

After Surgery.—(13 patients.) 5 cardiac failure (2 chronic congestive failure, 3 acute failure—possibly anaemic in origin); 4 bronchopneumonia; 1 diabetes, chronic urinary infection, and obesity; 1 leaking duodenal stump; 1 continued to bleed from D.U. after prepyloric section gastrectomy.

These figures emphasize the importance of intercurrent conditions other than ulceration, resulting in the patient's death. Bronchopneumonia, cardiac

failure, and the frailty of old age appear as major factors in the cause of death in 20 out of 29 fatalities.

It is possible that the six deaths due to acute cardiac failure were avoidable. Insufficient transfusion was probably an important factor in most of them, but some may have been precipitated by over-rapid transfusion.

Table V gives the mortality rate per decade.

TABLE V.—*Mortality by Age-groups, and Proportion of Cases by Age*

	<40	40–	50–	60–	70–	80+
Deaths ..	0	1	3	8	11	7
Percentage of age group	0	1.0	3.2	6.7	9.4	17.5

Proportion of patients: 40 and over 86.4%; 50 and over 68.4%; 60 and over 50.9%; 70 and over 29.0%; 80 and over 7.4%.

For comparison:

Avery Jones (1957) ..	60 and over 40%;	series mortality, 7.9%.
Cates (1959) ..	60 " " 33%;	" " 5.4%.
Ogilvie <i>et al.</i> (1952) ..	60 " " 30%;	" " 5.3%.
Fraenkel and Truelove (1955)	50 " " 65.8%;	" " 5.6%.

Discussion

Comparative Statistics.—As has been pointed out repeatedly in previous papers, the fact that the mortality rate rises steeply with advancing years makes exact comparison between one series and another unrealistic. The best that can be achieved is to state the proportions in the series of patients over certain specified ages, giving due allowance for this in making comparisons. In Table V these proportions are stated and compared with four recently published series. Another factor, even less easy to compare, is the severity of the haemorrhages. It is a policy in this hospital never to refuse admission to a surgical emergency; therefore it is possible that this series contains a higher proportion of minor haemorrhages than would be included by a hospital in an area where practitioners have difficulty in finding beds for their emergencies.

Selection of Patients for Surgery

The rational course to follow in treating gastro-intestinal haemorrhage is undoubtedly to select for operative treatment any patient who is judged to be in real danger of bleeding to death, and to treat all the rest medically. In recent years all published papers have followed this course, but have shown considerable variation in the proportion so selected for operation.

The basis for selection in this series was the same as that applied by Bohn (1949) in the first series reported from this hospital. Briefly, surgery may be indicated: (1) if the haemorrhage is unduly severe (indicating involvement of a major vessel); (2) if bleeding continues or recurs while on medical treatment; (3) if the patient is over 40 years of age; and (4) if the patient gives good evidence of having chronic ulceration.

Clearly these criteria are only indications, and leave wide scope for variation in interpretation and emphasis. In Bohn's series (from 1946 to 1948) 25% were operated on with a series mortality of 7.4%. In the next two years, during which the criteria were interpreted with more emphasis towards surgery in an attempt to improve on these figures, the emergency operation rate was increased to 54%, but the mortality rose to 8.3%. While these figures are not strictly comparable (35% of patients over 60 in the first series and 45% in the second), it is clear that the increased operation rate did not improve the results. In the present series the operation rate has fallen to 20%.

This figure of 20% is still higher than that for other series. The following are corresponding figures from

four recent papers: Avery Jones (1957) 10%, Ogilvie *et al.* (1952) 10%, Cates (1959) 13.5%, Fraenkel and Truelove (1955) 16.4%. In comparing these figures it is important to take into account the age factor. The average age of our patients, and those of Fraenkel and Truelove, seem to have been higher than those of all other series, so that more of our patients would fall into the group in which surgery is indicated.

Selection of the correct time for surgery may also vary considerably. In this paper "immediate surgery" has been classified as within 12 hours of admission, a time chosen to indicate that medical measures had played no real part in treatment. In the 1946–8 series 50% were immediate operations, while in the present series only 24% were so treated, without increasing the mortality rate. It is possible that in the immediate post-war years a higher proportion of really severe bleeding occurred, but it seems likely that the greater part of this fall is due to an increased tendency to wait and see if the bleeding recurs. It suggests that, provided blood transfusion is given to maintain the patient in a fit state for surgery at any moment, this initial wait-and-see policy is not unduly risky. It must, however, be stressed that in cases with a very severe haemorrhage, or with continued or recurrent haemorrhage, a wait-and-see policy might well prove disastrous.

Surgical Treatment of Acute Ulcers

It seems to be a widely accepted practice to perform a gastrectomy in all cases when palpation of the stomach fails to reveal a chronic ulcer or other definite cause for the haemorrhage. Ogilvie *et al.* (1952) state that they do not favour gastrectomy, because an acute ulcer is seldom found.

The experience in this hospital is at variance with this view. Since 1946, in 42 cases in which a gastrotomy has been performed because no chronic ulcer was palpable, a definite acute ulcer was found in 23, leaving the surgeon in no doubt about the cause of the bleeding. In these 23 patients the ulcer was excised down to the muscle coat, the bleeding vessel underrun, and the mucosa sewn over the defect. In no case did further bleeding occur.

It is argued that acute ulcers do not require surgery unless they bleed, and therefore the essential point of surgery is to stop the bleeding. Gastrectomy may be necessary to achieve this, but often it is not. To perform a gastrectomy when a lesser procedure will suffice to stop bleeding would seem quite illogical. Not only is a major operation being performed to remove a lesion which would in almost all cases heal spontaneously, but it does not prevent subsequent acute ulceration.

In none of our locally excised ulcers has chronic ulceration developed later at the site of excision. Possibly in a larger series of cases a few such chronic ulcers would develop, but this seems a weak argument for advising gastrectomy in all cases.

The situation is much more difficult when gastrotomy fails to reveal an ulcer with a definite bleeding-point, especially when the stomach and proximal jejunum are empty of blood. If the indications for surgery were unequivocal, then a gastrectomy is almost certainly justified, but, if not, it may be wisest to admit that an error has been made and close the abdomen.

General Improvement in Results

Since the war there has been a steady improvement in the reported results of treatment. In some series the mortality rate has fallen dramatically (Oxford 19% in

1949, 5.3% in 1955), while in others a low mortality rate has been maintained in spite of an increasing proportion of elderly patients (Avery Jones, 7.9% mortality rate maintained, with an increase from 33% over 60 in the earlier years to 40% later). The figures for the 11 years in this hospital are:

1946-48	..	7.4% mortality	..	35% over 60
1949-51	..	8.3% "	..	45% "
1951-53	..	7.0% "	..	49% "
1953-57	..	5.9% "	..	51% "

There is probably no single explanation for the general improvement. An expanded range of antibiotics and anaesthetics agents has undoubtedly helped, but probably increased understanding of the problem, crystallized by Avery Jones in 1947, has played the most important part.

Summary

Of 541 cases admitted suffering from gastro-duodenal haemorrhage 488 were due to acute or chronic peptic ulceration. Of these 488 cases, 97 were treated by emergency operation. The overall mortality was 5.9%.

The frequency and timing of emergency surgery are discussed. A frequency of 20% was found in this series. Comparing this with previous series from this hospital, it does not appear that the mortality rate will be reduced by resorting to surgery more often than this. The number of "immediate" operations (within 12 hours of admission) has tended to fall in this hospital, but nevertheless the dangers of delay in severe or repeated haemorrhage are stressed.

It is suggested that acute ulceration often occurs in the post-gastrectomy stomach. In this series probably all the post-gastrectomy haemorrhages were from acute rather than chronic recurrent ulceration.

It is argued that gastrectomy for haemorrhage from acute ulcers is often illogical and unnecessary.

I thank Mr. F. H. Aitken Walker, Mr. G. Bohn, Mr. R. G. Reid, and Mr. C. Latto for permission to study the records of their patients and to publish this paper. I am particularly grateful to Mr. G. Bohn whose precepts regarding gastro-duodenal haemorrhage I have attempted to follow. I also thank the records department of this hospital, who sorted out over 2,000 case-histories for my perusal.

REFERENCES

- Bohn, G. (1949). *Brit. med. J.*, 2, 630.
 Cates, J. E. (1959). *Ibid.*, 1, 206.
 Fraenkel, G. J., and Truelove, S. C. (1955). *Ibid.*, 1, 999.
 Jones, F. Avery (1957). *Ibid.*, 1, 719, 786.
 Ogilvie, A. G., Cardoe, N., and Bentley, F. H. (1952). *Ibid.*, 2, 304.

Medical Memoranda

"Contrecoup" Fracture of the Sternum

Fractures of the sternum are rare. Holderman (1928) found 62 cases among 46,237 fractures of all types; an incidence of 0.13%. They are usually caused by direct violence. Fractures of the sternum produced by indirect violence are exceedingly uncommon. They have been produced by muscular action in lifting, coughing, labour, and tetanus. "Contrecoup fractures of the sternum," so called by Tarnowsky (1905), are caused either by falls on the head and shoulders, or by the fall of a heavy object on the back of the cervico-dorsal spine, so that the thorax and neck are brought into forcible ante flexion. The fracture thus sustained in the adult is at or near the junction of the manubrium and gladiolus, the latter overriding the former. Tarnowsky, who added 7 cases

to 20 previously described by Servier (1889), summarized the mechanism producing these fractures as follows.

(1) Falls on the head and shoulders press the ribs forward and upward, the range of motion increasing from the first to the seventh pair. (2) The clavicles may sometimes act as a lever and help to wrench the manubrium from the gladiolus. (3) Intrathoracic pressure at the time of the fall exerts a positive pressure on the thoracic wall. (4) The second costal cartilages act as a wedge tending to separate the manubrium from the gladiolus.

Gurlt (1864), Brunn (1913), and Kirkham (1941) reported fractures of the sternum sustained in a similar manner. Many of the fractures were caused by violent injuries producing associated fractures of the cervico-thoracic vertebrae and ribs.

No associated bony injury was present in the following case. The fracture was produced indirectly by the patient falling on his back, the cervico-dorsal spine being acutely ante flexed.

CASE REPORT

A man aged 43 slipped while descending some stairs and fell backwards. He slid down about 16 stairs on his back, feet foremost. His cervico-dorsal spine was acutely ante flexed partly as a result of the fall and partly voluntarily, to avoid hitting the back of his head on the stairs during the descent.

On admission to hospital on December 21, 1958, he complained of severe pain over the front of the sternum which came on immediately after the fall. He was pale; pulse 76, B.P. 150/100. His breathing was mainly abdominal, and he was holding his head in one position to prevent pulling on the sternum through the sternomastoid. He was a well-built man. Localized tenderness and slight swelling were present over the sternum, just below the sterno-manubrial joint. No obvious deformity of the sternum was seen. There was bruising over the right buttock and left scapula. No abnormality was found in the heart and lungs. X-ray examination revealed a fracture of the sternum 2 cm. below the sterno-manubrial joint, with anterior displacement of the lower fragment. No abnormality was seen in the ribs, dorsal vertebrae, or intrathoracic organs.

He was treated by bed rest and analgesics. No attempt was made to reduce the displacement. Ten days after the accident he was discharged from hospital, and began work five weeks later. X-ray examination six weeks after the injury showed the position of the fracture to be unchanged. Bony union was occurring.

I thank Mr. G. O. Jelly for permission to publish this case.

MILES FOX, M.B., F.R.C.S.,
 Resident Surgical Officer,
 Ancoats Hospital, Manchester.

REFERENCES

- Brunn, W. von (1913). *Zbl. Chir.*, 40, 633.
 Gurlt, E. (1864). *Handbuch der Lehre von den Knochenbrüchen*, vol. 2, p. 264. Grote, Hamm.
 Holderman, H. H. (1928). *Ann. Surg.*, 88, 252.
 Kirkham, J. H. (1941). *Brit. med. J.*, 2, 614.
 Servier (1889). Cited by Tarnowsky (1905).
 Tarnowsky, G. de (1905). *Ann. Surg.*, 41, 252.

The World Health Organization has announced that in 1959 only 72,000 cases of smallpox were reported throughout the world as against 242,000 cases in 1958. These statistics, which do not include mainland China, show that the incidence of smallpox was much lower in India and East Pakistan. (United Nations Information Centre.)