passage of bulky fat-laden stools leads to rapid loss of weight. The absorption of protein and of carbohydrate is also impaired, but to lesser degree than that of fat. In the absence of treatment, vitamin deficiencies, hypocalcaemia, hypoproteinaemia, and anaemia may occur. Gradually, compensatory mechanisms assert themselves and halt this progressive starvation. The lowered body weight permits survival on reduced calorie requirements and reduces the demand for protein replacement; anatomical adaptation involving hypertrophy of the remaining small intestine and its mucosal villi augments absorption; the colon increases its ability to absorb water, amino-acids, and crystalloids; reduction in the rate of gastric emptying allows an increased time for contact between food and mucosal surface. This last form of adaptation is well illustrated by the following case-history.

In a man aged 40, massive intestinal strangulation in an internal hernia necessitated resection of the caecum and of the small bowel to within 3 ft. (91 cm.) of the duodeno-jejunal flexure. He developed the typical starvation-steatorrhoea syndrome. Six months after operation he noticed a sudden and striking improvement in his symptoms coincident with the onset of epigastric discomfort and occasional vomiting of retention type. He gave a previous history of duodenal ulceration and a clinical and radiological diagnosis of pyloric stenosis was made. The addition of a gastro-jejunostomy in this particular patient was clearly undesirable, and he was trained to perform nightly gastric lavage with half-strength saline; on this regime he continued in good health for about eight years. It is highly probable that the pyloric obstruction had improved intestinal absorption by promoting "drip feeding" from the stomach into the residual small intestine. An operation planned to produce a degree of pyloric obstruction in patients with insufficient small intestine is rational and worthy of further study.

Repeated haematological studies in patients having had extensive resection of the distal small bowel may shed light on the disputed site of absorption of vitamin-B₁₂. Recent experimental work in the rat suggests that the ileum is the chief site of absorption of this vitamin and that resection of this part of the gut predisposes to macrocytic anaemia (Booth *et al.*, 1957). The patient whose history is described above had a mild microcytic anaemia, but repeated examination of the peripheral blood and bone-marrow has not revealed any sign of a developing macrocytic anaemia; the entire distal small bowel was removed, and yet, almost nine years later, there is no evidence of vitamin-B₁₂ deficiency, and tests have shown satisfactory vitamin-B₁₂ absorption (intestinal continuity was restored by anastomosing the jejunum to the ascending colon). This may reflect either a species difference or the ability of the jejunum to assume the function of absorbing this vitamin.

Conclusion

While there has been substantial progress in our understanding of the physiology of the small intestine, our knowledge must still be regarded as rudimentary. However, cases showing varying degrees of malabsorption are by no means uncommon and an increasing number of research tools are becoming available for use in their study. The radioactive labelling of such food factors as vitamin B₁₂, iron, and fats, has already led to a fuller understanding of the absorption of these substances from the intestine. Biopsy of the small intestine (Shiner, 1956) permits a histological study of the mucosal pattern in various disease states; if it is employed before, during, and after specific therapy much new information should

be gained. This technique also provides adequate tissue for histochemical and electron-microscopical studies. A readily ingestible pill has now been devised which contains a small radio transmitter, capable of transmitting information about hydrogen-ion concentration, pressure, and temperature during its passage through the small intestine (MacKay and Jacobson, 1958). The challenge is clear; new techniques are available, and intensive study should result in significant progress.

REFERENCES

- Anderson, C. D., Gunn, R. T. S., and Watt, J. K. (1955). *Brit. med. J.*, **1**, 508.
 Atwater, J. S., Butt, H. R., and Priestley, J. T. (1943). *Ann. Surg.*, **117**, 414.
 Booth, C. C., Chanarin, I., Anderson, B. B., and Mollin, D. L. (1957). *Brit. J. Haemat.*, **3**, 253.
 Burge, H., and Clark, P. A. (1959). *Brit. med. J.*, **1**, 1142.
 Cameron, D. G., Watson, G. M., and Witts, L. J. (1949). *Blood*, **4**, 803.
 Connell, A. M. (1957). *Scot. med. J.*, **2**, 162.
 Duncan, I. B. R., Goudie, J. G., Mackie, L. M., and Howie, J. W. (1954). *J. Path. Bact.*, **67**, 282.
 Feggetter, G. Y. (1959). *Proc. roy. Soc. Med.*, **52**, 838.
 Fox, H. J., and Grimson, K. S. (1950). *J. Lab. clin. Med.*, **35**, 362.
 Lundh, G. (1957). *Acta chir. scand.*, **113**, 431.
 MacKay, S., and Jacobson, B. (1958). *Electronics*, **31**, Jan. 3, p. 51.
 Meurling, S. (1953). *Acta Soc. Med. upsalien*, Suppl. 3.
 Poulley, J. W., Fairweather, F. A., and Leeming, A. (1957). *Lancet*, **1**, 406.
 Shiner, M. (1956). *Ibid.*, **1**, 17.
 Watson, G. M., and Witts, L. J. (1952). *Brit. med. J.*, **1**, 13.
 Welbourn, R., Hughes, R., and Wells, C. (1951). *Lancet*, **1**, 939.
 Wollaefer, E. E. (1950). *Postgrad. Med.*, **8**, 251.

POST-PUBERTAL ADRENAL VIRILISM WITH BIOCHEMICAL DISTURBANCE OF THE CONGENITAL TYPE OF ADRENAL HYPERPLASIA

BY

R. V. BROOKS, Ph.D., F.R.I.C.

D. MATTINGLY, M.B., B.S., M.R.C.P.

IVOR H. MILLS, M.D., Ph.D., M.R.C.P.

AND

F. T. G. PRUNTY, M.D., F.R.C.P.

From the Department of Metabolic Diseases, St. Thomas's Hospital, and the Department of Chemical Pathology, St. Thomas's Hospital Medical School, London

That the adrenal cortex could be the seat of lesions resulting in virilizing changes was first established by Bulloch and Sequeira (1905); the cases with which they were concerned were prepubertal. In 1910 Apert reviewed the available material, and developed a classification involving all age-groups which is the basis of that in use to-day. He was responsible for the present term "adrenal virilism," while the term "adreno-genital syndrome" derives from Gallais (1912). The syndrome was intensively studied by Broster and Vines (1933) prior to the second world war. It is apparent that the terminology used in the classification of these conditions has become confused. Thus the term "adreno-genital syndrome" may include cases of virilism arising before or after puberty and cases of the Cushing type (Broster, 1953). On the other hand, Wilkins (1957) properly makes a sharp distinction between the adreno-genital syndrome and Cushing's syndrome, including them under the term "hyper-adrenocorticism."

Though similar clinical changes may be brought about by adrenal cortical tumours or adrenal cortical "hyperplasia" the present paper is restricted to a

discussion of those patients afflicted with the latter lesion. Adrenal virilism, adreno-genital syndrome, congenital pseudohermaphroditism, congenital adrenal hyperplasia, and hyperadrenocorticism have all been used to describe the same condition. Clearly the term "congenital (female) pseudohermaphroditism" should not be used in this restricted sense, since the clinical condition may be due to other causes (Wilkins, 1957).

As a better understanding of the chemical pathology of the adrenal cortex is gradually being gained, the time has come to urge precision in the nomenclature of these conditions. The following clinical classification will be used as a basis for discussion in the present paper.

ADRENO-GENITAL SYNDROME:

1. *Congenital adrenal hyperplasia* (female pseudohermaphroditism or male precocious puberty due to adrenal cortical hyperplasia).

2. *Post-natal adrenal virilism*: (a) Arising pre-pubertally. (b) Arising post-pubertally.

(The adrenals in these patients are often, but not necessarily, enlarged.)

Intensive biochemical studies of a large number of patients of the first group have been made, and it is generally accepted that the ability of the adrenal to synthesize cortisol normally is impaired. This is due to deficiency of 21- or 11-hydroxylation of the steroid nucleus, and results in the excretion of metabolites of accumulated 21-carbon intermediates, such as pregnanetriol and excess of 17-ketosteroids, including metabolites of androgens (Eberlein and Bongiovanni, 1955; Dorfman and Shipley, 1956; Bush *et al.*, 1957; Wilkins, 1957). These biochemical lesions are apparently inborn errors of metabolism and are thought to be inherited as recessive characters (Wilkins, 1957).

By contrast much less is known of the biochemistry of post-pubertal adrenal virilism. To begin with, there is a difference of opinion about the duration of the lesion. Wilkins (1957) implies that an isolated patient developing symptoms at the age of 5 years had a congenital lesion, and makes a similar implication with regard to a patient who reached puberty before she had become markedly virilized (Blizzard and Wilkins, 1957). Greenblatt (1958) is of the opinion that the congenital condition may be latent and have an onset even after puberty. But Dorfman and Shipley (1956) speak of the lesion as being one of "acquired hyperplasia," and Epstein and Kupperman (1959) distinguished a "post-pubertal adreno-genital syndrome" of a different type. Their patients had a late onset and lacked evidence of increased pregnanetriol excretion.

It has been shown that post-pubertal adrenal virilism may be difficult to distinguish by superficial biochemical methods from "simple hirsutism" and indeed from patients with the Stein-Leventhal syndrome. Patients with hirsutism developing post-pubertally showed a continuous spectrum of 17-ketosteroid excretion from normal to moderately high levels (Prunty *et al.*, 1958); out of a series of 32 patients with hirsutism, one was definitely distinguished by the excretion of excessive amounts of pregnanetriol in the urine. Since that time two further cases of a similar nature have been observed, and they are presented here (Cases 2 and 3). Observations on the case previously briefly mentioned are more fully reported here for completeness (Case 1).

Biochemical Methods.—Urinary 17-ketosteroids and 17-ketogenic steroids were determined by methods described in Prunty *et al.* (1959). Pregnanetriol was

determined by a slight modification of the method of Bongiovanni and Clayton (1954). Urine tetrahydrocortisone, tetrahydrocortisol, and 11-oxo-pregnanetriol were determined by chromatographic techniques (Brooks and Prunty, 1960).

Case 1

A girl aged 18 presented with severe hirsutism. This began at the age of 14 and included the face particularly, arms, legs, areolae, and linea alba. Menstruation started at the age of 10 and continued in a regular 21-day cycle of three to four days' duration. Temperature records indicated the periods were ovulatory. The clitoris was moderately enlarged and the mammae were small. She was short in stature—5 ft. 2 in. (157.5 cm.)—not obese, but very broad-shouldered (Fig. 1). There was a moderate degree of acne

FIG. 1.—Case 1. Before treatment. Note broad shoulders, small mammae, and short stature.

on her face and on the upper part of her back and shoulders. Vaginal examination indicated that the uterus and ovaries were of normal size, and endometrial biopsy showed the development of a late secretory endometrium. Mentally she was suffering from considerable tension with hysterical episodes.

The urinary steroid investigations are indicated in Table I. She was treated for eight months with a combination of 10 mg. of prednisone and 0.2 mg. of ethinyl oestradiol daily. (The latter was given in a cyclical fashion for three weeks in every four.) This produced definite improvement

TABLE I.—Urinary Steroid Excretion in mg. Per Day

	Case 1	Case 2	Case 3
17-Ketosteroid	14.8–20.8	27.8–29.4	14.2–17.5
17-Ketogenic steroid	10.6–19.0	10.6–12.2	9.2–19.3
Pregnanetriol	11.7	7.7	3.7
11-Oxo-pregnanetriol	0.9	—	1.0
Tetrahydrocortisone plus tetrahydrocortisol	0.7	—	1.2
After corticotrophin*:			
17-Ketosteroid	38	29	28
17-Ketogenic steroid	48	25	45
Pregnanetriol	54	15	21

* Third day; 20 units in gelatin intramuscularly twice daily.

in the hirsutism. Whereas she had shaved at least once daily and sometimes twice, it was barely necessary to do so once. The acne greatly improved. Her periods continued regularly. The effect of treatment on the urine steroids is shown in Fig. 2. She then refused treatment for 11 months

FIG. 2.—Treatment and steroid excretion of Case 1. Steroid excretions are recorded in mg./day. Horizontal bars for pregnanetriol values represent pooled specimens of urine. The break A indicates an eight-months period during which 10 mg. of prednisone and 0.2 mg. of ethinyl oestradiol were given. The break B represents an 11-months period during which no treatment was given.

FIG. 3.—Case 2. This patient was much less severely affected than Cases 1 and 3.

and the hirsutism relapsed. But on returning to 10 mg. of prednisone with 0.02 mg. of ethinyl oestradiol there was again similar improvement. In addition to reduced growth, the hair became much softer.

Case 2

This girl was also aged 18 and had been hirsute for only four years. She had a hairy elder sister. There was a similar distribution of hair growth to that in Case 1, but the condition was much milder. She was 5 ft. 6 in. (167.5 cm.) tall, the breasts were small, shoulders normal (Fig. 3), and the size of the clitoris was probably within normal limits. Periods were regular, but she had not become pregnant after six months of regular intercourse. X-ray examination after pelvic insufflation of air showed probable ovarian enlargement. Steroid results are shown in Table I. She was treated with 10 mg. of prednisone daily, which was sufficient to cause marked depression of 17-ketosteroids (Table II), but after 13 months the effects on the hirsutism were minimal. Temperature records before and after treatment suggested ovulation was normal.

Case 3

This girl was aged 20 and presented with very severe facial acne of six years' duration. This was followed by the development of severe facial hirsutism and also involved her body as in Case 1. She had two sisters who were not hirsute. She was 5 ft. 4 in. (162.5 cm.), had moderately small mammae, and the clitoris was definitely enlarged. She, too, was broad-shouldered (Fig. 4). Menstruation began at age 12, then stopped for six months. Finally, it became irregular, occurring at intervals of two to three months. The acne was very severe and produced scarring of her face and over her shoulders.

TABLE II.—Urine Steroid Response to Steroid Therapy

	Prednisone Dose (mg./Day)	Duration (Months)	Urine (mg./Day)		
			17-Ketosteroid	17-Ketogenic Steroid	Pregnanetriol
Case 2 ..	10	4 13	5.9 3.6	7.0 10.9	1.3 —
Case 3 ..	20	1	5.5	16.3	—
	15	2	8.3	14.3	—
	6	6	10.3	12.2	—
	20 15	1½ 1	3.0 5.4	11.8 13.8	— —

She was treated with 15 or 20 mg. of prednisone daily for 14 months (Table II). The larger dose was used partly on account of the severity of her condition and partly because there was some evidence that 15 mg./day was insufficient to cause adequate suppression of 17-ketosteroids. The treatment produced a dramatic and sustained effect on the acne, but much less effect on the hirsutism (Fig. 4). Periods became regular at 4-weeks intervals. The relatively large dose of prednisolone produced mooning of the face.

FIG. 4.—Case 3. Note the small mammae and broad shoulders. The pictures were taken after treatment and show the complete quiescence of facial acne with residual scarring.

Discussion

These three patients all developed hirsutism, and two of them acne, after normal menstruation had been established. All had poor mammary development, two clitoral enlargement, and two normal ovulatory periods. One patient was noticeably short and two were broad-shouldered. None had hypertension. The family history varied; one patient had no sisters and there was no history of hirsutism in the family; a second (Case 2) had a hairy sister, but unfortunately it was impossible to obtain a specimen of urine from this girl. The third patient had a normal family history, including two normal sisters.

The 17-ketosteroid levels varied from high normal to moderately elevated, and 17-ketogenic steroids were normal. However, in all three patients the urinary pregnanetriol was elevated, and in the two in whom it

was determined so was the 11-oxo-pregnanetriol. Since the determination of 17-ketogenic steroids includes both metabolites of cortisol and pregnanetriol with its 11-oxo-derivative, the urinary tetrahydrocortisone plus tetrahydrocortisol was determined and shown to be low in both instances. The effect of the injection of corticotrophin was evaluated in all three patients. The ratios 17-ketogenic steroids/17-ketosteroids were low in all cases (0.9–1.6) (Prunty *et al.*, 1958), but for unknown reasons the response to corticotrophin in Case 2 was definitely poor.

The results of therapy in these three patients were variable. Clinically the effect on the acne in Case 3 was dramatic and quite good in Case 1. In no case was the hirsutism abolished, but the alleviation varied from definite improvement in Case 1 to minimal improvement in Case 2. There was a definite tendency in all patients for the growth of hair to be softer, and this alone is often appreciated by patients. In general, the effects of treatment in prepubertal congenital adrenal hyperplasia are often gratifying and the acne and hirsutism which develop may be most favourably affected. Sometimes, however, probably especially if the inauguration of treatment is delayed, the reduction of hair growth is disappointing (Wilkins, 1957; de Mowbray *et al.*, 1959), as it often appears to be in post-pubertal hirsutism (de Mowbray *et al.*, 1959; Jefferies and Levy, 1959; Mattingly *et al.*, 1960) in spite of adequate suppression of 17-ketosteroid excretion by the treatment. This latter, of course, occurred in all three of the above patients (Table II, Fig. 2). In two of them 10 mg. of prednisone daily in divided doses was sufficient for the purpose, but in Case 3 it seemed necessary at times to raise the dosage to as much as 20 mg. daily. This dose is undoubtedly undesirably high and was sufficient to produce evidence of side-effects, whereas 10 mg. daily seems to be a perfectly safe dose.

Reference to Fig. 2 will show that there is a tendency for the administration of oestrogen alone to cause suppression of 17-ketosteroids in Case 1, and possibly also the pregnanetriol excretion. The addition of prednisone, of course, produces a more profound effect, but when the oestrogen is withdrawn on day 20 there is a temporary "rebound" of 17-ketosteroid excretion. Evidence of a similar effect of oestrogen has also been seen during the treatment of a patient aged 13 with congenital adrenal hyperplasia who normally excreted up to 100 mg. of 17-ketosteroid daily. On lowering the dose of cortisone to 37.5 mg. daily (Fig. 5) there was a steady rise in the excretion of 17-ketosteroid, which became markedly reduced on the simultaneous administration of 0.2 mg. of ethinyl oestradiol daily. There can be some similar effect of oestrogen in "simple hirsutism" (Mattingly *et al.*, 1960).

One of the most important aspects of these three cases is the nature of the lesion from which they are suffering. In spite of the late onset of clinical symptoms the biochemical pattern is typical of that of prepubertal congenital adrenal hyperplasia (Wilkins, 1957; Bush *et al.*, 1957; Finkelstein, 1959). Pregnanetriol excretion does not normally exceed about 1 mg./day and 11-oxo-pregnanetriol is below 50 μ g. daily. Pregnanetriol may be elevated somewhat in some cases of simple hirsutism, and especially when corticotrophin is given, but the present cases may be distinguished from these by the relatively higher proportion of pregnanetriol to cortisol metabolites in the urine as reported by Brooks and Prunty (1960). These authors also report more detailed

biochemical studies on these patients, indicating a pattern more typical of congenital adrenal hyperplasia than of simple hirsutism.

Description of patients of this type in the literature is scanty. It is not known what proportion of patients classified as having the adult adreno-genital syndrome by Broster and Vines (1953) have this abnormality of steroid metabolism.

Finkelstein (1956) found 1 out of 11 virilized patients to be of this category, while Greenblatt (1958) has seen four similar patients, three of whom were sisters. This strongly substantiates the latter's view that these cases had a latent form of congenital adrenal hyperplasia. In one of our cases there is a strong probability of a familial background. From the biochemical point of view it is not certain that there is in fact a complete break

between the pattern seen in the more severely abnormal cases of simple hirsutism and the least severely affected of the present cases (Brooks and Prunty, 1960), but there is no more direct evidence, other than the delay of clinical manifestation until puberty, that the lesion may be acquired.

It is not unusual for congenital biochemical lesions to occur in latent form, and there is some evidence that this may be true in parents of children with congenital adrenal hyperplasia (Childs *et al.*, 1956). Hence we are inclined to favour the view that patients with the adreno-genital syndrome of the type described here—that is, clinical post-natal adrenal virilism arising post-pubertally—have basically a congenital biochemical disorder of a type seen in congenital adrenal hyperplasia, where there is relatively greater or less deficiency of the synthesis of cortisol by the adrenal gland, and that this underlying defect could then possibly be brought to light by changing conditions at puberty. It is clear that, for the present at least, this group of patients can only be defined in biochemical terms and distinguished from patients with hirsutism of different origin (Prunty *et al.*, 1958).

Summary

Three cases are presented in which signs of hirsutism and virilism developed after the onset of puberty. This syndrome was accompanied by biochemical changes of the same nature as those found in congenital adrenal hyperplasia developing before puberty.

The urine contained an excess of pregnanetriol, which was accentuated by corticotrophin, excess of 11-oxo-pregnanetriol, and deficiency of metabolites of cortisol.

The nature of this condition and the importance of clearly clarifying it are discussed.

FIG. 5.—Excretion of 17-ketosteroid of a patient aged 13 with congenital adrenal hyperplasia. In the absence of treatment 17-ketosteroids reached 100 mg./day. Note the sharp drop in 17-ketosteroid when treatment with 0.2 mg. of ethinyl oestradiol daily was instituted in July, 1956.

REFERENCES

- Apert, M. (1910). *Bull. Soc. Pédiat. Paris*, **12**, 501.
- Blizzard, R. M., and Wilkins, L. (1957). *A.M.A. Arch. intern. Med.*, **100**, 729.
- Bongiovanni, A. M., and Clayton, G. W. (1954). *Bull. Johns Hopk. Hosp.*, **94**, 180.
- Brooks, R. V., and Prunty, F. T. G. (1960). *J. Endocr.* In press.
- Broster, L. R. (1953). In *The Suprarenal Cortex*, edited by J. M. Yoffey, p. 201. Butterworth, London.
- and Vines, H. W. C. (1933). *The Adrenal Cortex*. Lewis, London.
- Bulloch, W., and Sequeira, J. H. (1905). *Trans. path. Soc. Lond.*, **56**, 189.
- Bush, I. E., Swyer, G. I. M., Stern, M. I., and Willoughby, M. L. N. (1957). *J. Endocr.*, **15**, 430.
- Childs, B., Grumbach, M. M., and Van Wyk, J. J. (1956). *J. clin. Invest.*, **35**, 213.
- de Mowbray, R. R., Spence, A. W., Medvei, V. C., and Robinson, A. M. (1959). *Brit. med. J.*, **2**, 456.
- Dorfman, R. I., and Shipley, R. A. (1956). *The Androgens*. Wiley, New York.
- Eberlein, W. R., and Bongiovanni, A. M. (1955). *J. clin. Invest.*, **34**, 1337.
- Epstein, J. A., and Kupperman, H. S. (1959). *J. clin. Endocr.*, **19**, 1503.
- Finkelstein, M. (1959). *Acta endocr. (Kbh.)*, **30**, 489.
- Gallais, A. (1912). *Le Syndrome Génito-Surrenal*. Thesis, Paris. Reprinted Vigot Frères, Paris. 1914.
- Greenblatt, R. B. (1958). *Recent Progr. Hormone Res.*, **14**, 335.
- Jefferies, W. McK., and Levy, R. P. (1959). *J. clin. Endocr.*, **19**, 1069.
- Mattingly, D., Mills, I. H., and Prunty, F. T. G. (1960). *Brit. med. J.*, **1**, 1298.
- Prunty, F. T. G., Brooks, R. V., and Mattingly, D. (1958). *Ibid.*, **2**, 1554.
- McSwiney, R. R., and Hawkins, J. B. (1959). *A Laboratory Manual of Chemical Pathology*. Pergamon Press, London.
- Wilkins, L. (1957). *The Diagnosis and Treatment of Endocrine Disorders in Childhood and Adolescence*, 2nd ed. Thomas, Springfield, Illinois.

TREATMENT OF SIMPLE HIRSUTISM, INCLUDING THE HIRSUTE TYPE OF STEIN-LEVENTHAL SYNDROME

BY

D. MATTINGLY, M.B., B.S., M.R.C.P.

IVOR H. MILLS, M.D., Ph.D., M.R.C.P.

AND

F. T. G. PRUNTY, M.D., F.R.C.P.

From the Department of Metabolic Diseases, St. Thomas's Hospital, and the Department of Chemical Pathology, St. Thomas's Hospital Medical School, London

The results of diminishing androgen production by steroid therapy or adrenal surgery, in patients with congenital adrenal hyperplasia and adrenal cortical tumour respectively, from the point of view of the hirsutism are variable. They may be dramatic, useful, or even disappointing (Hamilton and Shepardson, 1952; Blizzard and Wilkins, 1957; Heinbecker *et al.*, 1957; Prunty *et al.*, 1958).

In these patients there is usually a clear and major overproduction of adrenal ketosteroids. Occasionally in patients developing post-pubertal hirsutism there is an adrenal lesion similar to that seen in congenital adrenal hyperplasia (Brooks *et al.*, 1960). Such patients have been excluded from the present study. However, most patients suffering from post-pubertal hirsutism do not have these lesions. Many may be described as having simple hirsutism and some have the hirsute type of Stein-Leventhal syndrome with accompanying anovulatory menstrual cycles and sterility. In some of both of these groups abnormalities of adrenal ketosteroids may be demonstrated (Prunty *et al.*, 1958; Gallagher *et al.*, 1958; Brooks and Prunty, 1960).

The present paper concerns attempts at treatment of patients of these two latter types. The effect of prednisone, often combined with ethinyl oestradiol, on the hirsutism of these patients has been observed. While sometimes there is an undoubted favourable response, it must be stated at the outset that it is more often most disappointing. Hirsute patients with the Stein-Leventhal syndrome, the diagnosis being established as described previously (Prunty *et al.*, 1958), have more often been treated by ovarian wedge resection (Stein *et al.*, 1949), since this appears to be the treatment of choice for sterile married women.

Material and Methods

The patients studied included 21 of those described under the heading of simple hirsutism by Prunty *et al.* (1958) and three patients with the Stein-Leventhal syndrome having hirsutism. One of these three patients was included in the earlier paper under the heading of simple hirsutism. The presence of ovaries of the Stein-Leventhal type had been suspected, but at laparotomy they were found to be normal. Two years later she presented with a twisted ovarian cyst, at operation for which the ovaries were found to be typical of the Stein-Leventhal syndrome. Meanwhile she had irregular periods, but temperature records suggested she was ovulating during the few months immediately after the first laparotomy. All these patients were treated with steroids. Their approximate ages at onset of the hirsutism are given in Table I. The precise age at which it occurred is usually difficult to determine.

TABLE I.—Age at Onset of Hirsutism

Age in years No. of patients	11-15 11	16-20 7	21-25 5	26-30 0	31-35 1
---------------------------------	-------------	------------	------------	------------	------------

An additional seven patients with the Stein-Leventhal syndrome were subjected to ovarian wedge resection and not treated with steroids.

Twenty patients received 10 mg. of prednisone daily in divided doses; one received 30 mg., reduced to 20 mg.; one had 15 mg.; and one had 5 mg. In addition, 19 received ethinyl oestradiol in dosage between 0.05 and 0.3 mg. daily for three weeks in every four. Of these, 17 received either 0.1 or 0.2 mg. daily. The duration of steroid treatment varied from 4 to 39 months.

The urinary 17-ketosteroids were usually determined, often with two consecutive days' collection for each determination, to ensure that adequate suppression of 17-ketosteroids was maintained.

The assessment of changes in the hirsutism is not an easy matter. It is naturally most difficult, and undesirable, to prevent patients from removing their facial hair for a sufficient time to make an objective assessment of its rate of growth; the observer must therefore rely on patients' statements about frequency of removal. Bearing this difficulty in mind, patients in category 1 were said to have improved when there was a significant reduction in the frequency of the necessity for removal. This could sometimes be confirmed objectively. Patients who appeared to be certain that the hair, on their limbs particularly, was less virile were placed in category 2. All patients of category 1 also noted this feature. It was usual for all these patients to state that hair had become softer and lighter in colour. Patients in category 3 were those who received no benefit.