haemorrhagic phenomena; and (5) no extramedullary haemopoiesis. In general the anaemia is severe, the haemoglobin level averaging 4.2 g. per 100 ml. in eight cases, the lowest being 1.5 g. per 100 ml.

In the present case there was never any suggestion of myasthenia gravis, nor was there any exposure to marrow toxins or irradiation. The serum proteins were normal, and the histological pattern conformed to that of previously reported cases. The small size of the thymoma, rendering it invisible on an antero-posterior chest radiograph, was a rather unusual feature.

The question whether the thymoma is the cause of the red-cell aplasia or is due to a common aetiological agent has not been settled. There have been four reported cures following thymectomy out of a total of 17 cases operated on (Jacobs *et al.*, 1959), and it has been noted that the tumour has preceded the development of anaemia in six cases. On the other hand, thymomas may be associated with several diverse disorders, including myasthenia gravis, pancytopenia, Cushing's disease, and agammaglobulinaemia, and in one case anaemia developed three years after removal of the thymoma (Clarkson and Prockop, 1958). Furthermore, 15 cases of erythrocytic hypoplasia that did not have thymomas and were not exposed to marrow toxins have been described (Tsai and Levin, 1957). It is possible in this present case that the patient did not live long enough to show any haematological improvement, but in the case reported by Jacobs *et al.* (1959) marrow response was evident before the third week.

Analysis of the cases in which the thymoma was removed surgically shows that four were cured (Jacobs *et al.*, 1959; Humphreys and Southworth, 1945; Chediak *et al.*, 1953; Bakker, 1954) and a further three needed splenectomy or corticotrophin therapy (Chalmers and Boheimer, 1954; Lambie *et al.*, 1957) to effect an improvement. Pre-operatively corticotrophin or corticoids had no effect on the anaemia. Tissue extracts from the thymoma have apparently no haematological effect when injected into prepared animals (Jacobs *et al.*, 1959).

Summary

A further case of thymoma and pure red-cell anaemia is presented. After removal of the tumour there was no improvement in the blood picture up to the time of death 18 days later.

My thanks are due to the Honorary Director of the Special Unit of the Prince of Wales Hospital, Dr. K. W. Starr, for permission to publish this report.

REFERENCES

- Bakker, P. M. (1954). *Ned. T. Geneesk.*, **98**, 386.
 Bayrd, E. D., and Bernatz, P. E. (1957). *J. Amer. med. Ass.*, **163**, 723.
 Chalmers, J. N. M., and Boheimer, K. (1954). *Brit. med. J.*, **2**, 1514.
 Chediak, A. B., Fuste, R., and Rosales, G. V. (1953). *Arch. Hosp. univ. (Habana)*, **5**, 27.
 Clarkson, B., and Prockop, D. J. (1958). *New Engl. J. Med.*, **59**, 253.
 Green, P. (1958). *Canad. med. Ass. J.*, **78**, 419.
 Humphreys, G. H. II, and Southworth, H. (1945). *Amer. J. med. Sci.*, **210**, 501.
 Jacobs, E. M., Hutter, R. V. P., Pool, J. L., and Ley, A. B. (1959). *Cancer*, **12**, 47.
 Lambie, A. T., Burrows, B. A., and Sommers, S. C. (1957). *Amer. J. clin. Path.*, **27**, 444.
 Ramos, A. J., and Loeb, V. (1956). *J. Amer. med. Ass.*, **160**, 1317.
 Seaman, A. J., and Koler, R. D. (1953). *Acta haemat. (Basel)*, **9**, 153.
 Tsai, S. Y., and Levin, W. C. (1957). *Amer. J. Med.*, **22**, 322.
 Weinbaum, J. G., and Thompson, R. F. (1955). *Amer. J. clin. Path.*, **25**, 761.

STERIOD ARTHROPATHY OF THE HIP

BY

D. R. SWEETNAM, M.A., F.R.C.S.

Senior Registrar, Orthopaedic and Accident Department, the London Hospital

R. M. MASON, M.A., D.M., M.R.C.P.

Assistant Physician, Department of Physical Medicine and Regional Rheumatism Centre, the London Hospital

AND

R. O. MURRAY, M.B.E., M.A., M.D., D.M.R.

Radiologist, Institute of Orthopaedics and Royal National Orthopaedic Hospital, London

[WITH SPECIAL PLATE]

Rapid destruction of the femoral head with consequent disorganization of the hip-joint rarely, if ever, occurs in uncomplicated rheumatoid arthritis or osteoarthritis. Recently we have seen four patients, all treated with corticosteroids, in whom such destruction developed. The striking feature in each case was the relative freedom from pain in the presence of severe joint disintegration.

The dangers of steroid therapy in patients with joint infection are well known. Chandler and Wright (1958) were the first, however, to draw attention to similar damage which might occur after intra-articular injection of hydrocortisone in the absence of infection. They demonstrated considerable radiological deterioration, in spite of relief of pain, after injection of hydrocortisone into the knees of patients suffering from rheumatoid arthritis, and concluded that such treatment was "potentially dangerous." More recently, in a comparative study of prednisolone and aspirin in the treatment of rheumatoid arthritis by the Medical Research Council and Nuffield Foundation (1959), it was shown that orally administered prednisolone actually delayed the progressive radiological joint degeneration in the hands and feet of patients suffering from rheumatoid arthritis. The hip, however, seems to react in a different way, for progressive deterioration of an apparently normal joint can occur during steroid therapy. Nevertheless, susceptibility may be increased by pre-existing rheumatoid arthritis or osteoarthritis. Pietrograndi and Mastro-marino (1957) have reported such hip destruction in a woman treated for four years with cortisone by mouth in a dose of 100 mg. daily. A similar change in the osteoarthritic hip of a patient who received 18 intra-articular injections of 50 mg. hydrocortisone at monthly intervals has been reported by Chandler *et al.* (1959). Others have been recorded by Murray (1960) and Sweetnam (1960).

From the relatively small number of similar cases reported in the literature it seems clear that rapid destructive changes may result from either oral or intra-articular steroid therapy. The hip-joint appears to be particularly susceptible, presumably because the enormous force transmitted through it, which often exceeds the total body weight (Denham, 1959), renders it most liable to damage when pain is relieved, while at the same time the bone strength is impaired by local or generalized osteoporosis.

The cases here described are, we believe, further examples of "steroid arthropathy" of the hip.

Case Reports

Case 1

A motor mechanic aged 64 attended the London Hospital in March, 1955, suffering from rheumatoid arthritis of recent onset. He failed to respond to physiotherapy and analgesics, and in August, 1955, treatment with cortisone, 62.5 mg. daily by mouth, was started. Rapid improvement followed and was maintained, but each attempt to reduce the dose was followed by a severe exacerbation of symptoms. The original dose was therefore continued, and, although severely inconvenienced by painful stiffness and recurrent effusions into many joints, he has remained independent.

In February, 1958, after 2½ years of continuous steroid therapy, he complained for the first time of pain in the left hip, and one intra-articular injection of hydrocortisone, 125 mg., was given with some temporary relief. He has been troubled by gradually increasing discomfort in the hip ever since, but the pain has never been severe. During the past four months, however, his walking has been helped by wearing a weight-relieving calliper; he attributes its beneficial effect mainly to a feeling of increased stability at the hip rather than to relief of pain.

Examination revealed the changes of generalized severe rheumatoid arthritis. Movements of the left hip were accompanied by coarse crepitus, but little discomfort except at the limits of movement. The hip flexed to 110 degrees, abducted and adducted 40 degrees, and had a total rotation range of 70 degrees. There was 1½ in. (3.77 cm.) of true shortening, and Trendelenburg's sign was positive on the left side. E.S.R. 29 mm. in 1 hour; Wassermann reaction negative. The latex test for the rheumatoid serum factor was positive in a titre of 1:5,120. Alkaline phosphatase, serum calcium and phosphorus, and plasma protein levels and anti-staphylococcal α -haemolysin titre were all within normal limits. Radiographs showed evidence of generalized osteoporosis. The changes in the left hip in March, 1959, are shown in Special Plate Fig. 1. A radiograph of this hip in April, 1957, showed no abnormality.

Case 2

The headmistress of a small school, a spinster aged 57, developed rheumatoid arthritis when 25 years old. She was severely crippled by joint stiffness and deformity, but with notable fortitude continued to manage her school. During the summer of 1957 what had previously been only discomfort in the left hip slowly gave place to definite pain while weight-bearing; this gradually increased until within a year it was severe enough to interfere further with her already considerable difficulty in walking. At this time (July, 1958) an injection of hydrocortisone, 100 mg., was given into the hip. No relief followed and another injection of 125 mg. was given, but again there was no improvement. In spite of a third injection, in September, 1958, of 30 mg. of prednisolone trimethyl acetate, the pain remained unchanged. No more steroid was given, but the pain has not further increased. During the past six months, however, she has become aware of a feeling of instability in the hip, and she now finds this to be as great an impediment to walking as her pain.

Examination showed the presence of generalized nodular rheumatoid arthritis and multiple severe joint deformities. Movements of the left hip were accompanied by coarse crepitus, but surprisingly little pain. The hip flexed to 120 degrees, adducted and abducted 10 degrees each, and had a total range of rotation of 60 degrees. Throughout this range movement was almost painless. There was 1½ in. (3.77 cm.) of true shortening of the left leg. E.S.R. 15 mm. in 1 hour; haemoglobin 97%; Wassermann reaction negative; differential agglutinin titre (Rose-Waaler test) 128.

The initial rheumatoid change in the left hip as seen radiologically (Special Plate Fig. 2) gave place in nine months to gross destruction of the whole hip-joint (Special Plate Fig. 3).

Case 3

This patient, a ship's steward aged 51, had been in good health until February, 1959. At this time he consulted his doctor about discomfort in both knees, and oral treatment with prednisolone, 5 mg. twice daily, and phenylbutazone ("butazolidin") was prescribed. This relieved the pain, and he remained well, while continuing steroid therapy, until May, 1959. Then without any warning his right hip suddenly "locked," and he was temporarily unable to move it. A radiograph (Special Plate Fig. 4) showed a degenerative arthritis in this joint with partial subluxation and a separate fragment of bone lying lateral to the acetabular margin. The appearance of this fragment suggested that a painless fracture had occurred causing its separation from the acetabulum. There was no clinical or radiological abnormality in the knees. Steroid therapy was stopped in June, 1959, and thereafter pain in the right hip developed with gradually increasing severity. Five months later he was admitted to the Henry Gauvain Hospital, Alton, for a cup arthroplasty to be performed. Examination at that time showed no evidence of generalized arthritis. All movements of the right hip were limited by pain and accompanied by marked muscle spasm. There was a 30-degree flexion deformity with a further flexion range of 20 degrees. Adduction and abduction were negligible, and any attempt to produce rotation was resisted. There was 1 in. (2.5 cm.) of real shortening of the right leg. The Wassermann reaction was negative. By that time he was able to walk only 10 yd. (9 m.) at a time, and the destruction of the hip had become so severe (Special Plate Fig. 5) that simple excision of the femoral head and neck had to be performed.

Case 4

A married woman of 34 was admitted urgently to the London Hospital in July, 1956, suffering from a severe exacerbation of pemphigus vulgaris. The skin eruption was brought under control only by a high dose of steroids, and in order to prevent further relapses an average dose of 20 mg. of prednisone daily was necessary. This has remained a necessity until the present time.

Recently she was referred to the orthopaedic department complaining of a "strange sensation" in the left hip and a feeling as though the left leg was shorter than the right; this had troubled her since January, 1959. She had begun to limp about four months later, and more recently had complained of an occasional dull aching pain in the joint. On examination movements of the left hip produced slight pain at the extremes only. There was no muscle spasm. The hip flexed to 120 degrees and had a total rotation range of 50 degrees. The legs were of equal length.

While the radiograph taken in June, 1957, did not reveal any abnormality, that taken after 2½ years of continuous steroid therapy (Special Plate Fig. 6) shows destructive changes and fragmentation in the head of the left femur with some generalized osteoporosis.

Discussion

We believe that steroid therapy may be responsible for the severity of hip disorganization seen in these patients. It is true that in rheumatoid arthritis and osteoarthritis, in the absence of such treatment, severe destruction of joints can occur. The changes are, however, more gradual and never as extensive, and pain is a prominent feature. Coste *et al.* (1956) have reported a small number of cases of rheumatoid arthritis in which severe destruction of the hips occurred. However, as these cases had also been treated with steroids we believe they are further examples of "steroid arthropathy," and similar to the cases we have described. In the most progressive type of osteoarthritis the changes differ from the severe local disorganization seen in our patients. Though rapid loss of joint space may occur in the early

stages, the pace of destruction soon slows and reactive new bone, both as osteophyte formation and as sub-chondral sclerosis, appears to limit the destruction. Special Plate Fig. 7 shows such a painful osteoarthritic hip-joint at the end of a period of rapid progression in a patient aged 67 not treated with steroids.

Our patients were notable, not only for the rapid destructive changes in the hip-joint, but also for their relative freedom from pain. In one of our cases (Case 3), however, severe hip pain gradually developed when oral steroid therapy was stopped. This suggests that the relief of pain produced by steroids is a significant factor, for it may well be that excessive and unguarded activity in such joints resulting from the relief of pain leads to excessive destruction—a process which in the normal way is held in check by reflex protective muscle spasm.

One patient (Case 4), and probably a second (Case 3), had normal hips before steroids were given, and in these instances it is more difficult to explain why the joints should have suffered in this manner. In Case 4 there was general osteoporosis, and similar changes may also account for the destruction of the femoral head in Case 3. It may be that the mechanism for the production of this arthropathy is a combination of loss of joint pain, osteoporosis, and trabecular stress fractures, which together produce the changes we have described. It is probably significant that very similar changes are seen in patients suffering from *tabes dorsalis*; indeed, the x-ray appearance of the hips in our first three cases is almost indistinguishable from that of a typical "Charcot's joint."

The dose of steroids required to produce destructive joint changes appears to vary greatly in those cases so far reported. It does seem that, when given by mouth, a relatively high dose is required over a prolonged period; though Case 3 is an exception to this, for severe joint disorganization occurred after only 15 weeks' administration of prednisone in a daily dose of 10 mg. Case 4, on the other hand, showed early destructive changes only after continuous oral steroid therapy in high dosage for 3½ years.

The effect of intra-articular administration is just as variable. Case 2 received only three injections into her hip, but nevertheless rapid and severe joint destruction developed. The only other patient recorded in the literature to have developed similar changes after intra-articular injection alone required 18 monthly injections before developing the typical arthropathy (Chandler *et al.*, 1959).

Conclusion

Ever since cortisone and its equivalents became freely available in Great Britain the use of corticosteroids in both hospital and general practice has increased. Clearly, therefore, it is important that the possible damaging effect upon joints should be widely known. It is only recently that such cases have been reported, but already it appears that the condition of steroid-induced joint degeneration is becoming more common. It may well be that as the number of patients receiving steroids increases we shall see more in whom arthritic, and even previously normal, hip-joints have silently crumbled away.

Careful observation of patients on long-term or high-dose steroid therapy is essential so that the early evidence of hip destruction may be detected and treatment adjusted accordingly. Before advising treatment with either oral or intra-articular administration of cortico-

steroids this possible complication should be borne in mind and the likelihood of accelerated joint destruction weighed against the benefit which the patient is likely to derive.

Summary

The development of relatively painless destruction of the hip-joint in four patients treated with corticosteroids is described. "Steroid arthropathy" of the hip is discussed, and the possibility of this complication developing in the arthritic, or even previously normal, hips of patients treated with corticosteroids emphasized.

We wish to express our gratitude to Dr. W. S. Tegner, Mr. C. M. Murray, and Dr. Brian Russell for allowing us to report Cases 1, 3, and 4. Cases 1 and 4 attended the Orthopaedic Department of the London Hospital under the care of Mr. O. Vaughan-Jackson. We should like to thank Mr. R. Ruddick and Mr. R. P. Whitley for preparing the figures. Case 1 was shown to the Orthopaedic Section of the Royal Society of Medicine in October, 1959, and we wish to thank the Honorary Editor of the *Proceedings of the Royal Society of Medicine* for permission to report this case in the present series.

REFERENCES

- Chandler, G. N., Jones, D. T., Wright, V., and Hartfall, S. J. (1959). *Brit. med. J.*, 1, 952.
— and Wright, V. (1958). *Lancet*, 2, 661.
Coste, F., Piguot, B., Delbarre, F., and Saggi (1956). *Rev. Rhum.*, 23, 451.
Denham, R. A. (1959). *J. Bone Jt Surg.*, 41B, 550.
Medical Research Council and Nuffield Foundation Joint Committee on Clinical Trials of Cortisone, A.C.T.H., and Other Therapeutic Measures in Chronic Rheumatic Diseases (1959). *Ann. Rheum. Dis.*, 18, 173.
Murray, R. O. (1960). *Brit. J. Radiol.*, 33, 1.
Pietrograndi, V., and Mastromarino, R. (1957). *Ortop. Traum. Appar. mol.*, 25, 791.
Sweetnam, D. R. (1960). *Proc. roy. Soc. Med.* In press.

VISION, VISUAL ACUITY, AND OCULAR REFRACTION OF YOUNG MEN

FINDINGS IN A SAMPLE OF 1,033 SUBJECTS

BY

ARNOLD SORSBY,* M.D., F.R.C.S.

MICHAEL SHERIDAN, F.B.O.A.

GEORGE A. LEARY, F.S.M.C.

Courage Laboratory, Royal Eye Hospital, London

AND

B. BENJAMIN, Ph.D., F.I.A.

Among the various anthropometric surveys available on the population of this country there is none on the refraction of the eye. Several attempts have been made to establish the distribution of visual acuity, among which the survey on employees in Royal Ordnance Factories during 1943–6 (Black, 1951) was perhaps the most complete. All such studies are, however, open to the objection that they deal with selected material. As for refraction, an acceptable survey requires assessment of the static refraction, and this involves cycloplegia, complicating a fairly tedious task still more. In the present analysis we have attempted to obtain an approach to an unselected series by examining young men called up for National Service. The results recorded here may therefore be taken as nearly representative of young men of 18 to 22 years of age in the general population as could be obtained.

*Working with a grant from the Alexander Pigott Wernher Memorial Trust, Medical Research Council.

D. R. SWEETNAM *ET AL.*: STEROID ARTHROPATHY OF HIP

FIG. 1.—Case 1, March, 1959. Male aged 64 with rheumatoid arthritis. Oral steroid treatment maintained since August, 1955. The severe destruction of the femoral head was accompanied by surprisingly little pain. Radiograph of this hip in April, 1957, showed no abnormality.

FIG. 2.—July, 1958.

FIG. 3.—April, 1959.

FIGS. 2 and 3.—Case 2, female aged 57 suffering from severe rheumatoid arthritis. Three intra-articular steroid injections were given into the hip. No other steroid therapy had been given at any time.

FIGS. 4 and 5.—Case 3, male aged 51 given oral steroid treatment for 15 weeks. In addition to the severe destruction and subluxation of the right hip there was noticeable progression of the arthritic changes in the left hip, which was painless.

FIG. 4.—May, 1959.

FIG. 5.—October, 1959.

FIG. 6

FIG. 7

FIG. 6.—Case 4, November, 1959. Female aged 34 treated with steroids by mouth for three and a half years for severe pemphigus. In spite of this prolonged course the destruction of the femoral head is not yet severe.

FIG. 7.—November, 1959. This patient, aged 67, with painful osteoarthritis was not treated with steroids. Destruction, though rapid at first, soon slowed, and considerable subchondral sclerosis has developed.