thermostable leucocytosis factor, provides a reasonable explanation for the mechanism of leucocytosis in inflammation.

The immunological implication of inflammation is discussed, and it is pointed out that the role of inflammation is that of a regulator of bacterial invasiveness. The pattern of injury in inflammation is ascribed to the liberation of a toxic euglobulin termed necrosin. Fever is shown to be due to two factors—namely, a thermostable component termed pyrexin, and a thermolabile factor.

The studies on the anti-inflammatory problem, first described by me in 1940 and 1942, are summarized; and it is shown that the mechanism of action is referable to a suppression of cellular activity. The injured cell is incapable of forming the mediators of inflammation as readily as under normal circumstances.

A summary of the development of the inflammatory reaction as well as concluding remarks on the significance of the biochemistry of inflammation are pointed out.

REFERENCES

- Bennett, I. L., jun., and Beeson, P. B. (1953). *J. exp. Med.*, **98**, 493.
- Cohnheim, J. (1889). *Lectures on General Pathology*. New Sydenham Society, London.
- Dillon, M. L., Cooper, G. R., and Menkin, V. (1947). *Proc. Soc. exp. Biol. (N.Y.)*, **65**, 187.
- Harris, H. (1954). *Physiol. Rev.*, **34**, 529.
- (1960). *Bact. Rev.*, **24**, 3.
- Hench, P. S., Kendall, E. C., Slocumb, C. H., and Polley, H. F. (1949). *Proc. Mayo Clin.*, **24**, 181.
- King, M. K., and Wood, W. B., jun. (1958). *J. exp. Med.*, **107**, 305.
- Menkin, V. (1931a). *Ibid.*, **53**, 647.
- (1931b). *Ibid.*, **53**, 171.
- (1933). *Ibid.*, **57**, 977.
- (1934). *Amer. J. Path.*, **10**, 193.
- (1935). *Amer. J. med. Sci.*, **190**, 583.
- (1936). *J. exp. Med.*, **64**, 485.
- (1938a). *Ibid.*, **67**, 129.
- (1938b). *Ibid.*, **67**, 145.
- (1940a, reprinted 1950). *Dynamics of Inflammation*. Macmillan, New York.
- (1940b). *Amer. J. Path.*, **16**, 13.
- (1940c). *Arch. Path. (Chicago)*, **30**, 363.
- (1940d). *Amer. J. Physiol.*, **129**, 691.
- (1942a). In *Medico-Surgical Tributes to Harold Brunn*, p. 275. Univ. Calif. Press, Berkeley and Los Angeles.
- (1942b). *Proc. Soc. exp. Biol. (N.Y.)*, **51**, 39.
- (1943a). *Amer. J. Path.*, **19**, 1021.
- (1943b). *Arch. Path. (Chicago)*, **36**, 269.
- (1945). *Ibid.*, **39**, 28.
- (1946a). *Ibid.*, **41**, 376.
- (1946b). *Amer. J. Physiol.*, **147**, 379.
- (1946c). *Arch. Path. (Chicago)*, **41**, 50.
- (1948a). *Blood*, **3**, 939.
- (1948b). *Arch. Path. (Chicago)*, **46**, 145.
- (1949). *Blood*, **4**, 1323.
- (1950a). *Proc. Soc. exp. Biol. (N.Y.)*, **75**, 378.
- (1950b). *Newer Concepts of Inflammation*. Thomas, Springfield, Ill., U.S.A.
- (1951a). *Amer. J. Physiol.*, **166**, 509.
- (1951b). *Ibid.*, **166**, 518.
- (1951c). *Proc. Soc. exp. Biol. (N.Y.)*, **77**, 592.
- (1952). *Arch. int. Pharmacodyn.*, **89**, 229.
- (1953a). *Naunyn-Schmiedeberg's Arch. exp. Path. Pharmacol.*, **219**, 473.
- (1953b). *Brit. J. exp. Path.*, **34**, 412.
- (1953c). *Ibid.*, **34**, 420.
- (1953d). *Proc. Soc. exp. Biol. (N.Y.)*, **82**, 189.
- (1954). *Science*, **120**, 1026.
- (1955). *Ann. N.Y. Acad. Sci.*, **59**, 956.
- (1956). *Biochemical Mechanisms in Inflammation*. Thomas, Springfield, Ill., U.S.A.
- Menkin, V. (1957a). *Amer. J. Physiol.*, **189**, 98.
- (1957b). *J. Physiol. (Paris)*, **49**, 761.
- (1958a). *Amer. J. Path.*, **34**, 921.
- (1958b). *Path. Biol. (Paris)*, **6**, 183.
- (1958c). *Fed. Proc.*, **17**, 448.
- (1959a). Reported at a Symposium on Polypeptides in Relation to Smooth Muscles and Blood Vessels, London, March 23–24, 1959.
- (1959b). *Fed. Proc.*, **18**, p. 494.
- (1960). *Ibid.*, **19**, 124.
- and Kadish, M. A. (1942). *Arch. Path. (Chicago)*, **33**, 193.
- and Warner, C. R. (1937). *Amer. J. Path.*, **13**, 25.
- Miles, A. A., and Wilhelm, D. L. (1955). *Brit. J. exp. Path.*, **36**, 71.
- Shedlovsky, T. (1943). *Ann. N.Y. Acad. Sci.*, **43**, 259.
- Spector, W. G. (1956). *J. Path. Bact.*, **72**, 367.
- Ungar, G. (1953). In *The Mechanism of Inflammation: An International Symposium*, p. 151, edited by G. Jasmin and A. Robert. Acta, Montreal.

NEONATAL HYPERBILIRUBINAEMIA*

BY

ALBERT E. CLAIREAUX, M.D., F.R.C.P.Ed.

Consultant Morbid Anatomist, Bernhard Baron Memorial
Research Laboratories, Queen Charlotte's Maternity
Hospital, London

[WITH SPECIAL PLATE]

Jaundice is commonly observed in newborn infants. It is probable that 40% of infants have a serum bilirubin level of at least 4 mg. per 100 ml. during the first week of life. For this reason the term "physiological jaundice" has crept into use. This is an unfortunate description, as it tends to minimize the risk to the patient. Hyperbilirubinaemia can be dangerous or even lethal in certain patients. The danger lies in the development of cerebral nuclear jaundice (kernicterus). This may either kill the infant or leave him with permanently damaged areas in the brain. It must be stressed, therefore, that in certain circumstances jaundice in the newborn period is to be regarded as pathological and of serious import for the future of the child. Fortunately, hyperbilirubinaemia of this magnitude is not common in the absence of an actual disease process such as haemolytic disease of the newborn. The danger is that the necessary corrective treatment will be instituted too late if jaundice in otherwise normal infants is regarded as being entirely physiological.

The nature of the bile pigment responsible for the hyperbilirubinaemia is of diagnostic and prognostic significance. Ehrlich's method for demonstrating the presence of bilirubin in urine was adapted by van den Bergh and Snapper (1913) for the estimation of bilirubin in plasma. It was later found (van den Bergh and Muller, 1916) that alcohol was not an essential for the development of the colour reaction when bile was used or if the patient was suffering from obstructive jaundice. In patients suffering from haemolytic jaundice alcohol was still necessary, and the reaction was called "indirect." The direct and indirect forms of the van den Bergh reaction have been used for many years in the diagnosis of obstructive and non-obstructive jaundice. It was not clear, however, whether two separate bile

*Read to the Section of Child Health at the Joint Annual Meeting of the British Medical Association and Canadian Medical Association, Edinburgh, 1959.

pigments were involved, or whether bilirubin was in some way bound to protein in haemolytic jaundice and had to be freed by alcohol before a reaction could occur. Some years earlier Schmörl (1904) had examined a number of cases of kernicterus and was struck by the fact that the yellow areas in the brain did not turn green when preserved in formalin, whereas liver tissue in cases of cirrhosis did so. He was among the first to suggest that there were probably two types of bile pigment in the body. It was also observed that neurological complications did not occur in congenital obstruction of the bile-ducts. Cole and Lathe (1953) were able to identify two separate bile pigments in plasma from jaundiced patients. One pigment gave a direct reaction with the van den Bergh test, and was water-soluble. It was present in the plasma of patients with obstructive jaundice, and was the main pigment in human bile. The other pigment was obtained from patients with haemolytic jaundice. This pigment gave an indirect reaction with the van den Bergh test, was insoluble in water, and proved to be bilirubin itself. The direct-reacting pigment was then found to be a glucuronide of bilirubin (Billing *et al.*, 1957). The nature of this pigment was confirmed by Schmid (1956) and Talafant (1957).

Meanwhile Claireaux, Cole, and Lathe (1953) were able to show that the pigment responsible for the yellow staining of areas in the brain in kernicterus was also bilirubin. Samples of brain from the pigmented areas were extracted in solvent systems, and the pigment was separated by reverse-phase partition chromatography. The extracted pigment was identified spectrophotometrically.

The probable metabolic pathways leading to the conjugation of bilirubin with glucuronic acid have been outlined by Billing and Lathe (1958) and Schmid (1958). Uridine diphosphate glucuronic acid and bilirubin are acted upon by an enzyme, bilirubin glucuronyl transferase, with the production of bilirubin glucuronide and uridine diphosphate. The enzyme system responsible for the conjugation of bilirubin is present in the microsomes of the liver cell (Lathe and Walker, 1957). Activity of this enzyme is low in the liver of the newborn baby (Vest, 1958), in congenital non-haemolytic jaundice (Peterson and Schmid, 1957), and in the Gunn strain of jaundiced rats (Axelrod *et al.*, 1957) and the newborn animal (Brown *et al.*, 1958). Conjugation would appear to be essential for the excretion of bilirubin, though Schmid (1958) points out that a small fraction of bilirubin glucuronide in human bile is possibly hydrolysed to bilirubin either as a result of enzyme activity or on account of the alkalinity of the medium. Recently Isselbacher and McCarthy (1959) have suggested an alternative pathway for the excretion of bilirubin. Bilirubin sulphate has been identified in the bile of rats, cats, and man. The sulphate appears to be linked at the hydroxyl group, in contrast to carboxyl-group linkage of the glucuronide. The sulphate-conjugated fraction contains no glucuronic acid and is alkali-stable. A further unidentified fraction of bile pigment which does not contain glucuronic acid and is alkali-labile would also appear to exist. It is unlikely, however, that these fractions together make up a large part of the total bile pigment, and the majority of the latter is excreted as a glucuronide.

The capacity of the newborn infant's liver to form glucuronides has been studied by Vest (1958). Acetanilide

was administered to full-term and premature babies and the acetyl *p*-aminophenol glucuronide conjugate estimated in the urine. He found that adults and older children are able to excrete 80% of the test substance in 24 hours. Newborn infants, especially if premature, excrete only a small amount in 24 hours, and less than 10% in 48 hours. Normal function is gradually achieved during the first three months of life. It was also observed that there was an inverse relationship between the ability to form glucuronide and the height of the serum bilirubin. The rate at which conjugation occurs in the body has also been studied in infants by the estimation of acetyl *p*-aminophenol glucuronide in the blood. After a dose of acetanilide the blood concentration of the conjugate reaches a peak in four hours in the adult and in 11 hours or more in the newborn infant (Vest and Streiff, 1959). Since the enzyme system of the infant's liver responsible for the conjugation of *p*-aminophenol with glucuronic acid is also thought to be responsible for conjugating bilirubin, it is assumed that the latter is similarly delayed in the newborn infant. The author thus concludes that jaundice in the newborn, in the absence of an actual disease process, is the result of an insufficiency of glucuronic acid transferase activity in the liver.

Bilirubin in the first place is derived from haemoglobin. The latter is broken down by the cells of the reticulo-endothelial system, and the original tetra-pyrrol ring is split up to form a chain of four pyrrol rings known as bilirubin. This can be described diagrammatically as follows:

Bilirubin is then transferred to the liver cells, where two molecules of glucuronic acid are attached. This achieves two things: (1) it renders bilirubin water-soluble, and so it can be readily excreted; and (2) it renders it non-toxic.

There is thus abundant justification for the old division of cases of neonatal jaundice into non-

Conditions Leading to Jaundice in the Newborn

Condition	Type of Blood Pigment	Neurological Complications
Non-obstructive jaundice:		
Functional immaturity of liver	Bilirubin	Kernicterus probable
Haemolytic disease (erythroblastosis)	"	" frequent
Congenital spherocytosis (acholuric jaundice)	"	" rare
Congenital familial non-haemolytic jaundice	"	" possible
Congenital myxoedema (cretinism)	"	" "
Obstructive jaundice:		
Congenital bile-duct atresia	Bilirubin glucuronide	None
Haemolytic disease (persistent jaundice)	" "	"
Tumour	" "	"
Cystic fibrosis of pancreas	" "	"
Galactosaemia	" "	No kernicterus, but mental defect
Infection (bacteria, viruses, protozoa)	Bilirubin and bilirubin glucuronide	Kernicterus rare

obstructive and obstructive types. It must be said, however, that prematurity itself tends to complicate the picture. If an infant has an organic type of disease leading to an obstructive form of jaundice and that infant is also premature, both forms of bilirubin will be found in the blood. The various conditions leading to jaundice in the newborn are shown in the Table.

A. NON-OBSTRUCTIVE JAUNDICE

So far as Great Britain and probably North America are concerned the two common forms of non-obstructive jaundice in the newborn are associated either with functional immaturity of the liver or with haemolytic disease.

1. Functional Immaturity of the Liver

This is the condition formerly known as physiological jaundice, and as a rule the more immature the infant the higher the serum bilirubin and the later the time at which normal liver conjugation of bilirubin to its glucuronide begins. Cunningham (1959) studied the bilirubin in 1,000 babies. Of these, 63 were premature, and in 11 the bilirubin level reached 15 mg. per 100 ml. or over. Among babies weighing more than 2,500 g. at birth the bilirubin level was over 10 mg. per 100 ml. in only 2%. Haemolytic disease had been excluded among the cases in the series. Bowman *et al.* (1957) reported a similar series with a rather higher incidence of bilirubin levels in the upper range: 14% were over 20 mg. and 7% over 25 mg. per 100 ml. As the lesion is thought to be a pure enzyme deficiency, no structural change is seen in the liver in fatal cases. Apart from a few foci of normoblasts, histological examination of the organ reveals nothing (Special Plate, Fig. 1). In a few infants the serum bilirubin rises to a high level and kernicterus develops. About one-quarter of the cases of kernicterus reported by Claireaux *et al.* (1953) were associated with functional immaturity of the liver only.

2. Haemolytic Disease of the Newborn

This results from blood incompatibility between mother and foetus. Usually the rhesus blood group is involved, but other systems such as A-B-O, Kell, Duffy, Lewis, etc., may occasionally be incriminated. In this condition serum bilirubin levels may become extremely high—up to 40 mg. per 100 ml. or more—and in the untreated case kernicterus is apt to develop. In fatal cases the liver is enormously enlarged and the cut surface is reddish-brown in colour. The spleen is also usually much enlarged. On histological examination the hepatic cells are almost obscured by enormous numbers of haemopoietic foci composed of normoblasts and erythroblasts (Special Plate, Fig. 2). Large leucopoietic collections of cells are seen in the portal tracts. The liver cells and Kupffer cells contain haemosiderin, and the Prussian-blue reaction is positive (Claireaux, 1950).

3. Congenital Spherocytosis (Acholuric Jaundice)

This disorder is seldom very severe in the newborn period. It is associated with a congenital defect in the red blood corpuscles which renders them unduly fragile. Haemolysis begins in 0.6% saline and is complete at 0.45%. The spleen is enlarged, and the bone-marrow

shows very active erythropoiesis. As a rule the anaemia is moderate, and jaundice seldom reaches dangerous proportions. Occasionally, however, very high serum bilirubin levels are observed. In the cases reported by Adams (1954) the bilirubin rose to 21 mg. per 100 ml., and a previous sibling had died with kernicterus. Stamey and Diamond (1957) have reported four cases of congenital spherocytosis where severe jaundice was encountered during the first 36 hours of life.

4. Congenital Familial Non-haemolytic Jaundice

This is a rare condition first described by Crigler and Najjar (1952). Seven infants of three consanguineous families were noted to have been severely jaundiced in the neonatal period. These infants were not premature, did not have haemolytic disease, and had no infection. They were jaundiced from birth and had a serum bilirubin level of 23–45 mg. per 100 ml. The majority died within one year, and had neurological abnormalities. One was shown to have kernicterus. Two survivors were reported by Childs and Najjar (1956). These have a permanent serum bilirubin level of 25 mg. per 100 ml. They are of normal height and weight for their age, and the liver and spleen are of normal size. Among the fatal cases necropsy revealed little abnormal. Some bile thrombi were present in the liver, and in one case a mild portal fibrosis was noted. Schmid *et al.* (1957) have investigated the bile in three children with this disorder. According to their report a general defect in glucuronide production would seem to be the basic abnormality. A similar group of patients has been studied by Childs *et al.* (1959). All patients showed a marked impairment in glucuronide conjugation. A conjugation defect midway between that of the patients and normal controls was observed in the majority of the relatives. The authors believe that the gene responsible for the disease is recessive in respect to the jaundice, but shows incomplete dominance so far as glucuronide conjugation is concerned.

5. Congenital Myxoedema (Cretinism)

Prolonged jaundice in three cases of congenital myxoedema was reported by Åkerrén (1954). The patients' general condition improved after treatment with thyroid extract and the jaundice vanished. Kernicterus has not been reported with certainty in this condition, but Christensen (1956) has described a case in which the serum bilirubin reached 21 mg. per 100 ml., and defective vision, hearing, and intellect were suspected. It has been suggested that lack of thyroid hormone has an inhibitory effect on the enzyme systems responsible for the conjugation of bilirubin (Lathe *et al.*, 1958).

B. OBSTRUCTIVE JAUNDICE

This group is less clearly defined than the non-obstructive group, and is considerably less common. In non-obstructive jaundice the serum bilirubin level is raised as a result of haemolysis or deficient liver-cell function. Neurological damage is not infrequent. In the group with obstructive jaundice there are such obvious conditions as biliary atresia and obstruction of the common bile-duct, but also a number of rather rare conditions which cause severe jaundice and in which the main pigment involved is bilirubin glucuronide. In a

very few instances, however, both types of bile pigment may be present, and if bilirubin should predominate kernicterus can occur. It is, however, extremely rare.

1. Congenital Atresia of the Bile-ducts

This occurs in about 1 in 20,000 deliveries (Moore, 1953). It is the commonest form of neonatal obstructive jaundice. The sex incidence is equal. Moore (1953) found other associated congenital abnormalities in five of his 31 cases, and four of these involved the urinary tract. There are two forms of the condition: (a) an extrahepatic atresia, and (b) an intrahepatic atresia.

(a) Extrahepatic Atresia

This is more common than the intrahepatic variety. There are many possibilities of maldevelopment of the biliary tree, from a stenosis of the common duct to absence of all extrahepatic biliary structures. The liver is enlarged and a biliary cirrhosis soon develops. On slicing, the cut surface of the organ is seen to be green. Histological examination shows a marked peribiliary fibrosis with proliferation of small bile-ducts. The intercellular canaliculi and small bile-ducts contain bile plugs. Some inflammatory-cell infiltration is present in the connective tissue; these cells are mainly lymphocytes and plasma cells. (Special Plate, Fig. 3).

(b) Intrahepatic Atresia

This is exceedingly rare. Curiously enough, it is associated with a longer period of survival than the extrahepatic form (Ahrens *et al.*, 1951; Haas and Dobbs, 1958). The liver is enlarged and green in colour, and it has a finely granular cut surface. The gall-bladder and common duct may be normal. Cirrhosis is a less-marked feature on histological examination. There is some increase in the fibrous tissue of the portal tracts, but no bile-ducts are seen. Bile plugs are present in the intercellular canaliculi.

The skin shows multiple xanthomatous lesions in this form of the disease (Ahrens *et al.*, 1951). In both types of atresia the predominant serum pigment is bilirubin glucuronide.

2. Haemolytic Disease

In a very few cases infants with haemolytic disease remain jaundiced for several weeks. The main pigment in the blood is now found to be bilirubin glucuronide, and apart from the jaundice the infants appear to be well. This is the so-called inspissated-bile syndrome. The liver is enlarged and very firm. The intercellular bile-channels are distended with dark inspissated bile. The portal tracts are normal, as are the main bile-ducts. As a rule the condition clears without treatment, and cirrhosis is almost unknown.

3. Tumour

Very rarely obstructive jaundice may result from the pressure of tumour masses on the common bile-duct. Gross (1953) reported two such cases. In one patient there was a haemangio-endothelioma of the head of the pancreas, and the other had Niemann-Pick disease with enlarged nodes in the gastro-hepatic ligament. Neoplasms of the liver itself are rare in the newborn. The

commonest are haemangiomas. These may be exceedingly large, and lead to a fatal haemorrhage if ruptured. The infants may or may not be jaundiced. Very rarely a malignant hepatoma may be found in the neonatal period. These cause great hepatomegaly and metastasize freely. They are associated with jaundice and ascites.

4. Cystic Fibrosis of the Pancreas

Sass-Kortsak *et al.* (1957) have reported a few cases of obstructive jaundice in babies with cystic fibrosis of the pancreas. Focal biliary cirrhosis in a stillborn infant has been described by Claireaux (1956). The lesion tends to be patchy, and in the early stages is mainly subcapsular in situation. In older children this may be followed by widespread hepatic cirrhosis and severe jaundice.

5. Galactosaemia

In severe cases of galactosaemia the infant may be jaundiced shortly after birth. Galactosaemia is a metabolic disorder resulting from a lack of the enzyme capable of converting galactose-1-phosphate to glucose-1-phosphate (Schwarz *et al.*, 1955). The damage to liver and other tissue cells is the result of the toxic action of some metabolite of galactose. In this disease the liver is enlarged and the cut surface has a pale-yellow colour. The liver cells on histological examination show severe fatty change with pronounced vacuolation (Special Plate, Fig. 4). Periportal fibrosis occurs and cirrhosis of the liver develops. The bile canaliculi contain plugs of inspissated bile. Occasional multinuclear giant cells are found. In a few cases liver-cell adenomas develop, and the cells taking part in this formation are full of glycogen (Edmonds *et al.*, 1952). In the untreated patient cataracts may develop after the third week of life.

6. Jaundice in Infection

(a) Bacterial Infection

Jaundice complicating neonatal sepsis is now a rare event. It is only infrequently very pronounced. The liver is enlarged and signs of generalized infection are present. Bilirubin and bilirubin glucuronide levels in the blood are both raised, but the jaundice rapidly disappears if the infection is controlled by the appropriate antibiotics. In a few instances the jaundice results from direct liver damage caused by abscess formation following umbilical sepsis. A rare cause of sepsis giving rise to miliary necrosis of the liver is infection with *Listeria monocytogenes* (Claireaux, Gerrard, and Marsland, 1955).

Severe congenital syphilis is also a rarity in Great Britain, and Nabarro (1954) has expressed some doubt whether it is ever a cause of jaundice in the newborn period. Cases of gross liver enlargement with jaundice previously ascribed to congenital syphilis are probably examples of some other condition. Syphilis can cause a severe intralobular cirrhosis of the liver in the infant, and the histological appearances are highly characteristic (Special Plate, Fig. 5). This may cause jaundice in the older infant.

(b) Viral Infection

(i) *Herpetic Hepatitis*.—A viraemia resulting from infection with herpes simplex virus causes widespread

visceral lesions, including a characteristic focal necrosis of the liver. Jaundice is usually present, but is not necessarily severe. The infant is frequently premature. The liver is enlarged, and multiple white nodules, some of which show umbilication, are raised above the surface of the organ. Similar nodules are seen on the cut surface when the liver is sliced. On histological examination the liver cells in the nodular zones are seen to be necrotic; those bordering the necrotic areas show nuclear swelling, and eosinophilic inclusion bodies may be present. Similar necrotic areas may be found elsewhere in the body, notably in the brain and suprarenal glands.

(ii) *Congenital Hepatitis*.—The viral nature of this condition is still in some doubt, but the probabilities are in favour of a virus being the aetiological agent. Stokes *et al.* (1951) reported that serum from an infant with hepatitis and from his mother caused hepatitis when injected into volunteers. The incubation period was such that either virus A (infective hepatitis) or virus B (homologous serum jaundice) could be responsible. It would seem unlikely that virus A is the infective agent, as viraemia in this condition is transient and the mother soon develops antibodies to the virus. These strictures do not apply in the case of virus B (Gellis *et al.*, 1954). Smetana and Johnson (1955) believe that the histological appearance of the liver in neonatal hepatitis suggests that the primary lesion is a congenital maldevelopment involving the intercellular bile canaliculi leading from the liver cells to the bile-channels. The "hepatitis" in such an event merely represents the liver cells' response to a variety of stimuli (Bain *et al.*, 1957). Miller (1957) has reported the condition in one of binocular twins. Liver-function tests in the mother would appear to indicate some degree of liver damage, which the author associates with hepatitis. According to Weller *et al.* (1957) the virus of cytomegalic inclusion disease has been recovered from a patient presenting with the signs of a neonatal hepatitis. Obstructive jaundice in 25 patients was studied by Sass-Kortsak *et al.* (1957). Biliary atresia, haemolytic disease, sepsis, and galactosaemia were excluded from their series. They found two examples of fibrocystic disease of the pancreas, and the remaining 23 cases were classified as idiopathic. Some of the idiopathic cases had negative liver-function tests, and the others gave positive results. It was the latter group which were thought to be affected by some virus. Hsia *et al.* (1958) made a new approach to the problem. They conceded that no certain aetiological agent had been demonstrated in neonatal hepatitis; they therefore investigated the genetic background of 59 families comprising 11 affected and 50 normal siblings. Their statistical data would suggest the theory that the trait responsible for neonatal "hepatitis" is an autosomal recessive. Shorter *et al.* (1959) were also unable to incriminate a single aetiological agent, and suggest that the histological response found in the infant's liver may be to a number of quite different stimuli.

At necropsy the liver may be enlarged, or less frequently it is smaller than usual. The organ is green and the cut surface has a granular appearance. The gall-bladder and main ducts are normal. Histologically, the normal lobular architecture is lost. Numerous exceedingly large multinucleated giant cells are seen (Special Plate, Fig. 6). These cells may have 30 or 40 nuclei. Their cytoplasm has a brownish tinge owing to the deposition of haemosiderin, and the Prussian-blue

reaction is positive. Beyond the necrotic areas some islands of regenerating liver cells are found, and these islands may contain haemopoietic foci. There is an increase in fibrous tissue round the necrotic liver cells and in the portal tracts. This connective tissue shows infiltration with lymphocytes and plasma cells. Some proliferation of small bile-channels occurs in the portal tracts, and many contain bile plugs. The origin of the multinuclear giant cells is obscure. Craig and Landing (1952) regarded them as degenerated liver cells fusing to form a syncytial mass. Stokes *et al.* (1951) suggested that they were a response of the liver cell to virus infection, and Dible *et al.* (1954) believed they represented rapid proliferation of liver cells during recovery from the virus infection. If the infant survives beyond the neonatal period a true cirrhosis develops. The infant is jaundiced, and both direct and indirect pigment levels are raised. Kernicterus has been reported in one case by Erhlich and Ratner (1955).

(iii) *Cytomegalic Inclusion Disease*.—Generalized inclusion-body disease is occasionally found in the newborn. The infective agent is said to be salivary-gland virus (Smith, 1956; Weller *et al.*, 1957). At necropsy the liver and spleen are found to be enlarged. On histological examination typical inclusion bodies are seen in the epithelial cells of salivary-gland acini, renal tubules, bile-ducts, pancreatic and thyroid acini, and lying free in the lung alveoli. Extramedullary haemopoiesis is marked in the liver if the infant dies shortly after birth. In the older infant these foci disappear. There is some increase in the connective tissue in the portal tracts accompanied by a small bile-duct proliferation. Bile thrombi are present in the canaliculi. The affected bile-duct epithelial cells are greatly enlarged, and contain both intranuclear and cytoplasmic inclusions. The intranuclear inclusion bodies are large acidophilic masses; cytoplasmic inclusions are small, multiple, and basophilic. They are usually less numerous in the liver than in the kidneys. The infants are jaundiced, and this may become severe. There is an increase in both bilirubin and bilirubin glucuronide in the blood. Kernicterus has occasionally been reported (Kinney, 1942; France, 1951).

(c) Protozoal Infection

An acute generalized infection caused by *Toxoplasma gondii* may involve the liver. The infants are jaundiced, and the liver and spleen are enlarged. Calcification of the brain and chorioretinitis are prominent features. On histological examination the liver shows marked erythropoietic activity together with leucocytic infiltration. Histiocytes containing toxoplasma parasites may be present. Bile thrombi are present in the intercellular canaliculi, and the liver cells contain haemosiderin. Areas of necrosis may be present in liver, heart, lungs, and spleen. In these necrotic areas toxoplasma may be found.

KERNICTERUS

The first report of yellow staining of the brain in jaundiced infants is ascribed to Orth (1875). Schmörl (1904) coined the term "kernicterus," and the association with erythroblastosis was soon established. Aiden *et al.* (1950) then reported the occurrence of kernicterus in jaundiced premature infants who did not suffer from haemolytic disease. Similar cases were described by

A. E. CLAIREAUX: NEONATAL HYPERBILIRUBINAEMIA

FIG. 1.—Liver. Functional immaturity. No structural defect present, and only an occasional focus of haemopoiesis. (Haematoxylin and eosin. $\times 60$.)

FIG. 2.—Liver. Haemolytic disease. Innumerable haemopoietic foci present. (Haematoxylin and eosin. $\times 60$.)

FIG. 3.—Liver. Congenital obliteration of bile-ducts. Increased fibrous tissue in portal tracts. Proliferation of small bile-ducts. Bile-plugs in canaliculi. (Haematoxylin and eosin. $\times 75$.)

FIG. 4.—Liver. Galactosaemia. (Haematoxylin and eosin. $\times 90$.)

FIG. 5.—Liver. Congenital syphilis. Pericellular intralobular cirrhosis. (Haematoxylin and eosin. $\times 90$.)

FIG. 6.—Liver. Neonatal hepatitis. Large number of multinucleated giant cells. (Haematoxylin and eosin. $\times 90$.)

Zuelzer and Mudgett (1950). Less frequent neurological damage may occur as a sequel to congenital spherocytosis, congenital non-haemolytic jaundice, cytomegalic inclusion disease, or neonatal hepatitis. The association between haemolytic disease and kernicterus was well established, but the risk to the premature infant suffering from "physiological jaundice" was less well known. Meyer (1956) studied the serum bilirubin levels in premature infants with this in mind. He found that in the infant under 2,000 g. birth weight the bilirubin was still rising on the sixth day of life, whereas in the mature infant the peak value was past and the level was falling. He noted that infants with kernicterus had a serum bilirubin level over 18 mg. per 100 ml. A larger group of premature infants was studied by Crosse *et al.* (1958). Of 1,320 premature infants, no fewer than 92 (8.4%) required an exchange transfusion on account of dangerously high levels of bilirubin in the blood. Five of the surviving infants have evidence of neurological damage. These workers found that no infant with a bilirubin level below 20 mg. per 100 ml. developed kernicterus. At levels of 22–27 mg. per 100 ml. the incidence of kernicterus was 19%, and at a level over 30 mg. per 100 ml. it was 33½%. Waters (1958) also stressed the danger of allowing the serum bilirubin level to rise above 20 mg. per 100 ml.

It would seem probable, however, that the incidence of very high levels of bilirubin is relatively rare in the premature infant in the absence of a disease process; it probably is not in excess of 1–2% of premature babies under normal conditions. However, it was observed that under certain circumstances the incidence of high serum bilirubin levels would be increased (Laurance, 1955). If excessive doses of vitamin-K analogue ("synkavit") are given, 20–30 mg. daily, then high bilirubin levels and severe jaundice will follow. Asteriadou-Samartzis and Leikin (1958) administered vitamin K to 119 premature infants. Serum bilirubin levels were higher in the infants given vitamin K than in control infants. Vitamin K, on the other hand, had little or no effect, nor, it must be emphasized, did vitamin K when given in correct therapeutic dosage. More recently Lucey and Dolan (1959) have reported hyperbilirubinaemia in newborn infants whose respective mothers received vitamin K prior to delivery. In seven cases there was a marked rise in the bilirubin in the first three days of life (20–47 mg. per 100 ml.), and one infant died with kernicterus and one survivor was permanently affected. The authors believe that vitamin K crosses the placental barrier and damages the infant's liver. Gardner (1959) is also of the opinion that vitamin K crosses the placental barrier.

The connexion between very high serum levels of bilirubin and kernicterus, even in the absence of haemolytic disease, was quite obvious, but Silverman *et al.* (1956) reported a number of cases of kernicterus where the serum bilirubin level was only around 15 mg. per 100 ml. They found that their patients had been on a regimen of penicillin and "gantrisin" (sulphafurazole). A further study of the bilirubin levels in infants treated with gantrisin was carried out by Harris *et al.* (1958), and they showed that kernicterus could occur when the bilirubin level was under 15 mg. per 100 ml. A possible explanation for this phenomenon is offered by Odell (1959). It is known that bilirubin is carried on the plasma albumin. If the plasma albumin concentration is low, as in the premature infant, the former becomes readily

saturated with bilirubin. If another organic anion is now introduced into the blood which will compete with bilirubin for attachment to albumin, then the amount of dissociated bilirubin in the blood will rise. This diffusible unbound bilirubin can attack the cells of the central nervous system. Gantrisin is known to displace albumin-bound bilirubin *in vitro* (Odell, 1958), and apparently will do so *in vivo*. Salicylates are apparently even more effective than the sulphonamides in this respect (Odell, 1959b).

In all the above examples the infants affected by kernicterus have been premature. According to Boon (1957) kernicterus is not uncommon in mature babies in Singapore even if haemolytic disease is excluded. This author ascribes the condition to poor liver function in the infant associated with maternal malnutrition. It may be, however, that some specific factors are involved which are not as yet known.

For many years it was believed that the brain was first damaged by anoxia or other agent, and bilirubin then stained the devitalized tissues. When the nature of the yellow pigment in the brain was demonstrated it was suggested that bilirubin itself was the toxic agent (Claireaux *et al.*, 1953). It is now generally accepted that bilirubin is a highly toxic substance, and one of the effects of its conjugation with glucuronic acid is to render it non-toxic. Ernster *et al.* (1957) have shown that bilirubin inhibits the enzymatic mechanism in brain cells by which phosphorylation is coupled to respiration. They consider that kernicterus is the result of metabolic damage to the central nervous system caused by bilirubin.

The brain is often stained a diffuse yellow colour, especially in the premature infant, but certain nuclear structures are found to be deeply discoloured. The pigmented area may be light yellow or deep orange in colour. The structures most liable to be affected are the corpus striatum (globus pallidus, putamen, lenticular nucleus), thalamus and subthalamic nuclei, hippocampus, nucleus of third cranial nerve, mammillary bodies, red nucleus, nuclei in floor of fourth ventricle, dentate nucleus, flocculus and vermis of cerebellum, and the olives. In some cases the grey matter of the spinal cord and cerebral cortex is also affected. The distribution of lesions varies from case to case, and is generally similar in jaundice caused by functional hepatic immaturity and in haemolytic disease. In the former, however, the yellow staining tends to be slightly paler, and diffuse staining in the region of the periventricular white matter is relatively common. The yellow colour gradually fades if the infant should survive, and in older infants only some slight shrinkage of affected nuclei may be noted at necropsy. On histological examination the brain from an infant dying in the first week of life may show little change. There may be some degree of chromatolysis in the neurones from stained areas, and some of the cells may contain yellow pigment. In older infants there is evidence of marked loss of neurones and replacement gliosis has occurred. Crome (1955) has reported similar changes in the cerebral cortex in patients with mental deficiency following neonatal jaundice.

Neurological damage in infants with severe jaundice in the neonatal period may result in mental deficiency, spasticity, athetosis, or nerve deafness. Modern methods of treatment are directed towards the prevention of such complications.

REFERENCES

- Adams, W. C. (1954). *J. Pediat.*, **44**, 213.
- Ahrens, E. H., Harris, R. C., and MacMahon, H. E. (1951). *Pediatrics*, **8**, 628.
- Aidin, R., Corner, B., and Tovey, G. (1950). *Lancet*, **1**, 1153.
- Åkerrén, Y. (1954). *Acta paediat. (Uppsala)*, **43**, 411.
- Asteriadou-Samartzis, E., and Leikin, S. (1958). *Pediatrics*, **21**, 397.
- Axelrod, J., Schmid, R., and Hammaker, L. (1957). *Nature (Lond.)*, **180**, 1426.
- Bain, G. O., Wang, G. C., and Misanik, L. F. (1957). *J. Pediat.*, **51**, 549.
- Billing, B. H., Cole, P. G., and Lathe, G. H. (1957). *Biochem. J.*, **65**, 774.
- and Lathe, G. H. (1958). *Amer. J. Med.*, **24**, 111.
- Boon, W. H. (1957). *Arch. Dis. Childh.*, **32**, 85.
- Bowman, J. M., Rapmund, G., and Harris, R. C. (1957). *Amer. J. Dis. Childh.*, **93**, 75.
- Brown, A. K., Zuelzer, W. W., and Burnett, H. H. (1958). *J. clin. Invest.*, **37**, 332.
- Childs, B., and Najjar, V. A. (1956). *Pediatrics*, **18**, 369.
- Sidbury, J. B., and Migeon, C. J. (1959). *Ibid.*, **23**, 903.
- Christensen, J. F. (1956). *Acta paediat. (Uppsala)*, **45**, 367.
- Claireaux, A. E. (1950). *Arch. Dis. Childh.*, **25**, 61.
- (1956). *Ibid.*, **31**, 22.
- Cole, P. G., and Lathe, G. H. (1953). *Lancet*, **2**, 1226.
- Gerrard, J. W., and Marsland, E. A. (1955). *Rev. int. Hépatol.*, **5**, 1153.
- Cole, P. G., and Lathe, G. H. (1953). *J. clin. Path.*, **6**, 99.
- Craig, J. M., and Landing, B. H. (1952). *A.M.A. Arch. Path.*, **54**, 321.
- Crigler, J. F. jun., and Najjar, V. A. (1952). *Pediatrics*, **10**, 169.
- Crome, L. (1955). *J. Neurol. Neurosurg. Psychiat.*, **18**, 17.
- Crosse, V. M., Wallis, P. G., and Walsh, A. M. (1958). *Arch. Dis. Childh.*, **33**, 403.
- Cunningham, A. A. (1959). *Ibid.*, **34**, 262.
- Dible, J. H., Hunt, W. E., Pugh, V. W., Steingold, L., and Wood, J. H. F. (1954). *J. Path. Bact.*, **67**, 195.
- Edmonds, A. M., Hennigar, G. R., and Crooks, R. (1952). *Pediatrics*, **10**, 40.
- Ehrlich, J. C., and Ratner, I. M. (1955). *Amer. J. Path.*, **31**, 1013.
- Ernster, L., Herlin, L., and Zetterström, R. (1957). *Pediatrics*, **20**, 647.
- France, N. E. (1951). *Arch. Dis. Childh.*, **26**, 588.
- Gardner, L. I. (1959). *Pediatrics*, **23**, 1017.
- Gellis, S. S., Craig, J. M., and Hsia, D. Y. Y. (1954). *Amer. J. Dis. Childh.*, **88**, 285.
- Gross, R. E. (1953). *The Surgery of Infancy and Childhood*. Saunders, Philadelphia.
- Haas, L., and Dobbs, R. H. (1958). *Arch. Dis. Childh.*, **33**, 396.
- Harris, R. C., Lucey, J. F., and MacLean, J. R. (1958). *Pediatrics*, **21**, 875.
- Hsia, D. Y. Y., Boggs, J. D., and Driscoll, S. G. (1958). *Amer. J. Dis. Childh.*, **95**, 485.
- Isselbacher, K. J., and McCarthy, E. A. (1959). *J. clin. Invest.*, **38**, 645.
- Kinney, T. D. (1942). *Amer. J. Path.*, **18**, 799.
- Lathe, G. H., Claireaux, A. E., and Norman, A. P. (1958). In *Recent Advances in Paediatrics*, 2nd ed., ed. D. Gairdner, p. 87. Churchill, London.
- and Walker, M. (1957). *Biochem. J.*, **67**, 9 P.
- Laurance, B. (1955). *Lancet*, **1**, 819.
- Lucey, J. F., and Dolan, R. G. (1959). *Pediatrics*, **23**, 553.
- Meyer, T. C. (1956). *Arch. Dis. Childh.*, **31**, 75.
- Miller, B. (1957). *Amer. J. Dis. Childh.*, **94**, 318.
- Moore, T. C. (1953). *Surg. Gynec. Obstet.*, **96**, 215.
- Nabarro, D. N. (1954). *Congenital Syphilis*. Arnold, London.
- Odell, G. B. (1958). *Amer. J. Dis. Childh.*, **96**, 535.
- (1959a). *J. Pediat.*, **55**, 268.
- (1959b). *J. clin. Invest.*, **38**, 823.
- Orth, J. (1875). *Virchows Arch. path. Anat.*, **63**, 447.
- Peterson, R. E., and Schmid, R. (1957). *J. clin. Endocr.*, **17**, 1485.
- Sass-Kortsak, A., Bowden, D. H., Sirek, O. V., and Cherniak, M. (1957). *Amer. J. Dis. Childh.*, **94**, 560.
- Schmid, R. (1956). *Science*, **124**, 76.
- (1958). *A.M.A. Arch. intern. Med.*, **101**, 669.
- Hammaker, L., and Axelrod, J. (1957). *Arch. Biochem.*, **70**, 285.
- Schmörl, G. (1904). *Verh. dtsch. path. Ges.*, 1903, **6**, 109.
- Schwarz, V., Golberg, L., Komrower, G. M., and Holzel, A. (1955). *Biochem. J.*, **59**, xxii.
- Shorter, R. G., Baggenstoss, A. H., and Logan, G. B. (1959). *Amer. J. Dis. Childh.*, **98**, 359.
- Silverman, W. A., Andersen, D. H., Blanc, W. A., and Crozier, D. N. (1956). *Pediatrics*, **18**, 614.
- Smetana, H. F., and Johnson, F. B. (1955). *Amer. J. Path.*, **31**, 747.
- Smith, M. G. (1956). *Proc. Soc. exp. Biol. (N.Y.)*, **92**, 424.
- Stamey, C. C., and Diamond, L. K. (1957). *Amer. J. Dis. Childh.*, **94**, 616.
- Stokes, J., Wolman, I. J., Blanchard, M. C., and Farquhar, J. D. (1951). *Ibid.*, **82**, 213.
- Talafant, E. (1957). *Coll. Trav. chim. Tchecosl.*, **22**, 661.
- van den Bergh, A. A. H., and Muller, P. (1916). *Biochem. Z.*, **77**, 90.
- and Snapper, I. (1913). *Dtsch. Arch. klin. Med.*, **110**, 540.
- Vest, M. (1958). *Arch. Dis. Childh.*, **33**, 473.
- and Streiff, R. R. (1959). *Amer. J. Dis. Childh.*, **98**, 688.
- Waters, W. J. (1958). *J. Pediat.*, **52**, 559.
- Weller, T. H., Macauley, J. C., Craig, J. M., and Wirth, P. (1957). *Proc. Soc. exp. Biol. (N.Y.)*, **94**, 4.
- Zuelzer, W. W., and Mudgett, R. T. (1950). *Pediatrics*, **6**, 452.

INTRACTABLE UTERINE HAEMORRHAGE DUE TO ABNORMAL SUBENDOMETRIAL BLOOD-VESSELS AFTER CHILDBIRTH AND MISCARRIAGE

BY

R. SALM, M.D., L.R.C.P.&S.Ed.

Consultant Pathologist

AND

P. N. SIMONS, F.R.C.S.Ed., M.R.C.O.G.

Consultant Gynaecologist

Camborne-Redruth Hospital, Redruth, Cornwall

[WITH SPECIAL PLATE]

Two cases of severe and intractable menorrhagia in young women are presented which, clinically, were considered to be functional in origin, as no organic cause was clinically demonstrable. Both patients were finally subjected to the drastic procedure of hysterectomy, as every conservative measure taken had failed to stop the recurrent severe uterine haemorrhages. Detailed examination of the operation specimens revealed the presence of large, thick-walled subendometrial blood-vessels, which were interpreted as being normal radial myometrial vessels, probably arteries, which had originally supplied the placental site, had increased in calibre during pregnancy, but had failed to involute during the puerperium. Erosion of these vessels during menstruation we regard as the cause of the exceptionally severe and recurrent menorrhagia—a cause which, to our knowledge, has not been recorded previously.

Case 1

A woman aged 25 was admitted as an emergency on February 15, 1958, because of sudden, severe, painless vaginal haemorrhage on the second day of a period which had begun normally. She had two children, the first being born in 1954. A miscarriage, concerning which no details were available, occurred in 1956, and in August, 1957, a normal delivery at home was followed by an uneventful puerperium. A few weeks afterwards she had a heavy period with the passage of clots. A regular 28-day cycle was re-established, but menorrhagia continued and was at times severe.

On examination she showed the clinical signs of acute blood loss. Morphine, 1/4 gr. (6 mg.), was given intramuscularly and 1,600 ml. of whole blood was transfused. When her general condition had improved sufficiently a quantity of blood-clot was removed from the vagina. No abnormality could be found in the cervix, uterine body, and adnexae, nor was there any evidence of systemic disease or blood dyscrasia. A full blood-count gave the following results: haemoglobin, 7.4 g. per 100 ml.; red cells, 2,800,000 per c.mm.; total white cells, 7,700 per c.mm., with polymorphs predominating; platelets, 433,000 per c.mm. Bleeding and clotting times were normal.