of this moderately rare condition, but was reported, I believe, in the recent review by Brain and Henson.* In this case, in a context of general weakness, there seems to have been, in addition, gross focal weakness and atrophy affecting the legs and buttocks. There was a history of sciatic pain followed by foot-drop, and both ankle jerks were absent. In the presence of multiple metastases in the lumbar vertebrae, involvement of the roots of the cauda equina seems a much more likely explanation.

Professor MCMICHAEL: I suppose we could account for the muscular weakness on the extreme disturbance of the electrolytes.

Professor RUSSELL FRASER: Yes. I think the idea that it depended only on serum potassium has been

*Brain, R., and Henson, R. A., *Lancet*, 1958, 2, 971.

eliminated, but hypercalcaemia itself is a great cause of neuromuscular weakness and I think perhaps that may have been the major factor.

Professor C. V. HARRISON: I think we could decide whether there was parathyroid hyperactivity—I don't mean at this moment, but I think we could tell by examining the sections of this patient's cancellous bone in sites clearly free of tumour. You would get osteoclastic resorption with marrow fibrosis in an area where the presence of tumour could not explain it. I don't suppose that has been done, because we were looking for tumour, but I think we could do it now.

We are grateful to Dr. J. P. Shillingford and Dr. B. E. Heard for assistance in preparing this report, and to the photographic department of the Postgraduate Medical School for the illustrations.

Drug Treatment of Disease

BACILLARY DYSENTERY*

BY

A. R. D. ADAMS, M.D., F.R.C.P.

Physician in Charge, Clinical Department, Liverpool School of Tropical Medicine

The bacillary dysenteries (shigelloses) are of world-wide distribution, but are particularly prevalent in warm countries where sanitation is defective. In Great Britain they are apt to occur epidemically in institutions, especially those for children or for mental patients. The organisms responsible are species of the genus *Shigella*; they include *Sh. shigae*, *Sh. schmitzii*, *Sh. flexneri*, *Sh. boydii*, and *Sh. sonnei*. There are several types and variants of *Sh. flexneri* and also *Sh. boydii*. The first to be identified, *Sh. shigae*, is the most virulent of these organisms and is more particularly found in the tropics; *Sh. sonnei*, the least virulent and one which rarely causes death, is that usually found in Britain. To an increasing extent *Sh. sonnei* is becoming the predominant species throughout Europe and the Middle East.

Sulphonamides

During the first world war, when there was no specific treatment for them, the bacillary dysenteries caused great morbidity and a considerable mortality in the forces, notably among those engaged in the Eastern Mediterranean theatres of war. At the beginning of the second world war sulphaguanidine, a sulphonamide which in passing through the bowel is but little absorbed, was shown to arrest clinical attacks of Flexner dysentery in children in a dramatic manner. It proved of equal value in adults serving in the forces in the Eastern Mediterranean; therefore it was produced on a large scale for the treatment of bacillary dysentery among the British forces wherever the disease was prevalent. As a result bacillary dysentery ceased to be a continuous major problem in the maintenance of operational activity by the troops.

While the drug initially was still in short supply some patients suffering from bacillary dysentery were treated with the freely absorbable sulphonamide sulphapyridine.

*The article by Dr. Adams on "Amoebiasis and Amoebic Dysentery" in this series appeared in the *British Medical Journal* of March 26 (p. 956).

This proved effective in controlling the dysenteric attacks but was prone to cause crystalluria, leading to anuria which, if not relieved, proved fatal. This hazard is particularly great in a patient suffering from a febrile and dehydrating disease, associated as this is with vomiting, diarrhoea, and sweating, in a hot climate. Consequently sulphaguanidine, to the exclusion of other sulphonamides, was adopted by the British forces for the treatment of bacillary dysentery. The American forces, when later engaged, preferred the freely absorbable sulphadiazine as their choice; when using it certain recommendations were laid down for the prevention of crystalluria, including dosage with alkalis to render the urine alkaline. The latter may not be possible even when the effort to achieve it is pushed to the stage of alkalosis.

Drug Resistance

After the sulphonamides had been in use for a few years, drug-resistance became evident during an outbreak of Flexner bacillary dysentery in children in an institution; sulphonamide treatment in the earlier cases proved effective, while in the later cases it failed to control the infection. The acquirement of drug-fastness by strains of *Shigella* is now a recognized risk attending their treatment with the sulphonamides.

Antibiotics

These sulphonamide-resistant infections were treated with streptomycin, given orally; it proved therapeutically active, though less so than had the sulphonamides in the earlier cases. Streptomycin when given orally is poorly absorbed; it therefore does not produce the toxic side-effects with which its parenteral administration is associated. It is not inactivated materially in the intestine. Other antibiotics, especially chloramphenicol and the tetracyclines, have since usually proved to be more efficient than is streptomycin for the treatment of shigellosis. Drug-resistance to the sulphonamides is

especially likely to be acquired by strains of *Sh. flexneri*; *Sh. sonnei* infections often do not respond to sulphonamide treatment, or, if they do so, very rapidly become resistant to it. Therefore in these infections, and especially the latter, antibiotic therapy is the treatment of choice on all occasions whether or not sulphonamides also are given. So far there has been no evidence of the development by strains of *Shigella* of drug-fastness to the wide-spectrum antibiotics.

DRUG THERAPY IN PRACTICE

Sulphonamides

Where cost and availability are of significance, as in the tropics, sulphaguanidine and the later drugs of the same type (succinylsulphathiazole, phthalylsulphathiazole) still are widely and effectively used for the treatment of the forms of bacillary dysentery amenable to them. The dosage advocated is about 0.1 g. per kg. of body weight by mouth initially; half this dose is given 4-hourly thereafter until at least 24 hours after the diarrhoea ceases. This same dose is then continued at 8-hourly intervals for at least another three days. Ten days is the maximum period over which these drugs should be given. In practice, if they are effective the result of treatment should be clearly evident within 48 hours of its start. In round figures the dosage for an adult with any of these three drugs is initially 6 g. orally, followed by half this amount at the appropriate intervals.

Other sulphonamides, particularly sulphadiazine, are equally effective; but their use is governed by the considerations to which reference has been made. In tropical practice, especially up-country, most British workers adhere to the view that they are better avoided if the less readily absorbable compounds are available.

Antibiotics

Where cost and availability are not of prime importance the broad-spectrum antibiotics, in the form of chloramphenicol or the tetracyclines, by mouth are the first choice. The tetracyclines are not so rapidly absorbed from the intestine nor so freely diffused as is chloramphenicol. The larger the oral dose the less, proportionately, are the tetracyclines absorbed from the bowel and the greater, therefore, is the concentration of these drugs in the stools. In bacillary dysentery, unlike typhoid, the causative organisms are localized in the gut; so this may be an advantage in the treatment of this condition if the causation is beyond doubt and is not complicated by the incidence—for example, during an epidemic in children—of cases of gastro-enteritis due to infection with *Salmonella* or coliform organisms.

Chloramphenicol generally is claimed to be more effective than are the tetracyclines. It can be given orally to infants or children at a daily dosage of 50 to 100 mg. per kg. body weight, in divided doses, for five days. Adults can be given 3 to 4 g. daily, by mouth in four divided doses, for the same period. It is effective against most strains of all species of *Shigella*, and these develop no acquired resistance to it. Adults suffering from sulphonamide-resistant Shiga or Flexner dysentery have responded rapidly to an initial dosage of 2 g. followed by 1 g. in 12 hours and again in 24 hours. Toxic side-effects in the form of stomatitis and glossitis, diarrhoea, and pruritus are less than those associated with tetracycline treatment. Chloramphenicol may

depress the bone marrow. Allergic skin manifestations, which may occur in a few sensitive patients, clear on stopping the drug.

A dosage scale of oxytetracycline ("terramycin") found satisfactory in one considerable epidemic was as follows: for children of 6 months or less, 25 to 50 mg.; up to 1 year, 100 mg.; from 2 to 5 years, 150 to 200 mg.; older, 250 mg.; and adults, 250–500 mg. These amounts were given daily by mouth, in divided dosage, for five days. Bigger daily doses than these have been given over longer periods without ill effect.

Sulphonamide-resistant strains of Shiga and of Flexner dysentery in adults responded readily to dosage initially of 2 g. and then 1 g. at 12 hours and 24 hours. Chlorotetracycline ("aureomycin") produced a similar result in this group of patients.

The toxic side-effects of tetracycline treatment, though very rarely seen, are allergic and neurotoxic, nausea, and blood dyscrasia. In addition, treatment with this drug by depressing the normal bacterial flora permits overgrowth of *Candida albicans*; this may lead to stomatitis, glossitis and a hairy tongue, and pruritus ani and vulvae. Very rarely after prolonged dosage this overgrowth can assume alarming proportions, with grave infection of the alimentary, respiratory, and genital tracts. The development of resistance to tetracycline has not been found in strains of *Shigella*.

Streptomycin, if prescribed, is given orally in doses of 0.5 g. twice daily regardless of age. It is not absorbed to a material extent from the bowel; therefore there is very little risk of the development of its classical toxic side-effects; here also there is no evidence of acquired drug-resistance.

A combination of antibiotics and of sulphonamides, such as streptomycin and sulphaguanidine ("guaninycin"), is very effective in treating gastro-enteritis of infants. The dosage is 20 mg. per lb. (45 mg. per kg.) of body weight daily. "Streptotriad" (each tablet containing streptomycin 65 mg., sulphathiazole 100 mg., sulphadiazine 100 mg., and sulphamerazine 65 mg.), in a dosage of 3 tablets eight-hourly for six days gave good results in the treatment of shigellosis in Cyprus. Chloramphenicol with sulphaphthalidine in combination has also been used successfully; other combinations of antibiotics with soluble and insoluble sulphonamides have their advocates. These and similar combinations possibly lessen the risk of development of sulphonamide-resistant strains of *Shigella*.

In the most severe forms of infantile gastro-enteritis of African children, much of it due to *Shigella* infection, a combination of streptomycin (20 mg./kg.) and chloramphenicol (40 mg./kg.) orally proved very effective as a chemotherapeutic agent, when adequate steps were promptly taken to rectify loss of fluid and electrolytes owing to the disease.

Supportive Treatment

In fulminating attacks of bacillary dysentery, as in other grave infective disorders, early supportive treatment with corticosteroids or gamma-globulin may carry a patient on long enough for the specific treatment to control the infection, and thereby be a life-saving measure. In no way does such treatment replace the specific treatment; the latter must not be delayed when the former is employed.

Dehydration is an accompaniment of severe bacillary dysentery; it is of particularly serious import in children. Therefore in the treatment of children, when there is evidence of its need, fluid additional to the oral intake should immediately be given parenterally to remedy the deficit of fluid and salt, to offset the continuing loss of these, and to meet the child's normal requirements. The estimated daily requirement of a normal child under 6 months of age is 70 ml. of fluid per lb. body weight (150 ml./kg.); up to 1 year, 60 ml./lb. (125 ml./kg.) body weight. At 3 years a total of 1 to 1.5 litres a day is needed, at 8 years 1.5 to 2 litres, and over 10 years at least 2 litres daily. A child moderately dehydrated has lost 5%, a severely dehydrated child 10%, and a very gravely dehydrated child up to 20% of its body weight in fluid. Thus a severely dehydrated child weighing 9 kg., whose estimated normal weight is 10 kg. (22 lb.), requires prompt infusion of 1 litre of fluid to correct his existing deficit. Dehydration in children demands immediate and adequate remedy; of itself it is a major cause of infant death. As an emergency measure 600 ml. of fluid can safely be given subcutaneously to a clinically dehydrated child suffering from gastro-enteritis or dysentery when it is first seen.

Fluid, under suitable conditions, is best given by drip infusion intravenously; for a child it should be in the form of one-fifth to one-half strength normal (physiological) saline, or similar-strength compound sodium lactate injection (*B.P.* 1958) (Hartmann's solution), with 5% added dextrose in each case. The infantile kidney does not excrete salt as efficiently as does the adult; overloading of the child with salt must therefore be avoided; excessive infusion can likewise be disastrous. If intravenous treatment is not possible, these solutions can be infused subcutaneously with hyaluronidase. They also should be given orally, and are palatable to a child. The efficient re-establishment and maintenance of fluid and electrolyte balance in children are matters needing judgment and clinical experience, and it is not so amenable to control by laboratory estimations as it is in adults.

Complications

The common complications of bacillary dysentery are inflammatory lesions of certain organs, notably the eyes, joints, and peripheral nerves; they are attributed to absorption of the toxins. They can be forestalled by early specific treatment. If they do occur they subside under palliative treatment without permanent defect.

Prophylaxis

There is no vaccine or similar protective inoculation against infection. Under very special conditions over limited periods of risk sulphonamides have been given to afford protection against the disease; this practice is undesirable other than as a most exceptional measure.

The Biological Journal is to be published twice a year by the Biological Society of the University of St. Andrews, and is a successor to a magazine first published by the Society for its own members in 1952. It will present reports of recent research and articles of general interest on all aspects of biology for both amateur and professional. Individual copies cost 2s. 6d.; details may be obtained from *The Biological Journal*, 67, Arden Street, Edinburgh, 9, Scotland.

THE TREATMENT OF PSYCHOPATHS*

BY

P. D. SCOTT, M.D., D.P.M.

Consultant Physician, Maudsley Hospital, London

Psychopathy is so ill defined that it is essential, before considering treatment, to develop some clinical classification. The first part of this paper is directed to that end.

If we consider the three best-known definitions—those of Henderson (1939), East (1949), and the Mental Health Act (1959), we find four, and only four, common elements, which may be set out as follows:

	Excluding Clause	Time Factor	Description of Behaviour	Requiring Treatment
Henderson (1937)	Not amounting to mental defect	Throughout their lives or from a comparatively early age. Usually recurrent or episodic	Antisocial or asocial conduct	Social, penal, and medical
East (1949)	Non-insane, psychoneurotic or mentally defective	Persistently	Unable to adapt to social requirements on account of abnormal peculiarities of impulse, temperament, or character	Specialized medical and rehabilitative treatment instead of or in addition to ordinary methods of punishment
Mental Health Act, 1959	Whether or not including subnormality of intelligence (i.e., independent of sub-normality and mental illness)	Persistent	Abnormally aggressive or seriously irresponsible conduct	Requires or is susceptible to medical treatment

We are therefore justified in considering the psychopath as one who does not fit readily into other psychiatric categories, who is persistently antisocial or asocial, and who needs specialized treatment. As soon as we try to go further, in what can be observed now, in the past, or in matters of prognosis, we are in difficulties.

The follow-up studies of Gibbens *et al.* (1959) have shown prognosis, even in aggressive psychopaths, to be very uncertain, in respect not only of frequency but also of quality of offences.

To look back into the earlier history for confirmation of the diagnosis is even less reliable. "Even in childhood," writes Noyes (1939), "the future psychopath usually shows signs of unwholesome personality traits." But the behaviour traits which he then instances are so very common in the histories of those who are never considered psychopathic that we are not afforded much help.

Many attempts have been made to link psychopathy with coexisting specific personality qualities. There is the danger, however, that qualities common to a wide range of non-psychopathic individuals will take on a spuriously ominous meaning when observed in clinics, remand homes, or prisons, and that such backgrounds might be causing or uncovering the qualities. Quoting Noyes again, the psychopath "often conceals his own motives from himself by a superficial rationalization";

*Modified from a paper read to the Royal Medico-Psychological Association in October, 1959.