Dehydration is an accompaniment of severe bacillary dysentery; it is of particularly serious import in children. Therefore in the treatment of children, when there is evidence of its need, fluid additional to the oral intake should immediately be given parenterally to remedy the deficit of fluid and salt, to offset the continuing loss of these, and to meet the child's normal requirements. The estimated daily requirement of a normal child under 6 months of age is 70 ml. of fluid per lb. body weight (150 ml./kg.); up to 1 year, 60 ml./lb. (125 ml./kg.) body weight. At 3 years a total of 1 to 1.5 litres a day is needed, at 8 years 1.5 to 2 litres, and over 10 years at least 2 litres daily. A child moderately dehydrated has lost 5%, a severely dehydrated child 10%, and a very gravely dehydrated child up to 20% of its body weight in fluid. Thus a severely dehydrated child weighing 9 kg., whose estimated normal weight is 10 kg. (22 lb.), requires prompt infusion of 1 litre of fluid to correct his existing deficit. Dehydration in children demands immediate and adequate remedy; of itself it is a major cause of infant death. As an emergency measure 600 ml. of fluid can safely be given subcutaneously to a clinically dehydrated child suffering from gastro-enteritis or dysentery when it is first seen.

Fluid, under suitable conditions, is best given by drip infusion intravenously; for a child it should be in the form of one-fifth to one-half strength normal (physiological) saline, or similar-strength compound sodium lactate injection (*B.P.* 1958) (Hartmann's solution), with 5% added dextrose in each case. The infantile kidney does not excrete salt as efficiently as does the adult; overloading of the child with salt must therefore be avoided; excessive infusion can likewise be disastrous. If intravenous treatment is not possible, these solutions can be infused subcutaneously with hyaluronidase. They also should be given orally, and are palatable to a child. The efficient re-establishment and maintenance of fluid and electrolyte balance in children are matters needing judgment and clinical experience, and it is not so amenable to control by laboratory estimations as it is in adults.

Complications

The common complications of bacillary dysentery are inflammatory lesions of certain organs, notably the eyes, joints, and peripheral nerves; they are attributed to absorption of the toxins. They can be forestalled by early specific treatment. If they do occur they subside under palliative treatment without permanent defect.

Prophylaxis

There is no vaccine or similar protective inoculation against infection. Under very special conditions over limited periods of risk sulphonamides have been given to afford protection against the disease; this practice is undesirable other than as a most exceptional measure.

The Biological Journal is to be published twice a year by the Biological Society of the University of St. Andrews, and is a successor to a magazine first published by the Society for its own members in 1952. It will present reports of recent research and articles of general interest on all aspects of biology for both amateur and professional. Individual copies cost 2s. 6d.; details may be obtained from *The Biological Journal*, 67, Arden Street, Edinburgh, 9, Scotland.

THE TREATMENT OF PSYCHOPATHS*

BY

P. D. SCOTT, M.D., D.P.M.

Consultant Physician, Maudsley Hospital, London

Psychopathy is so ill defined that it is essential, before considering treatment, to develop some clinical classification. The first part of this paper is directed to that end.

If we consider the three best-known definitions—those of Henderson (1939), East (1949), and the Mental Health Act (1959), we find four, and only four, common elements, which may be set out as follows:

	Excluding Clause	Time Factor	Description of Behaviour	Requiring Treatment
Henderson (1937)	Not amounting to mental defect	Throughout their lives or from a comparatively early age. Usually recurrent or episodic	Antisocial or asocial conduct	Social, penal, and medical
East (1949)	Non-insane, psychoneurotic or mentally defective	Persistently	Unable to adapt to social requirements on account of abnormal peculiarities of impulse, temperament, or character	Specialized medical and rehabilitative treatment instead of or in addition to ordinary methods of punishment
Mental Health Act, 1959	Whether or not including subnormality of intelligence (i.e., independent of sub-normality and mental illness)	Persistent	Abnormally aggressive or seriously irresponsible conduct	Requires or is susceptible to medical treatment

We are therefore justified in considering the psychopath as one who does not fit readily into other psychiatric categories, who is persistently antisocial or asocial, and who needs specialized treatment. As soon as we try to go further, in what can be observed now, in the past, or in matters of prognosis, we are in difficulties.

The follow-up studies of Gibbens *et al.* (1959) have shown prognosis, even in aggressive psychopaths, to be very uncertain, in respect not only of frequency but also of quality of offences.

To look back into the earlier history for confirmation of the diagnosis is even less reliable. "Even in childhood," writes Noyes (1939), "the future psychopath usually shows signs of unwholesome personality traits." But the behaviour traits which he then instances are so very common in the histories of those who are never considered psychopathic that we are not afforded much help.

Many attempts have been made to link psychopathy with coexisting specific personality qualities. There is the danger, however, that qualities common to a wide range of non-psychopathic individuals will take on a spuriously ominous meaning when observed in clinics, remand homes, or prisons, and that such backgrounds might be causing or uncovering the qualities. Quoting Noyes again, the psychopath "often conceals his own motives from himself by a superficial rationalization";

*Modified from a paper read to the Royal Medico-Psychological Association in October, 1959.

this surely might be observed in all of us. Other supposed qualities—for example, “a defect in keenness and delicacy of sentiment”—need no further comment. Even the well-known and oft-quoted qualities such as lack of real sincerity and co-operation in treatment (Cleckley, 1941), lack of anxiety and inability to feel guilt, no emotional feelings towards anyone (Bender, 1947), completely egocentric, hedonistic, opportunistic, unable to foresee consequences—all these are very suspect in that they can be observed only in some psychopaths and sometimes in non-psychopaths.

Examples could readily be quoted of persistent serious antisocial behaviour, not related to psychosis or mental deficiency, in which there is a high degree of anxiety and guilt, and a genuine wish for treatment. Lack of guilt, lack of caution or discretion, failure to reckon with consequences, are far more often seen in relation to the offence than as factors pervading the whole personality. “No emotional feelings towards anybody” implies a very rare state. This may be illustrated from experience in approved schools. It is known that a big proportion of chronic adult recidivists, whom most would agree to label psychopathic, have been through juvenile correctional establishments, so that a proportion of the inmates of these establishments must be psychopaths or embryo psychopaths. Yet when the population of an approved school is examined it is found not that the children are unable to make emotional contacts but that they are already so deeply involved in an unsatisfactory, disappointing, or conflictful relationship, usually with the parents, that it hinders them in their other relationships.

Nevertheless it is rare to find young offenders who cannot make a meaningful relationship either with a staff member who sets out to help or, in the very rebellious, with another handicapped or delinquent individual. It is particularly impressive to observe that, in such an institution, individuals who have been considered at staff conferences or in classifying procedures to be “completely egocentric” can suddenly (perhaps on coming into contact with a new member of the staff) develop or uncover precisely opposite qualities. This has its counterpart in the often-cited fact that even apparently intractable psychopaths suddenly settle down. In ordinary psychiatric practice, and indeed in ordinary life, qualities of egocentricity, opportunism, and affective coldness can be observed, without any history or particular likelihood of antisocial behaviour, in individuals who can only be classified as psychopathic if the bounds of the entity are stretched to a very wide and arbitrary extent.

Differences Between Chronic Offenders and Psychopaths

In practice nothing is gained by trying to draw a distinction between psychopathic and chronic offenders. The criteria often depend upon the tradition in the diagnostic service or on the available treatment methods. Psychopaths must be studied in prisons as well as in hospital; at any rate, lessons learnt from psychopaths will be applicable to certain chronic offenders.

It is also difficult to draw the line between aggressive and inadequate psychopaths: they seem to have many more resemblances than differences. Inadequate psychopathy can sometimes be observed to change into aggressive psychopathy, especially in adolescence. It has been shown (Gibbens *et al.*, 1959) that the subsequent offences of “aggressive psychopaths” are often not aggressive.

Inadequate psychopaths sometimes break out into what have been called “catathymic crises” (Wertham, 1959), or autonomous affective crises (Gould, 1959), while in other inadequate psychopaths the motivation and pattern of offences closely resemble those of aggressive psychopaths. Thus, as Menninger (1938) has pointed out, some murders by aggressive psychopaths resemble some suicides by inadequate psychopaths.

Considerations for Research

Throughout the range of antisocial acts, whether set out legalistically or, more simply, as offences of a sexual nature, against property, against persons, or against the State, each may be the final common expression of widely differing individuals and motivation. As a corollary of this, whatever the abnormal or antisocial impulse may be—whether to kill, to steal, to practise some perversion—it will always be possible to find examples of those who do not act it out, of those who might if certain precipitating factors were to act strongly, and of those who do, or would, whenever the opportunity arises. This, together with the hard fact that, of all possible criteria for prediction, nothing seems so reliable as the simple record of offences, suggests that it may be advisable deliberately to study mechanisms of control rather than what is controlled. Possibly the comparison of offenders who have not repeated their offences for 10 years with recidivists would be valuable.

The very useful concept of Mayer-Gross *et al.* (1954) and of Schneider (1958) of psychopathy simply as a deviation from the statistical normal, which therefore includes the study of the whole range of abnormal and unusual personalities, is likely to be so wide as to be an unrewarding or impracticable approach from the point of view of research.

To study specific types of offenders—that is, selected according to their offence—would be useful, but limited for the reasons stated above.

Statistical factorial analysis would seem to be essential but as yet likely to be disappointing, for the reason that the method is dependent on the nature and accuracy of the material fed to it. One of the major obstacles in research will be the difficulty, in many cases the impossibility, of obtaining crucial information; for instance, on the early learning environment.

While the various analytical and deep psychological approaches have been disappointing from the treatment point of view, they have opened up areas of experience of which the importance had never been suspected—for example, the implications of the child-parent relationship. It may be that the immediate value of this approach will be in the choice of questions to be asked and in the technique of obtaining information. Such techniques could be operated most effectively with children whose parents would also be available, as in the Fels investigation into parental attitudes.

As compared with the more objective approaches (psychometric, physiological, anthropological, genetic) the investigation of hidden environmental factors has lagged far behind. Yet it would seem that enough evidence has now accumulated to demonstrate its importance and to warrant the concentration of effort in this direction.

Sources of Material

The available sources of material are the 25,000 individuals in prisons and Borstal institutions, the 8,000

young people in approved schools, the children in special schools under the Education Act ("maladjusted" and some of the educationally subnormal), patients in the mental deficiency and mental hospitals, the three State hospitals, possibly the reception centres (formerly casual wards) where the asocial drifters tend to accumulate.

It is suggested that the approved schools might prove a very valuable place for research operations for these reasons: (1) there is legal provision for detaining them, and for following them up over a statutory period of six years; (2) they are required to pass through (with rare exceptions) a classifying centre where the research team might operate both for initial assessment and for subsequent reassessment during the follow-up periods; (3) their parents will usually be available; (4) these schools are small enough to facilitate the study of every inmate, and sufficiently numerous and differently managed as to vary greatly in the sort of training and treatment provided; and (5) at this age the approved school is likely to be the first long-stay correctional institution which the inmates have experienced, so that there will be less likelihood of secondary "processing" by the penal machinery (a complication which may be very important in corrective training or preventive detention).

Classification of Psychopaths

Before studying the effects of treatment it seems essential to classify the material to be treated; experience shows that this cannot usefully be done by description of behaviour alone, because, as stated above, a piece of behaviour can be the final expression of very different motivations and personalities.

Excluding organic, epileptic, and psychotic illness, the following four subdivisions suggest a practical departure point for such a classification. Each is considered briefly under four headings: (a) the offender's attitude to his offence, (b) observations about the offence, (c) its impact on the observer, and (d) treatment required.

1. Trained, But to Antisocial Standards

Such offenders are behaving as they were taught to behave; they are doing what is "normal" in their families or districts. Some families, consciously or not, train their children to get what they want irrespective of the considerations of others; this is particularly likely to happen in large families, in which the siblings battle for supremacy and are checked from so doing by similar forceful means by their parents. This is what has been called subcultural crime, or "socialized delinquency" (Hewitt and Jenkins, 1946). A more subtle variant is where the parent is obtaining a vicarious satisfaction from the misbehaviour, as described by Johnson and Szurek (1952).

(a) The offender has learnt his behaviour consistently, he will not feel guilt about doing what he was taught to do, his family or circle will not be critical. This sort of crime is not inconsistent with a well-formed character in the individual.

(b) The offence is strictly goal-motivated, intended to gain something materially or as mastery or prestige.

(c) The observer may easily understand the behaviour and feel sympathy in the sense of finding it possible to identify with the offender and his setting.

(d) The treatment required is retraining; in its simplest form punishment is enough, especially if detection is certain and if the others of the subculture are treated at the same time. As with any punishment, it is likely to be most

effective if not too severe or too lenient, and if there is some complementary "advice, assistance, and befriending" to make sure that the offender understands the wider implications of his behaviour and the rationale of society's reaction to it. Because the offender's personality is intact, his learning process is unfettered, and there will be no positive defence against learning another pattern of behaviour.

2. Reparative Behaviour

The offence here is part of an intelligently, laboriously, and often unconsciously worked out policy aimed at adjustment of the individual to difficulties in his environment or to the personal handicaps which that environment has produced in him. For example, some offenders are compensating through their crimes for a sense of inadequacy and inferiority; they tend to be assertive, arrogant, sensitive to slight, and are predisposed to defensive assaults, perhaps to breaking and entering, or to stealing powerful objects, especially cars and motor-cycles. Some sexual offenders—for instance, the homosexual—may fall into this category; the child learns that, in some way, women are dangerous, and therefore substitutes some other sexual outlet. Rado (1949) has used the term reparative in his classification of sex offenders.

(a) Such an offender is identified with his pattern of misbehaviour, even proud of it, and may value it because it protects him from something more dangerous. He may feel guilt about himself, may neglect consequences, but is unlikely to feel guilt in relation to the offence. His personality is to some extent defective.

(b) The offence is distinctly goal-motivated, gaining something in the emotional field though there may be secondary material advantages which may come to take an important part. The offences are not rigidly stereotyped, though habit formation can make them so.

(c) The trained observer can readily understand and sympathize.

(d) Treatment again is retraining; but here there will always be a positive defensive system opposed to treatment, so that some form of psychotherapy will be required. It is often possible for the individual, spontaneously or with help, to change the reparative pattern to one which does not conflict with the law, so that resolution of the personality defect is not always necessary; this is why punishment sometimes works even in this category.

3. The Untrained Offender

Those in the above categories have learnt effectively; in this category they have not. Because there have been no steady standards of behaviour provided, or because there has been a lack of inducements (material or affectionate) to learn, the resulting adult is correspondingly without standards and weak in character. These are the individuals whose behaviour or conduct difficulties show themselves diffusely and from early childhood. They are likely to have been handled in an inconsistent manner with an unpredictable succession of punishment, connivance, or encouragement either from the one parent or as between several adult figures.

(a) The offender often genuinely wishes he could resist his impulses, tends to be confused by punitive measures, but lacks the ability to act with firmness and consistency.

(b) The offences are goal-motivated, often suggested by others, likely to be mixed in type, certainly not stereotyped, and not maladaptive—they could have "come off," and often do; when they do they are enjoyed.

(c) Again the pattern is understandable in terms of the weak personality and specific background of the offender.

(d) Treatment consists of long-term patient supervision with a background of sanctions. Here, especially, heavy punishment tends to lead to the fourth category, and thus

to defeat its objective. In mild cases ambulant supervision and supportive treatment may be sufficient (psychotherapy is not very effective), but in severe cases a sheltered environment with gradual reintroduction of responsibility is indicated.

4. Rigid Fixations

These are the individuals in whom frustration has been so overwhelming, probably at a critical point in their learning process, that learning has broken down and been replaced by a fixed, "maladaptive" pattern of response. The theory rests on ethological observations such as those of Maier (1949) on rats which, when forced to act in a no-solution situation, developed a stereotyped persistent pattern of response even though it resulted, every time, in punishment. Patrick (1934) observed human subjects in a comparable experimental situation, and found that they, too, gave up rational searching for a solution and fell back upon a stereotyped pattern of behaviour. Marquart's (1948) students showed similar maladaptive behaviour when faced with no-solution problems. These fixations persist in animals and humans after "vacations" outside the experimental situation.

Theoretical as these considerations may be, it is a matter of direct observation that there are certain offenders who, from an early age, show a stereotyped, non-adaptive, and non-goal-directed pattern of behaviour which persists unaltered despite long periods within a controlled environment, and is utterly unimproved by even severe punishment. Furthermore, the background always shows early experience of "frustration" consistent with the theory.

(a) This offender often helplessly dissociates himself from his behaviour, but may be genuinely remorseful and anxious about it. Despite attempts to rationalize it, he tends to feel bewildered about it, and the more intelligent covertly believe themselves to be insane.

(b) The offence is stereotyped, gains nothing but relief from immediate tension, and seems out of context with the situation—for example, in the degree of violence expressed.

(c) The observer cannot feel sympathy with the offender or the offence which is outside his experience. The behaviour appears to be compulsive but is not so, for it is not resisted.

(d) Little is known about the treatment, but experience confirms that punishment is useless. It is certainly possible to prevent the building up of the "no-solution" situation, and it is useful to find out to what extent the frustrating precipitating situation is specific or general. This is particularly important for those who are confined within institutions, for it may frequently be the case that, for example, in one-sex institutions the specific frustrating experience never arises, so that it is impossible, in that institution, to help the individual to overcome his particular difficulty.

Theoretical Consideration

To return to theoretical consideration, the rats could be cured of their fixation by "manual guidance," and, concerning humans, Marquart (1948) writes: "Guidance is of value for the elimination of an abnormal fixation but not a habit, and continued failure or punishment eliminates a habit but not a fixation." One supposes that the specific frustrating situation should be reproduced and the individual should be shown how to cope with it by the therapist at his elbow, in the situation with him. Perhaps this is what Stürup (1952) does when holding a person indeterminately in Herstedvester, or Maxwell Jones *et al.* (1952) when holding a person within a critical group situation.

At all events we should take account of Marquart's (1948, p. 137) remark: "Resistance should be handled by attempts to determine its source rather than by pressure designed to break it." Ashley Weeks (1958) wrote: "The idea that one can control future behaviour in a desired direction by setting a person apart and doing something to change him is a fallacy exploded by the recidivist record." Control is the operative word here, and we are bound to agree with him; but with a different approach there is no theoretical reason why offenders of this type should not be altered in such a way as to modify their stereotyped behaviour.

The difficulty, of course, with any such elementary classification as this, is that a given offender almost invariably falls awkwardly into several of the categories. It may well be that the third category (the untrained) is almost invariably combined with the fourth (the fixed patterns). And, of course, the second category (reparative patterns), containing as it does the whole range of antisocial neuroses and antisocial character disorders, is grossly oversimplified. Yet such a scheme fits many clinical facts, and may prevent wasteful and fruitless attempts at assessing treatment methods on material of quite different types. Bearing in mind some such classification, it may now be useful to review the available therapeutic methods.

Available Therapeutic Methods

Our treatment methods are as yet so ineffective that the primary need, not to do further harm, still requires emphasis. Whenever possible, therefore, psychopaths should be kept out of custody, for detention carries risks of its own. As soon as offenders or the antisocially inclined are segregated there is the tendency for staff and inmates to consolidate at opposite poles; a hierarchy tends to develop among the offenders; a threat is thus offered to the staff which calls out a repressive authoritarian regime and the possibility of a vicious circle of resentment and counter-resentment.

The standard methods of combating this are summarized in the following headings brought up in discussion by a group of approved-school headmasters in Bristol: A full programme of constructive and recreational activities in which the staff participate; this often means a succession of not-too-long-term "projects," so that the individuals can experience some return for their efforts. Some hardship or real lack in an institution is a great boon, provided that it is within the community's capacity to correct; this of course was well known to all the pioneers but not, apparently, to the rich authorities who tend to supply their "problem" children on a lavish material scale. Divide the inmates into as many different sorts of groups as practicable; in some the staff should be active, but in others strictly in the background; every member of the community must be involved.

There should be opportunity to air complaints and grievances in free discussion groups (Fenton, 1957; *Lancet*, 1959) (one headmaster described what he called "natter-groups" for his junior approved-school boys). Some self-determination should be entrusted to the inmates, depending on the maturity and morale of the group. Staff should be of both sexes and should be kept keen and tolerant through staff meetings, access to those with greater experience or a different approach from

*Natter "is a Scots word of long standing, probably derived from the Icelandic *knetta*, to grumble" (*Brewer's Dictionary of Phrase and Fable*, p. 640).

theirs, exchanges with other establishments, research projects, and refresher courses. Outside contacts of inmates and staff with the community should be encouraged and not made difficult by geographical isolation.

Such methods are very generally, if somewhat unevenly, found in approved schools, Borstal institutions, and prisons. They should be looked upon as essentials for community living, to which specific methods of treatment may be added.

Greater use of and experimentation with hostels would reduce the population and length of stay in closed institutions.

Confining our further considerations to treatment methods within hospitals, prisons, and other remedial institutions, we have first to consider the commonly held view as expressed, for example, by Dr. R. Bennett (1959) in the House of Commons, that the only treatment for the psychopath is custodial and the lapse of time, "ageing in custody," as he put it. This is unlikely to be true if only by virtue of the great variation within the category of psychopaths. For those individuals for whom it is true two necessities arise: to maintain as healthy and productive a regime as possible, and to make certain that the age at which release could take place is not overshot; this would best be done by grading of security within the institution, or range of institutions, with a flexible means of transfer from one grade to the other, and by having an effective aftercare system. Much devoted work in institutions is thrown away through lack of aftercare.

The remaining principles of treatment are the familiar methods of individual psychotherapy, often difficult to apply to psychopaths, especially in prison, but maintaining a useful place. In a valuable paper, Mackwood (1949) describes the careful piecing together of the patient's history—the therapist, as it were, holding a mirror before him and enabling him to see his misbehaviour as part of his life pattern, a method which one would suppose to be useful, especially in the milder forms of our third category.

The difficulty is often in getting patients, especially those in the second (reparative) category, to the point at which they will co-operate. To this end the indeterminate sentence has been held to be of value or the indispensable corner-stone of treatment as in Scandinavia (Stürup, 1952; Aungle, 1959) and in some of the States of the U.S.A. (Boslow *et al.*, 1959). Undoubtedly the indeterminate sentence is valuable, but it would seem to be of doubtful value for the fourth (fixed) category—the one most likely to call it into operation.

Stürup (1959) describes his dynamic growth therapy and the utilization of "affective moments" (spontaneous catharsis) in changing a patient's point of view, while Maxwell Jones *et al.* (1952) use group pressures to bring a new awareness of his behaviour to the patient: both methods may be applicable to the second, possibly also to the third and fourth, of our categories.

Non-directive therapy (Rogers, 1942; Snyder *et al.*, 1947), group-guided interaction (Weeks, 1958), psychodrama (Moreno, 1946), and closed-group analytical psychotherapy in prison (Landers *et al.*, 1954) are now familiar terms indicating a considerable accumulation of experience, but largely unassessed in their efficacy.

Therapy based directly on Pavlovian theory—for example, Wolpe's (1958) therapy by reciprocal inhibition,

or Raymond's (1956) aversion therapy for sexual perversion—may have a useful place, but is unlikely to be effective with the fourth category.

Pharmacological (oestrogens, disulfram, sedatives) and surgical (castration, leucotomy) treatment of physically normal individuals is favoured in some quarters, and may be regarded as another form of physical rather than psychological restriction or control, and is thus open to all the shortcomings of such methods. Such measures should not be confused with the pharmacological or surgical treatment of specific psychiatric conditions—for example, epilepsy or brain damage—which is an entirely different matter.

This may be the point at which to acknowledge that, with psychopaths, highly refined psychotherapeutic procedures, applied by medical men, are often no more successful, sometimes less successful, than the simpler and less esoteric approaches of certain social workers, probation officers, institutional staff, who may have had little or no training apart from on-the-job experience. Many psychiatrists with experience of working in remedial establishments would agree that a sympathetic member of the staff capable of winning the individual's respect and able to be with him when day-to-day crises appear is able to effect more than a psychotherapist who sees that individual at stated times. It is an established fact that patients who leave hospital with a bad prognosis after failing to improve as regards their psychopathy may then do well under probationary supervision. Such matters need further investigation; it may be that those in category four (the fixed patterns) are more responsive to this direct guidance from someone at their elbow and sharing the experience than they would be to many psychotherapeutic procedures. Some workers intuitively obtain good results with certain psychopaths; it should be possible to find out how they do it.

Conclusion

All forms of treatment of psychopaths are as yet inadequately, or not at all, assessed, and, furthermore, it would seem that a variety of methods are being applied without adequate assessment of the individuals to be treated.

Some such classification as put forward here and applied uniformly in observation and classifying centres (Scott, 1959) would seem essential for further progress.

REFERENCES

- Aungle, P. G. (1959). *J. ment. Sci.*, **105**, 428.
- Bender, L. (1947). In *Handbook of Correctional Psychology*, edited by R. M. Lindner and R. V. Seliger, p. 360. Philosophical Library, New York.
- Bennett, R. (1959). *Hansard*, January 26, 1959, col. 783. H.M.S.O., London.
- Boslow, H. M., Rosenthal, D., and Gliedman, L. H. (1959). *Brit. J. Delinq.*, **10**, 5.
- Cleckley, H. (1941). *The Mask of Sanity*. Mosby, St. Louis.
- Dollard, J., Doob, L. W., Miller, N. E., Mowrer, O. H., and Sears, R. R. (1939). *Frustration and Aggression*. New Haven, Yale Univ. Press, New Haven.
- East, Sir Norwood (1949). *Society and the Criminal*, p. 41. H.M.S.O., London.
- Fenton, N. (1957). "An Introduction to Group Counselling in State Correctional Service." State of California Dept. of Correction Training Manual, VII.
- Gibbins, T. C. N., Pond, D. A., and Stafford-Clark, D. (1959). *J. ment. Sci.*, **105**, 108.
- Gould, J. (1959). In *Recent Progress in Psychiatry*, edited by G. W. T. H. Fleming and A. W. Walk, 3, 317. Churchill, London.
- Henderson, D. K. (1939). *Psychopathic States*. Norton, New York.
- Hewitt, L. E., and Jenkins, R. L. (1946). *Fundamental Patterns of Maladjustment*. State of Illinois.
- Johnson, A. M., and Szurek, S. A. (1952). *Psychoanal. Quart.*, **21**, 323.

- Jones, M., *et al.* (1952). *Social Psychiatry*. Tavistock Publications, London.
- Lancet, 1959, 2, 1022.
- Landers, J. J., Macphail, D. S., and Simpson, R. C. (1954). *J. ment. Sci.*, 100, 953.
- Mackwood, J. C. (1949). *Brit. J. Psychol. (Gen. Section)*, 40, 5.
- Maier, N. R. F. (1949). *Frustration: The Study of Behavior Without a Goal*. McGraw-Hill, New York.
- Marquart, D. I. (1948). *J. Gen. Psychol.*, 39, 107.
- Mayer-Gross, W., Slater, E., and Roth, M. (1954). *Clinical Psychiatry*, p. 94. Cassell, London.
- Menninger, K. (1938). *Man Against Himself*. Harrap, London.
- Mental Health Act (1959). Section 4 (4). H.M.S.O., London.
- Moreno, J. (1946). *Psychodrama*. Beacon Press, New York.
- Noyes, A. P. (1939). *Modern Clinical Psychiatry*, 2nd ed., p. 505. Saunders, Philadelphia.
- Patrick, J. R. (1934). *J. comp. Psychol.*, 18, 1, 153.
- Rado, Sandoz (1949). In *Psychosexual Development in Health and Disease*, edited by P. H. Hoch and J. Zubin, p. 159. Grune and Stratton, New York.
- Raymond, M. J. (1956). *Brit. med. J.*, 2, 854.
- Rogers, C. R. (1942). *Counseling and Psychotherapy*. Houghton, Boston.
- Schneider, K. (1958). *Psychopathic Personalities*. Cassell, London.
- Scott, P. D. (1959). *Brit. J. Delinq.*, 9, 298.
- Snyder, W., and others (1947). *Casebook of Non-directive Counseling*. Houghton, Boston.
- Stürup, G. K. (1952). *Brit. J. med. Psychol.*, 25, 31.
- (1959). *Teamwork in the Treatment of Psychopathic Criminals*. Herstedvester.
- Weeks, A. (1958). *Youthful Offenders at Highfields*, p. 130. Univ. Michigan Press, Ann Arbor.
- Wertham, F. (1947). *Dark Legend*. Gollancz, London.
- Wolpe, J. (1958). *Psychotherapy by Reciprocal Inhibition*. Stanford Univ. Press, California.

HORMONES AND CELLS

[FROM A SPECIAL CORRESPONDENT]

The Society for Endocrinology's symposium on "Cell Mechanisms in Hormone Production and Action" was held in London on May 3 and 4. By its terms of reference the symposium was to be an ordered inquiry into the synthesis and transport of hormones and the mechanisms whereby their influence is exerted in the cells of the target organ; its premise was that the hormone could be regarded as the bearer of partially interpreted genetic information.

Genetic Control

The importance of genetic control of hormone production was borne out by the content of the first two papers. W. R. TROTTER pointed out that the defects of single-gene mutations were well seen in certain forms of thyroid defect in man, notably in Pandred's syndrome, which could be shown to be inherited with a typical Mendelian 3:1 ratio. The condition is attributed to an inability of the thyroid gland to concentrate and take up iodine from the circulating blood, a fault that in turn is possibly due to a deficiency in the production of the enzyme peroxidase. Four other steps in thyroid-hormone synthesis have become recognized as gene-determined, and for one of these—the step involving the removal of iodine from thyroxine—the relevant enzyme "deiodinase" has actually been isolated.

A. C. ALLISON emphasized the central position of the protein enzymes and protein hormones in mediating gene effect. He maintained that genic influence found expression basically in two ways: in controlling amino-acid substitution in protein chains, and in making possible the occurrence of polymerization or aggregation of protein molecules. The small yet highly significant variations in the arrangement of amino-acids in some pituitary hormones could well have been evolved through the ages as the result of mutation of specific genes. Genic control would be exerted during the synthesis of the enzyme or the protein hormone, and the cell structures most probably involved in these syntheses were identified as the minute R.N.A.-containing granules lining the cytomembranes, granules now referred to as ribosomes.

Studies in the fine structure of the cell revealing the varied distribution and abundance of ribosomes in different cells were described by SYLVIA FITTON JACKSON, while P. N. CAMPBELL provided impressive biochemical evidence of the part played by rat-liver ribosomes in the synthesis of serum albumin. But ribosomes, it seems, do not function alone, and some part in synthesis is played by the cytomembranes, and probably other structures too, notably the mitochondria. Indeed, as ROSALIND V. PITT-RIVERS pointed out, though thyroxine synthesis *in vitro* can occur in tissue slices and even in isolated cells, homogenates produced only moniodotyrosine, which suggests that in some way the organization of the living cell as a whole is essential to the process. SYLVIA FITTON JACKSON suggested that secretion of substances of higher molecular weight than could reasonably be held to pass through the plasma membranes might take place through the production of relatively simple components which aggregated on reaching the outer surface of the cell. Variations of cell surface with function were mentioned by J. K. GRANT, who described how stress-invoked changes in adrenal cortical cells included the formation of microvilli, probably associated with increased hormone secretion.

The forms taken by some hormones during transport in the blood-stream were discussed by some speakers. I. H. MILLS reported that cortisol, and probably also other steroid hormones, are largely bound to plasma proteins during transport, and the same, according to J. R. TATA, was true of the thyroid hormones. In both instances the α -globulin fraction of the plasma was implicated.

Do Hormones Enter Cells?

When the hormone reaches the target cells it must somehow exert its effect on or through the cell surface. The need for a clearer understanding of the nature of the cell surface was stressed by L. WEISS, who argued that heterogeneity in cell-surface structure was probably much greater than commonly realized and that actual entry of the hormone into the cell need not necessarily occur; information could conceivably be imparted through a molecular reorientation process, possibly at specific sites on the surface.

Whether or not the hormones enter cells, various factors may modify their influence. F. G. YOUNG noted that the degree of oxygenation of the blood apparently affected the activity of the circulating insulin, a relationship that might hold with other hormones also. Metabolism of thyroid hormones yielded simpler compounds which still had the capacity to stimulate metabolism, but J. R. TATA believed that the true hormonal influence is carried only by thyroxine and triiodothyronine.

Hormones and Cell Metabolism

Several reports were concerned with the mode of action of hormones on cell metabolism. The action of insulin was considered by G. HOWARD SMITH to be essentially that of increasing the permeability of cells to glucose; he postulated a sugar-carrying mechanism in the cell which was controlled through inhibition by adenosine triphosphate, the action of insulin being to inhibit A.T.P. inhibition. A. KORNER, K. L. MANCHESTER, and E. REID described excellent independent investigations bearing upon the action of various hormones on amino-acid incorporation, protein synthesis, and components of protein-synthesizing systems. Effective agents were growth hormone, A.C.T.H., insulin, L-thyroxine, and certain corticosteroids.

These studies have the special significance that they are concerned with ways in which hormones might influence the production of hormones, and thus may ultimately provide a biochemical explanation for the integration of the endocrine system.

The mode of action of oestrogens remains as deep a mystery as ever, but the painstaking studies with oestrogen antagonists, described by C. W. EMMENS, hold promise of yielding valuable clues in the near future.