The extreme degrees of oedema seen in some of the patients (especially the children with glomerulonephritis) were thought to be associated with gross hypoalbuminaemia. The cause of this was partly albuminuria, which was usually heavy (in one child, loss of urinary protein reached a level of 22 g./litre at a time when the serum albumin level was 0.49%). In some patients the serum albumin level continued to fall after admission to hospital, presumably because the patients were admitted early in the course of their illness before the drain of albumin in their urine had fully depleted the serum protein level. Another factor in the development of gross oedema may have been that the serum protein levels of even apparently healthy Africans in Uganda are different from European normals (Holmes, Stanier, Semambo, and Jones, 1951), the albumin levels being lower and the globulin levels higher than in Europeans. The serum albumin levels in these patients, especially in the children, were therefore probably low before the urinary loss began.

Summary

Thirty-three African patients suffering from glomerulonephritis were studied. In all but the youngest cases the diagnosis was confirmed by renal biopsy.

The 12 children in the series presented with massive oedema and ascites associated with heavy albuminuria. Haematuria was only microscopical. A rise in blood-pressure was common. The serum albumin level was constantly low and the serum cholesterol usually raised. Electrophoresis of serum protein characteristically revealed a "nephrotic" pattern. Some rise in the blood-urea level was common in the early stages of the illness. Changes of acute nephritis were generally seen on renal biopsy, though in three out of eight cases in which this was done chronic nephritis was present. The illness ran a protracted course, abruptly curtailed in one case by steroid therapy. The overall prognosis appeared to be poor.

Renal biopsy revealed acute nephritis in 6 of the 21 adult patients. One presented with the nephrotic syndrome and made an apparently good recovery. In two others with a raised blood-pressure, macroscopic haematuria, and a raised blood urea, some improvement took place in hospital. In the remaining three the disease was rapidly progressive, two dying in hospital. Severe hypertension, retinopathy, renal insufficiency, and heart failure were characteristic of these cases in which severe changes of recent origin were present on renal biopsy.

Chronic nephritis was present in 10 cases, and in these hypertensive heart failure was the commonest mode of presentation. Advanced retinopathy was usual in the presence of severe hypertension. The blood urea was raised in all cases, and in four patients who died while under observation death was due to heart failure associated with insufficiency.

I wish to acknowledge the constant help and encouragement I received from Professor A. W. Williams, of Makerere College, in whose department this study was made. My thanks are also due to Professor J. N. P. Davies, of Makerere College, and Professor T. F. Hewer, of Bristol University, for assistance in the interpretation of histological specimens; and to Professor C. Bruce Perry, of Bristol University, for advice in the preparation of the paper. J. Kyobe, of Makerere College, acted as interpreter throughout the

study and gave much technical assistance; and Miss J. Summerscales, of the Uganda Medical Service, carried out the electrophoretic studies. Photomicrographs were prepared by Mr. D. N. White, of Bristol.

REFERENCES

- Becker, B. J. P. (1946). *S. Afr. J. med. Sci.*, **11**, 107.
 Carothers, J. C. (1934). *E. Afr. med. J.*, **10**, 335.
 Davies, J. N. P. (1947). *Ibid.*, **24**, 223.
 — (1949). *Ibid.*, **26**, 76.
 Donnison, C. P. (1928). *Kenya E. Afr. med. J.*, **5**, 210.
 Ellis, A. (1942). *Lancet*, **1**, 1.
 Furman, K. I. (1955). *S. Afr. med. J.*, **29**, 590.
 Gelfand, M. (1957). *The Sick African*, 3rd ed., pp. 592, 595. Juta, Capetown.
 Giglioli, G. (1932). *Trans. roy. Soc. trop. Med. Hyg.*, **26**, 177.
 Goldie, H. (1930). *Ibid.*, **23**, 503.
 Holmes, E. G., Stanier, M. W., Semambo, Y. B., and Jones, E. R. (1951). *Ibid.*, **45**, 371.
 James, C. S. (1939). *Med. J. Aust.*, **1**, 759.
 Jelliffe, D. B., and Reed, F. S. (1953). *J. trop. Med. Hyg.*, **56**, 33.
 Leather, H. M. (1959). *E. Afr. med. J.*, **36**, 305.
 Luetscher, J. A. (1940). *J. clin. Invest.*, **19**, 313.
 MacInness, D. A., and Longworth, L. G. (1939). *Science*, **89**, 438.
 Manson-Bahr, P., and Maybury, L. M. (1927). *Trans. roy. Soc. trop. Med. Hyg.*, **21**, 131.
 Raper, A. B. (1953). *E. Afr. med. J.*, **30**, 49.
 Shaper, A. G., and Shaper, L. (1958). *Ibid.*, **35**, 647.
 Stückler, G. B., Burke, E. C., and McKenzie, B. F. (1954). *Proc. Mayo Clin.*, **29**, 555.
 Uys, C. J. (1954). *S. Afr. J. clin. Sci.*, **5**, 206.
 Williams, A. W. (1939). *E. Afr. med. J.*, **16**, 2.

ACUTE APPENDICITIS IN PREGNANCY

BY

WILLIAM P. BLACK, M.B., F.R.C.S.ED., M.R.C.O.G.

Lecturer, Department of Obstetrics and Gynaecology,
University of Leeds

Acute appendicitis in pregnancy is a subject about which much has been written in American journals but which has had little attention in British literature. The incidence is between 0.5 and 1 per 1,000 pregnancies, and mismanagement of this condition carries a significant maternal mortality.

The diagnosis of appendicitis is frequently difficult in the pregnant, as in the very young, the very old, and the obese. Particularly in pregnancy is the early diagnosis most important, because the mortality of appendicitis complicating pregnancy and the puerperium is the mortality of delay.

Of the 25 cases of acute appendicitis in pregnancy here presented, 10 occurred in Glasgow Royal Maternity and Women's Hospital between 1942 and 1953 (an incidence of 0.26 per 1,000 births), 3 occurred in Eastern District Hospital, Glasgow (an incidence of 1.4 per 1,000 births), and 8 were treated in the Maternity Hospital at Leeds between 1948 and 1959 (an incidence of 0.36 per 1,000 births). Four were treated in the general surgical wards of Glasgow Royal Infirmary.

From the literature 348 cases of appendicitis in pregnancy were collected, giving a total of 373 cases, including the present series.

Incidence of Acute Appendicitis

In acute appendicitis in general, two-thirds of cases occur between 11 and 30 years of age, and two-thirds are male (Tashiro and Zininger, 1946). One-third of the male patients have perforative appendicitis, compared with one-fifth of the females. Three-quarters of all cases of so-called chronic appendicitis occur between 11 and 30 years of age, and two-thirds of these are females (Shelley, 1938). Appendicitis, therefore, is

a less severe condition in women, resulting in fewer and less advanced cases of acute appendicitis and more cases of chronic appendicitis of doubtful significance. The incidence of pregnancy among women with acute appendicitis is about 2% (Baer *et al.*, 1932).

Acute appendicitis in pregnancy occurs most often in nulliparae and primiparae under 30 years of age, probably as the result of the age incidence of acute appendicitis in general. In the present series 18 patients were under 30 years of age and 16 were in their first or second pregnancy.

Duration of Pregnancy

Whereas the stage of pregnancy affects the appendix because of the elevation of the caecum and appendix caused by the enlarging uterus (Baer *et al.*, 1932), it does not affect the incidence of appendicitis in pregnancy, which is approximately the same in each trimester. The distribution of the total cases according to trimester is shown in Table I, 35.1% occurring in the first trimester, 33.9% in the second, and 30.9% in the third trimester and the puerperium.

TABLE I.—*Stage of Pregnancy*

	Present Series	Literature	Total	Percentage
First trimester ..	2	122	124	35.1
Second „ ..	13	107	120	33.9
Third „ ..	6	89	95	26.9
Labour ..	2	4	6	1.7
Puerperium ..	2	6	8	2.3
Total ..	25	328	353	

In 20 cases from literature no maturity was given.

TABLE II.—*Maternal Mortality*

	Present Series	Literature	Total	No. of Cases	Mortality (%)
First trimester ..	—	—	—	132	0
Second „ ..	—	5	5	126	3.9
Third „ ..	—	11	11	101	10.9
Labour ..	1	—	1	6	16.7
Puerperium ..	—	—	—	8	0
Total ..	1	16	17		4.6
Cases ..	25	348		373	

The stage of pregnancy does affect the severity of the appendicitis. In Table II the maternal mortality is calculated for each trimester. No maternal death occurred in the first trimester, but the maternal mortality is 3.9% in the second, 10.9% in the third, 16.7% intrapartum, and nil in the puerperium. The maternal mortality for the first two trimesters is 1.9%, which has to be compared with the maternal mortality of 10.9% in the third trimester and a mortality rate for acute appendicitis in general of approximately 1% (Aird, 1957). One maternal death occurred in the present series.

The increasing maturity of the pregnancy adversely affects the severity of the appendicitis in three ways. First, the upward displacement of the appendix in pregnancy takes it out of the pelvis into the general peritoneal cavity, where localization of any infection is less likely. The enlarged uterus prevents the omentum from gaining satisfactory and unrestricted access to the site of the infection. The Braxton Hicks contractions of the neighbouring uterus prevent the infection from being quickly localized by the formation of adhesions, and the general increase in vascularity of the abdomen, with a consequent increase in the amount and rate

of lymphatic drainage, results in a more rapid progress of the inflammation.

Secondly, the upward migration of the appendix may be lessened or prevented by periappendicular adhesions, low attachment of the caecum, or a retrocaecal appendix. Hoffman and Suzuki (1954) showed that the mobility of the appendix as noted at operation has a direct influence on the severity of the appendicitis, the retrocaecal appendices and those fixed by adhesions resulting in more severe infection. A previous attack of appendicitis resulting in adhesions is therefore an adverse factor in cases of appendicitis in pregnancy. When the diagnosis of appendicitis has been made in a female, appendectomy should be performed. Subsequent appendectomy should be carefully considered for a patient who has had appendicitis which was not treated by operation in the first instance, because the risk of such an operation is negligible and the risk of any flare-up of appendicitis in a subsequent pregnancy is removed.

Thirdly, the diagnosis of acute appendicitis becomes more difficult as pregnancy approaches term. If the diagnosis is delayed, and perforation and peritonitis follow, the prognosis is made worse by a concomitant pregnancy. There is a high circulatory level of adrenocorticoids in pregnancy which tends to diminish the tissue response to inflammation and so mask the early signs of infection and hinder localization. There is also a general atony of visceral smooth muscle which may cause a diminished reaction of the appendicular muscle coat to infection with a less severe midgut colic of slower onset.

Foetal Loss

The foetal loss in acute appendicitis in pregnancy is considerable, and Table III shows the foetal loss for the series where details are available. The foetal mortality rate was 12.5% for the first trimester, 15.7% for the second, and 20% for the third, with an overall foetal loss of 17%. Premature interruption of pregnancy becomes more frequent as the severity of the appendicitis increases and the nearer the patient is to term, the severity of the appendicitis being more important than the maturity of the pregnancy. Abortion and premature

TABLE III.—*Foetal Loss*

	Present Series	Literature	Total	Mortality (%)
First trimester ..	1	7	8	12.5
Second „ ..	4	4	8	15.7
Third „ ..	3	9	12	20
Total ..	8	45	53	17

labour themselves have little adverse effect on the maternal prognosis. In the present series, five patients aborted and four had premature labours, three infants dying in the neonatal period. This is a foetal loss of 8 in 22 pregnancies (36.4%), since three patients have to be excluded because two developed appendicitis in the puerperium and in one patient the result of the pregnancy is not known.

Diagnosis

The symptoms of acute appendicitis are the same in the gravid as in the non-gravid. Smith and Bartlett (1940) found the commonest and earliest symptoms to be abdominal pain and the commonest sign to be local tenderness over the appendix. Vomiting occurring after the fifth lunar month requires careful investigation.

Physical findings may be minimal after the first trimester, so that the diagnosis becomes progressively more difficult near term. As pregnancy advances, the pain is situated higher and more laterally in the abdomen. In the first three months the clinical picture is no different from the non-gravid, but the elevation of the appendix, which usually occurs thereafter, is associated with a masking of the signs of muscle rigidity, rebound tenderness, psoas spasms, and maximal tenderness in the iliac fossa as elicited by rectal and vaginal examination. Abdominal examination is equivocal in many cases, particularly in the later months. McDonald (1929) stressed the significance of nausea and vomiting associated with pain, fever, and local tenderness.

The patient's temperature is of little help, the absence of any marked pyrexia being in keeping with appendicitis, whereas a rising pulse rate is of importance. The white-blood-cell count in pregnancy is normally 8,700 to 15,500/c.mm. (Priddle and Hesselstine, 1951), and the erythrocyte sedimentation rate is normally raised, with an average of 36 mm. (Westergren) at one hour in cases at term (Hamilton, 1953). Meharg and Loop (1953) placed little value on the white-cell count in cases of appendicitis, and Parker (1954) regarded only a high white-cell count to be significant.

Examination of the urine is important to differentiate pyelitis from appendicitis. In the early stages of pyelitis the urine may be quite clear microscopically despite suggestive symptoms, and elevation of the foot of the patient's bed may be helpful in allowing the flow of urine from the affected atonic ureter to improve. Only a catheter specimen of urine with a large number of pus cells present is an adequate pointer to pyelitis where the diagnosis is in doubt.

In the present series the most important diagnostic factor was the history given by the patient, though the duration of the history did not always coincide with the severity of the appendicitis at operation. White-blood-cell counts of 19,000/c.mm. were associated with appendicitis with perforation and peritonitis, but were not important in the diagnosis.

Treatment

The pregnant woman with acute appendicitis is first a surgical problem and secondly an obstetric problem. During the first two trimesters early diagnosis and immediate appendicectomy are the best treatment, the uterus being manipulated as little as possible during the operation, which should be performed rapidly. After operation every effort should be made to prevent abortion and premature labour by combating dehydration, acidosis, and abdominal infection. A carefully performed abdominal operation not involving the uterus should not cause abortion or premature labour, and removal of an acutely inflamed appendix in pregnancy will diminish the risk of pregnancy otherwise terminating prematurely.

When acute appendicitis is diagnosed in the last trimester, most authorities favour appendicectomy alone and leave the pregnancy to take its own course. Lower-segment caesarean section followed by appendicectomy may allow rapid delivery of the infant, reduction of the time of exposure of the infant to toxæmia, and easier access to the appendix, but it increases the danger of widespread infection of the uterine cavity and reduces the resistance of the peritoneum by additional trauma. Indeed, the removal of an appendix at the

time of caesarean section, of hysterectomy, or of removal of an ectopic pregnancy increases the operative risk. If the patient is in labour at the time of diagnosis, and vaginal delivery can be achieved without much delay, this is by far the best management, the appendix being removed immediately afterwards.

With a patient in early labour without obstetric reasons for abdominal delivery, in whose case it is considered unwise to await vaginal delivery before performing appendicectomy because a long labour is expected, the decision on caesarean section at the time of appendicectomy would have to be made on the merits of the individual case. In this country, to-day, lower-uterine-segment caesarean section is the operation of choice, should abdominal delivery be advisable at the same time as appendicectomy. Theoretically the best procedure would be extraperitoneal caesarean section followed by appendicectomy through an incision over the appendix. However, this would be a longer and more difficult procedure, and should the peritoneum be punctured during the extraperitoneal approach to the uterus the risk of spread of infection would be increased.

In 24 of the 25 cases reported, appendicectomy was performed without other surgical procedure, apart from drainage of the peritoneal cavity, which was decided on accepted surgical principles. In one case a localized appendix abscess was drained without removal of the appendix, this being eventually removed some months after delivery. The fatal case is presented in more detail.

Case 6.—A para-3 aged 28 was admitted in labour with a five-day history of diarrhoea followed by vomiting and upper abdominal pain prior to the onset of labour. On the third day of symptoms the pain became localized in the right iliac fossa. A spontaneous vertex delivery lasting six and a half hours occurred on the fifth day of symptoms and was followed by a marked increase in the severity of the pain in the right iliac fossa. Laparotomy on the sixth day of symptoms revealed a ruptured retrocaecal appendix, acutely inflamed and gangrenous, with generalized peritonitis. The patient died next day despite penicillin and intravenous fluids. During the course of her illness she had a very rapid pulse, rising to 160 a minute, and her temperature on admission was 100.4° F. (38° C.), rising to 102° F. (38.9° C.). Her urine remained microscopically and bacteriologically sterile.

Summary

A series of 25 cases of acute appendicitis in pregnancy are reported. One maternal death occurred, and there were five abortions and four premature labours with three neonatal deaths. Twenty-four cases were treated by appendicectomy, one by drainage of an appendix abscess.

From the literature 348 cases of acute appendicitis in pregnancy were collected. From a study of the total 373 cases the figures shown in Tables I to III were obtained.

Diagnosis, prognosis, and treatment of acute appendicitis are discussed and the following points noted. (1) Appendicitis appears to be no more frequent in pregnancy than otherwise, being commonest under 30 years of age, during the first and second pregnancies, and arising with about equal incidence in each trimester. (2) The condition in general is no more severe in pregnancy than otherwise, unless the diagnosis is delayed, when the inflammation is more acute and less well localized because of pregnancy. The diagnosis is

more difficult the nearer the patient is to term; consequently the risk to the patient is greater near term. (3) The foetal mortality is related to the severity of the appendicitis rather than to the duration of the pregnancy. Abortion or premature labour has little adverse effect on the maternal prognosis. (4) Previous appendicitis with adhesions will adversely affect the prognosis of subsequent appendicular inflammation in pregnancy. When appendicitis is diagnosed in a woman the appendix should be removed. (5) The treatment of choice is appendectomy, without interference with the pregnancy.

I thank my consultant colleagues for their permission to refer to the case-records of their patients, and for the help and advice of Sir Andrew Claye and Miss U. M. Lister in the preparation of this article.

BIBLIOGRAPHY

- Aird, I. (1957). *A Companion in Surgical Studies*, 2nd ed. Livingstone, Edinburgh and London.
- Baer, J. L., Reis, R. A., and Arens, R. A. (1932). *J. Amer. med. Ass.*, **98**, 1359.
- Child, C. G., and Douglas, R. G. (1944). *Amer. J. Obstet. Gynec.*, **47**, 213.
- Cocke, N. P., and Mason, J. M. (1920). *J. Amer. med. Ass.*, **75**, 95.
- Cosgrove, S. A. (1937). *Amer. J. Obstet. Gynec.*, **34**, 469.
- Hamilton, G. M. (1953). *J. Obstet. Gynaec. Brit. Emp.*, **60**, 409.
- Hoffman, E. S., and Suzuki, M. (1954). *Amer. J. Obstet. Gynec.*, **67**, 1338.
- Krieg, E. G. M. (1949). *Ibid.*, **57**, 736.
- McDonald, A. L. (1929). *Ibid.*, **18**, 110.
- Maes, U. (1934). *Ibid.*, **27**, 214.
- Meharg, J. G., and Loop, F. A. (1953). *Obstet. and Gynec.*, **1**, 460.
- Meiling, R. L. (1947). *Surg. Gynec. Obstet.*, **85**, 512.
- Parker, R. B. (1954). *Lancet*, **1**, 1252.
- Priddle, H. D., and Hesselstine, H. C. (1951). *Amer. J. Obstet. Gynec.*, **62**, 150.
- Priest, F. O. (1936). *Ibid.*, **31**, 878.
- Renn, A. C., Douglass, L. P., and Cushman, G. F. (1951). *Ibid.*, **62**, 1343.
- Shelley, H. J. (1938). *Arch. Surg. (Chicago)*, **37**, 17.
- Smith, J. A., and Bartlett, M. K. (1940). *New Engl. J. Med.*, **223**, 529.
- Tashiro, S., and Zininger, M. M. (1946). *Arch. Surg. (Chicago)*, **53**, 545.

A NEW MASK FOR DELIVERING OXYGEN OR OTHER GASES

BY

MARY CATTERALL, M.B., B.S., D.M.R.T.

Research Assistant to the Departments of Medicine and
Thoracic Surgery, General Infirmary at Leeds

The M.C. mask, described below, differs from other masks in that, by virtue of its shape and volume, a valve effect is obtained without the presence of a mechanical valve. Thus there is no rebreathing of carbon dioxide, and a cone of inflowing gas is delivered direct to the region of the patient's nostrils or mouth. The concentration of the inflowing gas can be regulated by the rate of flow from the delivery cylinder. A flow of 2 litres of oxygen a minute gives an average concentration of 35%.

The mask is comfortable, very portable, and does not interfere unduly with talking. It is suitable for the conscious or the unconscious patient and can be worn in the sitting or lying position. As it is made in a cup shape of semi-rigid plastic, it can be held to the face by ambulatory patients who require oxygen quickly for the making of some effort. It is suitable for the administration of other gases—for example, self-administered anaesthesia in labour. It is so constructed as to avoid

the disadvantages of the polymask described in a previous paper (Catterall and Snow, 1960).

The mask is worn as shown in the diagram, and the main part of it consists of a cup-shaped member of semi-rigid plastic through the centre of which runs the gas-delivery tube of the same material. The inner end of this extends to a point in the same plane as the rim of the cup-shaped member, which is in contact with the

Diagram showing M.C. mask in position.

nose and the chin. It is therefore in close proximity with the patient's upper lip, to which the inflowing gas is directed. The outer end of the gas-delivery tube extends sufficiently beyond the dome of the cup-shaped member to attach to it the supply tubing from the gas cylinder.

The sides of the rim are made deeper than the parts which touch the nose and the chin to provide more stability when the mask is worn.

A small hole is situated in the cup-shaped member about halfway between the rim and the gas-delivery tube; this is for the emergence of expired carbon dioxide and, at low rates of flow, for the inspiration of room air.

The cup-shaped member is kept in place by a skirt of thin plastic sheeting in the hem of which is a pipe cleaner. This is moulded over the bridge of the nose and the cheeks and beneath the chin. The covering of cotton fibres prevents any cutting in of the wire to the skin. Adjustable straps around the head fix the mask in position.

Principle of the M.C. Mask

The inflowing gas is delivered very close to the upper lip, which it strikes, and is deflected upwards to the nostrils and downwards to the mouth. In addition to this, on inhalation the inflowing gas is drawn towards the mouth or nostrils. Thus a central cone is produced which contains a high concentration of the inflowing gas. Around this, the atmosphere within the cup-shaped member contains smaller concentrations of the gas, until at the periphery, immediately proximal to the hole