The diets prescribed were basic in character. Their cost could be increased greatly by using special low-sodium preparations or higher-quality protein foods in the appropriate diets. Special diabetic food preparations employed to make individual diets more attractive would naturally add to the weekly expenditure. In the case of the 1,000-calorie diet, the one prescribed was thought to be adequate; the patient's income and standard of living may affect the quality and quantity of permitted foods on this type of diet. It may be that some supplementary foods are essential to make a 1,000-calorie diet less monotonous, with some consequent increase in cost. Financial assistance for this purpose may be sought by the patient from public funds, such as the special nourishment allowance allowed at the discretion of the National Assistance Board.

Those members of the population living on an old-age pension have a special problem with respect to the cost of their normal diet. The pensioner group spend two shillings less than the average weekly expenditure on food; whereas the average weekly expenditure on food is 30% of total expenditure on consumer goods, with the pensioner group this appears to be approximately 56% (National Food Survey Committee, 1959). The prescription of a special diet for pensioners is more likely to entail financial strain than in a normal population. However, financial aid and social services alleviate this strain somewhat. The problem is complicated, firstly, by factors such as pensioners' teeth, and, secondly, by the reluctance and physical inability of many older people to buy and prepare their food.

Summary

The estimated cost of prescribed diets was compared with that of the British Medical Association's recommended normal diet. Diabetic diets in most frequent use, the low-sodium diet, and gastric diets did not appear to entail any extra cost for an adult woman doing a sedentary job. Diabetic diets with a protein content of 100 g. or more, the 1,000-calorie diet, the 100-g. high-protein diet, and the low-animal-fat diet appeared to cost more, but none exceeded the estimated cost for a man doing medium work.

We are grateful to Miss E. Wilkinson-Hughes and Dr. J. D. N. Nabarro for making material for this paper available to us.

REFERENCES

- British Medical Association (1950). *Report of the Committee on Nutrition*. B.M.A., London.
 Ministry of Labour Gazette, 1958, 66, No. 3.
 Monthly Digest of Statistics, 1959, 159, 3.
 National Food Survey Committee (1959). *Domestic Food Consumption and Expenditure: 1957*. H.M.S.O., London.

The World Health Organization has provided Queen's University, Belfast, with a grant to assist in a year-long investigation which will be carried out by the University's department of social and preventive medicine and the Health Department of Belfast Corporation. The aim of the investigation is to find out what part air pollution plays in causing lung cancer. Belfast Corporation's Health Department will use five smoke- and dust-collecting instruments situated throughout the city; the samples collected will be sent for analysis to the air pollution unit of the Medical Research Council at St. Bartholomew's Hospital, London, where the proportion of cancer-producing substances will be determined. An investigation of smoking habits in Belfast is also being planned.

Preliminary Communications

Chlorothiazide in Treatment of Carpal-tunnel Syndrome

While it is now generally agreed that the major cause of the syndrome of acroparaesthesia in the upper limb is compression of the median nerve in the carpal tunnel, there is no such unanimity regarding treatment.

Although surgical decompression is the treatment of choice in long-standing cases with motor and sensory signs, milder cases respond to conservative methods of treatment, such as splinting of the wrist (Heathfield, 1957; Hierons, 1959), local injections of hydrocortisone (Crow, 1960; Foster, 1960), and systemic prednisone (Lees and Liversedge, 1959). None of these methods, however, is entirely satisfactory, for splinting is somewhat clumsy, hydrocortisone injections have a high relapse rate, and systemic steroids may not be of lasting benefit nor are they free from undesirable side-effects—so much so that the major complications which may occur with them should preclude their use in a condition which never threatens life.

In those cases associated with pregnancy there is evidence that tissue-fluid retention plays a part in the aetiology, and it is reasonable to suppose that this is a factor in other patients. In favour of this supposition is the common assertion by patients that their hands are swollen at the time when symptoms are maximal. McCallum (1959) and Wood (1959) have stated that cases of acroparaesthesia in pregnancy have been relieved by diuretic drugs. We thought it would be worth trying the administration of a highly effective diuretic, such as chlorothiazide, in an unselected group of patients with acroparaesthesia; so in October, 1959, we started treating with this drug all patients with the carpal-tunnel syndrome not severe enough to warrant surgical decompression. As the series was collected from a wide area covering several hospitals a double-blind trial was impracticable.

METHOD AND MATERIAL

All patients were initially given chlorothiazide 0.5 g. twice daily for six days in the week, and this was supplemented with potassium chloride 0.5 g. twice a day; they were seen again at the end of a month. In patients who had responded to the drug the dosage was progressively reduced and the cases were followed up until symptom-free for two months; patients were then asked to report back if symptoms recurred. At the end of the investigation all the apparently "cured" cases were contacted for review.

The series comprised 35 cases of typical acroparaesthesia due to compression of the median nerve in the carpal tunnel. As in other published series, there was a great preponderance of women. The age range was from 22 to 60 and was maximal in the decades on either side of 45. Symptoms had been present for a variable time from one month to 15 years, and were mainly nocturnal. Cases with marked signs of median-nerve paralysis were excluded from the series as being more suitable for surgery. Physical signs were therefore confined to minimal weakness of the thenar muscles and slight sensory impairment of median-nerve distribution. In 13 cases there were no abnormal physical signs except for a positive tourniquet test in nine. In no

cases were there any reflex changes in the upper limbs, and, although x-ray films of the cervical spine were taken, evidence of cervical spondylosis was found in only one case.

RESULTS

At the end of one month's treatment 29 out of the 35 patients were either symptom-free or much improved, but six showed no improvement; thus the treatment was initially successful in nearly 90% of cases. In one case treatment was combined with splinting of the wrist. Of the six cases in which the treatment proved unsuccessful four were cured by splinting and two eventually underwent operation. At the end of three months 19 of the 29 patients were still symptom-free, but 10 had relapsed on the treatment.

At a subsequent follow-up, varying from 3 to 13 months, there were three further relapses, so that 16 remained symptom-free without drug therapy. In one of the late relapses chlorothiazide was recommenced and the patient again became symptom-free.

The patients were not weighed as a routine, but 14 were weighed during the course of treatment and striking discrepancies were found; eight lost an average of 5 lb. (2.3 kg.), four gained an average of 10 lb. (4.5 kg.), and in two the weight remained stationary. Thus there was no significant change in the average weights during treatment.

It is evident that many cases of the carpal-tunnel syndrome can be successfully treated with chlorothiazide. There was no ultimate difference between cases with physical signs and those without, though the latter seemed to respond better initially; nor was duration of symptoms of importance in determining response, since two cases with symptoms for 15 years responded.

Patients with purely nocturnal symptoms and those with nocturnal and diurnal symptoms seemed to respond equally well to treatment, but those with continuous symptoms suggesting permanent nerve damage rather than recurrent ischaemia did not respond.

Although improvement appeared to be correlated with diuresis it was surprising to us that so few patients lost weight. This might be explained by the fact that some admitted to drinking more. Removal of oedema beneath the transverse carpal ligament need only be very small in amount to be sufficient to break a recurrent cycle of nerve ischaemia. However, it is possible that improvement may have been due more to redistribution of body fluid than to generalized dehydration. Such an explanation has been put forward to explain improvement with systemic steroid therapy, which may even cause oedema.

Side-effects were few; only two patients complained of slight nausea at the beginning of treatment. We encountered no clinical instances of electrolyte imbalance, nor was postural hypotension observed. Given with a weekly rest day and with a potassium supplement the drug appears to be safe, even when administered over a period of months.

Splinting and chlorothiazide can be combined: this was done successfully in three cases. Combined treatment might well have succeeded in more cases, but was purposely avoided.

Obesity appeared to be of importance in more than a quarter of this series, and many patients commented that symptoms began after a sudden increase in weight.

Adipose tissue in the carpal tunnel might be a precipitating cause in these patients or, alternatively, subcutaneous fat may compress the transverse carpal ligament from outside. It seems possible that treatment might have been successful in more cases had the patients not been simultaneously gaining weight, as these comprised a number of our failures.

K. W. G. HEATHFIELD, M.D., M.R.C.P.,
Consultant Neurologist,
Barnet General, Harold Wood, and Whipps Cross
Hospitals.

J. A. R. TIBBLES, M.B., D.C.H., M.R.C.P.,
Senior House Officer,
West End Hospital for Nervous Diseases, London.

REFERENCES

- Crow R. S. (1960). *Brit. med. J.*, **1**, 1611.
Foster, J. B. (1960). *Lancet*, **1**, 454.
Heathfield, K. W. G. (1957). *Ibid.*, **2**, 663.
Hierons, R. (1959). *Practitioner*, **182**, 590.
Lees, F., and Liversedge, L. A. (1959). *Lancet*, **2**, 1120.
McCallum, M. J. (1959). *Brit. med. J.*, **2**, 1095.
Wood, E. C. (1959). *Ibid.*, **2**, 1254.

The Scribner Arteriovenous Fistula for Haemodialysis

Experience has shown that some patients with acute renal failure reach artificial kidney units with veins inadequate for repeated cannulation for haemodialysis; and when such patients require haemodialysis every second or third day the deficiency may significantly affect their practical management. Alwall *et al.* (1948-9) attempted to resolve the problem by the employment of arterial and venous glass cannulae connected by a fistula of capillary glass tubing. Despite continuous heparinization, however, the technique proved unsuccessful. The indwelling cannulae and fistula recently introduced by Scribner (Quinton *et al.*, 1960; Scribner *et al.*, 1960) for the treatment of patients with chronic renal failure by repeated haemodialysis are made of polytetra fluoroethylene (P.T.F.E.), which is inert in body tissues and does not provoke clotting: patency is maintained without anticoagulants.

THE APPARATUS*

The cannulae and fistula are prepared from thin-walled P.T.F.E. tubing† of 2.84 mm. external and 2.76 mm. internal diameter. The cannulae are heat-drawn to sizes suitable for cannulation of a radial artery and a superficial radial vein: they are bent to a J-shape with the short tapered arm 5 cm. and the long untapered arm 10 cm. in length. "Swagelok" fittings are used to connect the ends of the long arms of the cannulae separately to two short stainless-steel conduits which are part of a stainless-steel plate: the opposite ends of the stainless-steel conduits are similarly connected to a single short (6 cm.) U-shaped P.T.F.E. tube previously heat-drawn so that its middle third is constricted down to approximately 1.25 mm. internal diameter (Fig. 1). With the cannulae in a radial artery and superficial radial vein, blood can pass freely from artery to vein through the fistula. When haemodialysis is carried out it is necessary only to disconnect the fistula temporarily and to connect the long arms of the arterial and venous cannulae to standard polyvinyl chloride (P.V.C.) tubing

*All items of equipment, including apparatus for moulding P.T.F.E., may be obtained from the Medical Division, Tally Register Corporation, 1310 Mercer Street, Seattle 9, Washington.
†"CLIX" TWT/10, supplied by Siemens Edison Swan Ltd., 132 Long Acre, London W.C.2.