as well as in the adrenal cortex. Crooke's change was present in the basophil cells of the anterior pituitary of the first case, but not in Case 2. Certainly there was time in Case 1 for the full picture of Cushing's syndrome to develop, and it is interesting that this hyalinization of the basophil cells was so marked. In Case 2 it would appear that there was not time for this to develop. As Allott and Skelton (1960) point out, this finding has been an inconstant feature of previously reported cases; it probably takes some time to develop, and is therefore absent from those cases in which the disease is rapid in onset and progression.

At least four theories may be advanced to account for this observed association of carcinoma with adrenal overactivity. (1) It may be due entirely to chance. Against this is the observation that these cases fall into a distinct pattern, with variations from the usually observed incidence in females of the carcinomata concerned, together with an unusual age distribution. (2) The carcinoma is stimulated by adrenal overactivity. There is little evidence that this should be so. Though the experimental evidence of Pomeroy (1954) in mice supports this theory, neither Kaliss *et al.* (1954) nor Fisher and Fisher (1959) were able to find experimental support for this in rats. (3) There may be stimulation by the cancer of the adrenal glands or adrenal cortical adenomata to hyperactivity. It is tempting to suggest that this may have happened in Case 2, where metastatic carcinomatous growth occurred in an adrenal cortical adenoma. Such adenomata are not uncommonly seen at necropsy in patients not suspected clinically of having Cushing's syndrome, and are presumably often, therefore, non-active. (4) Representatives of any or all of the above explanations may be found in any series of patients showing this association. Though increased aldosterone production has been postulated as a possible explanation of the findings of this association (Spaulding *et al.*, 1955), only one case has so far been reported where this has been found to be so (Webster *et al.*, 1959). Conceivably Case 1 could be explained on this basis on the somewhat tenuous evidence of the low Na/K ratio, but this is doubtful, to say the least of it.

We are no nearer to explaining the mystery of why hypokalaemic Cushing's syndrome should occur in association with malignant disease involving such widely separated organs as the thymus (Duguid and Kennedy, 1930), colon (Warren, 1945), bronchus (Brown, 1928), and prostate (Webster *et al.*, 1959). We feel, however, that it is important to go on reporting these cases as they occur, since by comparing their similarities and differences a common factor may emerge to explain this syndrome.

Summary

Two cases in which carcinoma of the bronchus was associated with Cushing's syndrome are reported. They are compared with similar cases in the literature where this association between malignant disease and adrenal overactivity has been observed. These two cases are of interest because one presented with a florid psychosis and purpura, which proved to be due to bone-marrow metastases, and the other with an asymmetrical neuro-miopathy. Various theories are advanced in an endeavour to explain this association.

We thank Dr. M. G. Ashby and Dr. E. Idris Jones for permission to publish information about their

patients; also Dr. A. G. Stansfeld, of St. Bartholomew's Hospital, for the histological report and photographs of Case 2, and Dr. Jones-Williams, of Mount Vernon Hospital, for the α - and β -fraction estimations on Case 1. Thanks are also due to Dr. J. D. O'D. Lavertine for the post-mortem report on Case 1, and Mr. Macdonald for the photographs of this case, taken at Harefield Hospital.

REFERENCES

- Albright, F. (1938). *N. Engl. J. Med.*, **218**, 813.
 Allott, E. N., and Skelton, M. O. (1960). *Lancet*, **2**, 278.
 Bagshawe, K. D. (1960a). *Ibid.*, **2**, 284.
 — (1960b). *Ibid.*, **2**, 287.
 Brain, W. R., and Henson, R. A. (1958). *Ibid.*, **2**, 971.
 Brown, W. H. (1928). *Ibid.*, **2**, 1022.
 Cope, C. L., and Black, E. G. (1959). *Brit. med. J.*, **2**, 1117.
 Duguid, J. B., and Kennedy, A. M. (1930). *J. Path. Bact.*, **33**, 93.
 Edmunds, A. W. B., McKeown, K. C., and Coleman, P. N. (1958). *J. clin. Path.*, **11**, 237.
 Farrant, P. C., and Insley, J. (1960). *Lancet*, **2**, 434.
 Fisher, T. N., and Fisher, E., jun. (1959). *Proc. Soc. exp. Biol. (N.Y.)*, **100**, 780.
 Kaliss, N., Borges, P. R. F., and Day, E. D. (1954). *Cancer Res.*, **14**, 210.
 Parrish, J. A. (1960). *Lancet*, **2**, 603.
 Pomeroy, T. C. (1954). *Cancer Res.*, **14**, 201.
 Spaulding, W. B., Oille, W. A., and Gornall, A. G. (1955). *Ann. intern. Med.*, **42**, 444.
 Warren, F. L. (1945). *Cancer Res.*, **5**, 49.
 Webster, G. D., jun., Touchstone, J. C., and Suzuki, M. (1959). *J. clin. Endocr.*, **19**, 967.

ACUTE CARBON TETRACHLORIDE POISONING

TRANSAMINASE AND BIOPSY STUDIES

BY

J. K. DAWBORN, M.B., B.S., M.R.A.C.P.
Drug Houses of Australia Research Fellow

MARY RALSTON, M.B., B.S., D.C.P., M.C.P.A.
Pathologist to the Clinical Research Unit

AND

S. WEIDEN, M.Sc., Ph.D.
Biochemist to the Clinical Research Unit

*From the Walter and Eliza Hall Institute of Medical
 Research and the Royal Melbourne Hospital*

[WITH SPECIAL PLATE]

Many features of acute carbon tetrachloride poisoning are well established (Cameron and Karunaratne, 1936; Council on Industrial Health, 1946; Hardin, 1954). However, the following case of carbon tetrachloride poisoning provided a unique opportunity for studying the course of hepatic damage by means of serial estimations of the serum glutamic oxalacetic transaminase (S.G.O.T.) level (method of Reitman and Frankel, 1957) and serial liver biopsies.

Case Report

A storeman aged 22 years accidentally swallowed a mouthful of carbon tetrachloride. Immediately his head began to throb, he felt faint, and two hours later he vomited and passed dark urine. The following day he suffered increasing anorexia and abdominal pain. On the third day he was admitted to hospital, and clinical examination disclosed pronounced tenderness and rigidity in the epigastrium. The urine contained granular casts, and there was moderate proteinuria but no bilirubinuria. The initial biochemical analyses were as follows: serum bilirubin 1.4 mg. per 100 ml.; serum albumin 2.7 g. and globulin 2.3 g.

per 100 ml.; S.G.O.T. 2,120 units; serum alkaline phosphatase 13 King-Armstrong units; cephalin flocculation+++; prothrombin index 19% of normal; serum vitamin B₁₂ 1,000 µg. per ml.; serum amylase 266 units; urinary diastase 80 units; and blood urea 27 mg. per 100 ml.

Treatment consisted in a full diet, with administration of choline hydrochloride, 3.0 g. daily, antihistamine ("benadryl"), 200 mg. daily, nicotinic acid, 400 mg. daily, and vitamin K. On the fifth day he felt well and was no longer tender in the epigastrium; the S.G.O.T. level had fallen from 2,120 to 520 units and continued to fall (see Graph). On

Graph showing transaminase (S.G.O.T.) levels in moderately severe carbon tetrachloride poisoning during phase of recovery. There was a striking fall in level corresponding with clinical improvement. First estimation on third day probably did not represent highest level reached.

the ninth day "bromsulphalein" retention was 38%, serum amylase less than 100 units, and urinary diastase 20 units. All other biochemical tests gave normal findings. The prothrombin index was 100%.

Liver biopsy on the ninth day revealed centrilobular zonal necrosis of one-third to one-half of each liver lobule (Special Plate, Fig. 1). In these areas there was an inflammatory infiltrate of lymphocytes, plasma cells, and macrophages containing bile pigment. Scattered surviving liver cells were present near the central vein. The reticulin framework was preserved (Special

Plate, Fig. 2). Renal biopsy (tenth day) showed only pigment in the epithelium of occasional medullary tubules. Gastric biopsy (eleventh day) was normal.

By the twenty-eighth day liver biopsy revealed considerable improvement. There were still centrilobular damage and fatty change, but liver-cell regeneration was present (Special Plate, Fig. 3). In a third liver biopsy, obtained on the seventieth day, there was almost complete restoration to normal; only a few inflammatory cells remained in the sinusoids and some fatty change persisted.

Discussion

Carbon tetrachloride causes swelling of mitochondria in liver cells and increased permeability of the mitochondrial membrane. There are disorganization of enzyme activity and loss of co-factors and enzymes from the mitochondria (Christie and Judah, 1954; Gallagher *et al.*, 1956). Choline and antihistamines, which lessen this damage to mitochondria (Christie and Judah, 1954; Judah and Rees, 1959; Judah, 1960), appeared to be effective in our case.

Raised levels of S.G.O.T. in carbon tetrachloride poisoning have been reported in human beings and in animals (Wróblewski *et al.*, 1956; Block and Cornish, 1958). In animal experiments S.G.O.T. was the most sensitive of the three enzymes measured, and the rise in S.G.O.T. was proportional to the extent of liver damage. In a study at the Royal Melbourne Hospital of 48 patients with various forms of liver disease (O'Brien *et al.*, 1960) there was an approximate correlation between S.G.O.T. levels and the extent of liver-cell necrosis. In the present case an S.G.O.T. of at least 2,120 units was associated with necrosis of one-third to one-half of every liver lobule in the biopsy. The prompt fall in S.G.O.T. level was associated with liver-cell regeneration after cessation of injury. The administration of choline and antihistamines may have contributed

to this fall by lessening release of enzymes from the liver cells.

The liver histology was consistent with previously described changes (Himsworth, 1950). The reticulin framework remained intact despite extensive centrilobular necrosis, so that full recovery could be anticipated (Special Plate, Fig. 2). Damage to the kidney was slight, as is usually the case when poisoning follows ingestion of carbon tetrachloride.

The present case illustrates the value of serial S.G.O.T. estimations in assessing the extent and duration of liver damage and the rate of recovery in carbon tetrachloride poisoning. There was a rapid and extensive cell necrosis which was demonstrated by liver biopsy to involve approximately half the liver lobule.

Summary

A non-fatal case of carbon tetrachloride poisoning is reported. S.G.O.T. levels and liver histology were followed at intervals for up to 70 days. The former returned to near normal after nine days, while the latter was still abnormal at 28 days. Clinical recovery was complete by 10 days and histological recovery by 70 days. The results are discussed.

We wish to thank Professor G. S. Christie for his advice in the clinical management of this case, and Dr. I. J. Wood for permission to report it. The illustrations were prepared by Mr. E. Matthaei, of the University of Melbourne.

REFERENCES

- Block, W. D., and Cornish, H. H. (1958). *Proc. Soc. exp. Biol. (N.Y.)*, **97**, 178.
- Cameron, G. R., and Karunaratne, W. A. E. (1936). *J. Path. Bact.*, **42**, 1.
- Christie, G. S., and Judah, J. D. (1954). *Proc. roy. Soc. B.*, **142**, 241.
- Council on Industrial Health (1946). *J. Amer. med. Ass.*, **132**, 786.
- Gallagher, C. H., Gupta, D. N., Judah, J. D., and Rees, K. R. (1956). *J. Path. Bact.*, **72**, 193.
- Hardin, B. L., jun. (1954). *Industr. Med. Surg.*, **23**, 93.
- Himsworth, H. P. (1950). *Lectures on the Liver and its Diseases*, 2nd ed. Blackwell, Oxford.
- Judah, J. D. (1960). *Nature (Lond.)*, **185**, 390.
- and Rees, K. R. (1959). *Biochemical Society Symposia*, No. 16, p. 94.
- O'Brien, E. N., Paterson, B., Ralston, M., and Mackay, I. R. (1960). *Aust. Ann. Med.*, **9**, 295.
- Reitman, S., and Frankel, S. (1957). *Amer. J. clin. Path.*, **28**, 56.
- Wróblewski, F., Jervis, G., and LaDue, J. S. (1956). *Ann. intern. Med.*, **45**, 782.

The formation of a medical faculty of Auckland University as a further step in the establishment of a medical school has been announced by the university council. The following were constituted the faculty: the professor and other full-time members of the Postgraduate School of Obstetrics and Gynaecology; Professor H. M. CAREY, Associate-professor G. H. GREEN, and one other to be appointed; the sub-dean of the branch medical faculty of the University of Otago, Dr. W. E. HENLY; three full-time tutors in medicine, surgery, and paediatrics at Auckland Hospital, Dr. H. K. IBBERTSON, Dr. J. C. GILLMAN, and Dr. R. H. CAUGHEY; the physician-in-charge of the medical unit at Auckland Hospital, Dr. J. D. K. NORTH; three members of the postgraduate committee of medical education in Auckland, to be appointed; three professors from the departments of chemistry, physics, and zoology, to be appointed; and the dean of the science faculty, Professor F. H. SEGAR.

J. R. BILLINGHURST *ET AL.*: CARCINOMA AND CUSHING'S SYNDROME

FIG. 1.—Case 2. Secondary carcinoma in adrenal gland. (H. and E. $\times 45$.)

FIG. 2.—Primary bronchial carcinoma. (H. and E. $\times 45$.)

J. K. DAWBORN *ET AL.*: ACUTE CARBON TETRACHLORIDE POISONING

FIG. 1.—Liver biopsy obtained on ninth day after swallowing carbon tetrachloride. There is pronounced centrilobular zonal necrosis. (Periodic acid-Schiff. $\times 80$.)

FIG. 2.—Section of liver biopsy obtained on ninth day, showing preservation of reticulin framework despite cellular necrosis. (Silver impregnation stain. $\times 80$.)

FIG. 3.—Section of liver biopsy obtained on 28th day. There are large regenerating liver cells with hyperchromatic nuclei and considerable inflammatory-cell infiltrate. Final biopsy on 70th day showed restoration of normal liver histology. (H. and E. $\times 400$.)

C. N. BEST AND P. B. COOK: MESENTERIC RETICULOSARCOMA AND STEATORRHOEA

FIG. 1.—Mass of largely necrotic reticulum cells. (H. and E. $\times 700$.)

FIG. 2.—Reticulin preparation, showing dense network. ($\times 130$.)