BRITISH MEDICAL JOURNAL

LONDON SATURDAY SEPTEMBER 16 1961

PROTECTION OF THE RESPIRATORY TRACT*

BY

Sir VICTOR NEGUS, D.Sc., M.S., F.R.C.S.

[WITH SPECIAL PLATE]

The memory of Walter Jobson Horne will be kept alive by the institution of this lecture. I am proud to be the first to deliver it, particularly as I remember a number of meetings at which his somewhat controversial observations gave life to the discussions. He was in appearance rotund and reminiscent of a character from Dickens; in intellect he was erudite and a good speaker. He was, in brief, a character.

The respiratory tract, and particularly inflammatory lesions of the larynx, were of great interest to him; his inclination was toward pathology and away from operations, characteristics to be remembered and imitated. A keen supporter of the B.M.A. and President of a Section at the Meeting in Bradford, he also had the honour of giving the Semon Lecture in 1921.

I have chosen as my subject: "Protection of the Respiratory Tract."

Consideration of this problem allows no distinction to be drawn between the upper and lower respiratory passages, as frequently occurs in the education of medical students; the former is often classed as a special subject and the latter as part of general medicine, with confusing results.

Since infections of the respiratory tract are among the most common of complaints, with the loss of a vast number of working days, it is important to have some understanding of the complicated physiological processes entailed.

After a medical career devoted to the study of this subject I regret to find myself in some confusion, but I will attempt to outline certain aspects which I hope will serve as a basis for therapeutic principles.

The respiratory tract must be considered from the nostrils to the alveoli, and in addition the peculiar anatomy of man makes it essential to include the mouth and pharynx, since mouth-breathing is frequently used, especially during speech. It will be best to consider the various dangers against which protection must be afforded, and foremost among these is the avoidance of entry of water and food into the trachea and bronchi; this is a primitive function which concerns lung-breathing fish, amphibians, reptiles, birds, and mammals.

When the pulmonary apparatus was evolved it was essential to provide a protective valve at its origin from the pharynx; this valve is the larynx, often considered, incorrectly, to function solely as the organ of voice.

*The Walter Jobson Horne Memorial Lecture delivered at the Adjourned Annual General Meeting of the British Medical Association, Sheffield, July 20, 1961.

Phonation is, however, facilitated in animals with a larynx, this being a convenient organ for the purpose, without anatomical adaptation. The simple larynx of an alligator, for example, lying flat in the floor of the pharynx, remains closed when water or food is being swallowed; reflex stimulation removes the danger of inundation of the trachea or lungs through an open glottis, and the same applies to the uncomplicated larynx of an amphibian such as the newt.

Crossing of the Air and Food Passages

Animals with a pulmonary apparatus inspire through the nose. This route is not designed for the benefit of respiration, but has as its object the drawing of air over the olfactory mucosa in order to maintain the acuity of the sense of smell. Since the nasal passages open in the roof of the mouth or pharynx, while the larynx opens from the floor of the pharynx, there is, of necessity, crossing of the two tracts for air and for food. In lung-fish, amphibians, reptiles, and birds it is essential that respiration shall cease while food and water are passing down the gullet; but in certain mammals the desirability of swallowing and breathing simultaneously has been overcome when the food is semi-liquid. This occurs with herbage, which must be taken in over lengthy periods of time (Text Fig. 1). Crossing of the passages for air and food is a particular cause of trouble in man.

Requirements of Olfaction

An important anatomical structure to be considered now is the epiglottis, which rises at the anterior end of the larynx; it is developed in animals of keen powers of scent to subserve the olfactory sense by its approximation to the soft palate, thereby shutting off the mouth from the air-stream and thus ensuring passage of all of the inspired air over the olfactory mucosa of the nose (Text Fig. 1).

The presence of the epiglottis does not of itself solve the problem of simultaneous breathing and swallowing; but a further modification connected with the epiglottis is of help. This modification is the lateral food-channel.

Lateral Food Channel

This channel is bounded medially by the aryteno-epiglottic fold and is shut off above by prolongation of the soft palate; this divides to form a shelf around the upstanding larynx, which projects into the posterior nasal passage; the food-passage is thereby separated

from the air-passages. In herbivorous animals such as deer and antelopes the channel is very efficient; the most efficient examples are found in cetaceans such as dolphins, porpoises, and some whales (Text Fig. 2).

Man, of herbivorous descent, has lateral food-passages of a sort along which liquids or semi-liquids may travel. These passages, though extremely efficient in true herbivora such as ungulates and many rodents, are in man not very effective in protecting the respiratory tract. It therefore happens that, though the infant can swallow milk along the lateral channels without complete interruption of respiration, the adult runs a risk if he attempts to swallow and breathe at the same time; it is common experience to find that fluid has gone the wrong way, and in individuals with paralysis of the larynx this experience is dangerous and sometimes fatal.

Entrance of Solid Objects

The inefficiency of the lateral passages is a danger to man as regards entrance of liquids, but a further hazard exists in his faculty of breathing through the mouth. The close apposition of epiglottis and soft palate of keen-scented animals is no longer required in higher apes and man, because of regression of the sense of smell in favour of sight. Evolutionary flexion of the head on the vertebral column with, in man, shortening of the snout leads to separation of the epiglottis from the palate, allowing air to pass freely through the mouth when it is open. This is, of course, to the advantage of speech, though it reduces the safety of the respiratory tract (Text Fig. 3). If some object is held between the lips or teeth, or is in the buccal cavity, a sudden inspiration may draw it through the larynx into the trachea or one of the bronchi; this applies especially to such light objects as pieces of nut or fragments of vegetables, substances which by irritative effects on the bronchial mucous membrane are very dangerous and often fatal. Sudden laughter or a cry may precipitate this accident.

TEXT FIG. 1.—Diagram to illustrate modifications to subserve olfaction. In most gill-breathing fish there is no communication between the olfactory organ and the food-tract; the early human embryo is similar. In air-breathing lung-fish and in amphibians, reptiles, most birds, and all mammals the nasal passage opens into the mouth or pharynx. Coaptation of nasal and pulmonary air-passages is illustrated in the crocodile, cat, and antelope, the epiglottis of mammals assisting in this process. Man is seen to have a gap between nasal and pulmonary air-tracts. Cetaceans, such as some whales, have a permanently intranasal larynx, of great benefit for simultaneous breathing and swallowing.

Entrance of Bacteria

The larynx is incapable of preventing entrance of bacteria coming from the external air or from the nasal passages, the sinuses, or the pharynx; each respiration may carry in and out innumerable organisms, a strong reason for the wearing of a mask by those with respiratory infections. The nose itself, when infected, becomes a danger instead of a protection to the respiratory passages. It is most important, therefore, to deal effectively with inflammatory conditions of the nose, and especially with collections of pus in the paranasal cavities; the maxillary sinuses are usually the key to the situation, and puncture and lavage at an appropriate time may be indicated as a means of avoiding chronic infections or spread to other sinuses.

Stagnation of secretions in the posterior nasal passage, usually known as the nasopharynx, is frequently due to the obstructive presence of adenoids and can be of considerable danger; surgical clearance may be necessary, but with special care to avoid damaging submucosal tissues, such as the pharyngeal aponeurosis and muscle fibres, for fear of subsequent scarring and elimination of ciliary streams, leading to droplet infection of the lungs. Normally ciliary streams propel a mucoid blanket which is then carried by swallowing movements, starting in the posterior nasal passage, into the stomach, where organisms are killed by gastric juices.

Removal of adenoids is a difficult operation if it is to succeed in clearing obstruction without inflicting damage on the submucosa. Removal of tonsils, if it

TEXT FIG. 2.—Diagram to illustrate the lateral food-channel. In each case the pharynx and larynx are seen in transverse section above and laterally below. Ph=pharynx; Tr=trachea; Ep=epiglottis; V=laryngeal ventricle; LFC=lateral food-channel; NP=nasal passage; Arc=arcus palato-pharyngeus. The larynx of the lung-fish lies flat in the floor of the pharynx with no lateral protection; the arrangement in the lizard is similar, the glottis being at some distance from the nasal passage. In the lion there is an epiglottis for olfactory purposes, but no protection of the pulmonary air-passages except when the glottis is closed. The monkey has aryteno-epiglottic folds with a lateral food-channel of no great efficiency. Herbivores, such as deer and oxen, have an upstanding larynx and an extremely efficient food-channel, guarded above by a complete arcus; cetaceans, such as dolphins, have complete separation of air and food passages.

leads to scarring and contraction of the fauces, may itself prevent efficient removal of infected secretions by interfering with swallowing movements of the superior constrictor muscles.

Inhalation of Dust

A certain amount of filtering is attained by vibrissae at the nostrils, and still more dust is entangled in the mucoid blanket which covers the nasal mucosa, to be removed by ciliary action; the larynx is not capable of preventing entrance of small particles unless it is closed. Irritation of the mucosa may promote cough, which may momentarily expel some of the debris. Subsequently ejection of dust and bacteria depends on ciliary action in the trachea and bronchi, to be considered later.

Harmful Gases and Vapours

Since respiration can be interrupted only for a very brief period, it is obvious that poisonous gases will reach the finer bronchi in spite of cough. This is noticeable in the slight or absent damage to the larynx after exposure, while inflammation and ulceration may be produced in the bronchi. In warfare reliance has, therefore, to be placed on gas-masks.

Coal-gas may give warning by its smell, and those with anosmia are consequently in danger of asphyxiation from undetected leakages.

Winter fogs have in recent years been accused of causing widespread respiratory infections, and inhalation of sulphuric or other irritating substances cannot be excluded except momentarily, since respiration cannot be discontinued; here again, as with the gases of warfare, masks may be of use. Contaminated secretions in the bronchi rely on ciliary action for their ejection.

Protection by the Nose

Reference has been made to the warning given by the sense of smell—a warning which can do no more than cause sudden and temporary spasmodic closure of the larynx or may lead to the wearing of a gas-mask.

Certain animals—for example, newts and frogs, crocodiles, water buffaloes, and whales—can prevent entrance of water by closure of the nostrils; in man this is not possible.

Functions of the Nasal Organ

At this stage it will be best to discuss the relationship of the nose to the lower respiratory passages; it is sometimes assumed that the main function of the nose is to help in maintaining the health of the tracheo-bronchial tree by arresting bacteria and dust and by warming and moistening inspired air. But when the nasal organ is examined in its primitive stages of evolution it is found to have a purely olfactory function; this is obvious in a fish such as the pike, where water entering during forward progression reaches the olfactory rosette. Various methods for drawing water over the rosette are found, some depending on the mechanism which helps to cause circulation of water through the gills; but in no instance is the nasal organ of any use in protection of the respiratory apparatus.

Lung-fish, which breathe air when the water in which they live dries up, have a communication between the nasal passage and the pharynx, but here again for the sole benefit of the olfactory sense; there is obviously no necessity for warming or moistening air.

When mammals are examined the nose is found to be divided into two sections: one anteriorly for the warming and moistening of air, and the other posteriorly for olfactory stimulation. If all mammals are examined it is found that the two sections are complementary and parallel in complexity; keen-scented animals, and especially carnivores, have a wide olfactory area and an extensive air-conditioning system, while species with feeble powers of smell, such as the higher apes and man, have a very restricted olfactory area, with corresponding reduction for warming and moistening air.

Though the lower respiratory tract benefits from passage of air through the nose, yet the nose was not designed for this purpose and does not adapt itself efficiently to alterations of climate; its primitive function is olfactory.

Air-conditioning

Respiratory exchanges of oxygen and carbon dioxide cannot continue in the alveoli unless the surface of the epithelium is moistened. The supply of moisture comes partly from the atmosphere and partly from transudation and secretion through the epithelium lining the nose, trachea, and bronchi. Air reaching the alveoli is at body temperature and is saturated with moisture; heat is essential if the air is to carry much water vapour.

TEXT FIG. 3.—Relation of the palate and the larynx. In keen-scented animals such as the Indian barking deer or muntjac the epiglottis lies above the soft palate and ensures that all air shall enter and leave through the nose. Carnivora, of which the tiger is an example, have a similar arrangement, of great value when tracking prey. The gorilla relies on sight and not on scent, and no longer needs to direct all inspired air through the nose; there is a gap between the soft palate and the epiglottis. This is still more marked in man.

In animals such as dogs, cats, and rabbits a very elaborate maxillo-turbinal body with an enormous surface, capable of alteration in area, can bring air to body temperature and to complete saturation for olfactory purposes and with incidental benefit to the respiratory tract. Man, with his degenerate sense of smell and restricted air-conditioning area, is not similarly equipped, and consequently much moisture, in addition to that from the nose, is given up in the passage of air through the trachea and bronchi (Special Plate, Fig. 1).

Knowing the importance of this function and realizing the deficiencies of the human nose, great care must be taken to preserve the nasal mucosa, limited as it is; removal of tissue from the maxillo-turbinals is highly improper. If the nose is dry from thyroid insufficiency, or rhinitis sicca or atrophica, the taking of small doses of potassium iodide is of benefit; a simple spray of sodium bicarbonate 0.5% and glycerin 10%, with chlorotone 0.5% as a preservative and normal saline as a base, may be of help in encouraging production of secretion.

Air-conditioning of rooms is important, and many cases of recurrent sinusitis are caused by over-dry air; a relative humidity of 50% or even more should be attained, but is not always attained in winter.

Atmospheric air in a cold winter carries little moisture, and when warmed up in a room or hospital ward the relative humidity—which varies inversely with the temperature, falls to an unhealthy level, perhaps as low as 20%. Alternately, moist air in the summer with a high water content, when brought into a room and artificially cooled, causes the relative humidity to rise to an unpleasant degree.

Supply of Moisture by the Nose

Nothing must be done to reduce the capabilities of the nasal mucosa for air-conditioning, but, on the other hand, steps may be necessary to improve matters. Of these, straightening of the nose or the septum is sometimes required, but only after estimating the relative air-flow through each nasal fossa; other forms of obstruction, as by polypi, may need attention.

Response of the Nose to Hot and Cold Climates

If the nose were directly concerned with the protection of the lower respiratory passages and lungs there would be dilatation of the arterioles and capillaries when inspired air is cold; such is not the case, since inhalation of very cold air at a temperature as low as -40°C . leads to slight vascular constriction and shrinkage of the mucosa, as shown by lowered resistance in the nose. This is similar to constriction of the skin capillaries designed to prevent loss of heat from the body. Very cold air, being dry, should theoretically lead to filling of sinusoids to provide additional moisture by transudation, but such is not the case (Text Fig. 4 A, B).

The nose retains much of the warm moisture coming from the bronchi on expiration, with advantage to the conditioning of subsequent inspirations; because of this, nasal respiration is superior to mouth-breathing, which has not the same faculty of retaining heat or moisture on expiration.

Nocturnal mouth-breathing in those whose nose is clear by day can often be prevented by the use of an

oral shield placed between the lips and teeth. Bringing an obstructed nostril uppermost by turning over in bed may clear the nose by virtue of the naso-pulmonary reflex. In man there is a rising temperature gradient from the nose to the trachea and bronchi; cold, dry air reaches body temperature and saturation only when it is in the alveoli; there is a corresponding fall on expiration.

TEXT FIG. 4.—Kymographic tracings to illustrate phases of respiration (above), temperature of air entering and leaving the nose (middle), and resistance in the nose as registered by the nasopharyngeal pressure (below). A. The subject breathes air reaching the nose at about -20°C .; the temperature rises during each expiration and falls during inspiration, but is still above zero. The pressure in the nasopharynx is $\pm 26\text{ mm. H}_2\text{O}$, which is only slightly above normal. B. Hot air causes increased congestion of the nasal mucous membrane, as shown by raised pressure in the nasopharynx to a level of $\pm 50\text{ mm. H}_2\text{O}$. The air enters and leaves the nose at a level equal to body temperature.

LEGENDS TO SPECIAL PLATE

FIG. 1.—Relative size of maxillo-turbinals in man (left) and cat (right). In man the ethmo- and maxillo-turbinals are flat plates lying close to the nasal septum and with hollow unwanted spaces laterally, the maxillary sinuses; the areas for air-conditioning and for olfaction are restricted. In the cat and other keen-scented carnivores the maxillo-turbinal is widely branching, occupying the whole breadth of the snout and affording a large area for air-conditioning; the olfactory area is correspondingly extensive.

FIG. 2.—Transverse sections of cilia photographed by electron microscopy and reproduced by kind permission of Professor Fawcett. ($\times 40,000$.) Each cilium has stiffening fibrils in a protoplasmic core, nine pairs around the periphery and one pair in the centre. There is remarkable similarity between the cilia of a mollusc (A), an amphibian (B), and a mammal (C). The centre pairs of fibrils are all orientated on one axis, which is no doubt designed to ensure uniformity of bending at a right-angle to this axis, thus avoiding interference.

FIG. 3.—High-power view of part of the lamella in the olfactory recess of a carp (*Cyprinus carpio*), illustrating the large number of goblet cells secreting mucus; some cells have reached the surface and have discharged their contents. When the fish is placed in distilled water or in saline solution similar to salt water there is increased activity of the goblet cells, no doubt intended to counteract inwardly directed osmosis or outwardly directed transudation.

SIR VICTOR NEGUS: PROTECTION OF THE RESPIRATORY TRACT

FIG. 1

FIG. 2A

FIG. 2B

FIG. 2C

FIG. 3

Retention of heat and moisture on expiration should be remembered with the subjects of tracheostomy or laryngectomy, who should wear a shield over the tracheostome, covered with at least six layers of gauze as a partial replacement for the nose.

Hot air is reduced in temperature as it passes through the nose on inspiration; and on exertion expiration by the mouth is of advantage as a means of losing heat.

In a very cold climate mouth-breathing is highly undesirable. Very slow inspirations allow chilling of the lower air-passages, since the preliminary warming influence of expired air is much reduced (Text Fig. 5).

TEXT. FIG. 5.—Slow breathing at six respirations per minute removes most of the warming effect of expired air; the current of air reaching the nose at -40°C . falls to -35°C . at the end of inspiration. There is then a sudden rise to $+10^{\circ}\text{C}$. at the end of expiration.

Ciliary Action

The health of the respiratory passages depends in great part on the action of ciliary streams, which are capable of expelling bacteria and small particles of dust if these happen to be entangled in the blanket of diluted mucus which overlies the epithelium. Cilia are of primitive origin and their structure is of great interest; the cilia of molluscs, amphibians, and mammals are almost indistinguishable. All have stiffening fibrils, nine pairs round the periphery and one pair in the centre; the latter are orientated on the same axis, so that waving of the cilia shall be in one direction only, with no clashing (Special Plate Fig. 2).

Cilia line the nasal fossa and sinuses, the posterior nasal passages, the trachea, bronchi, and bronchioles; there are none in the atria, air-sacs, or alveoli. Since the bronchial tree of man divides, according to Hilding, into about 50,000 branches, the task of cilia is great in cleansing these minute air-passages. Plugging of smaller bronchi or bronchioles by thick mucus may cause pulmonary collapse, requiring aspiration of the obstructing secretion.

Nothing must be done to upset ciliary action. Drastic operations on the intranasal structures, the sinuses, or posterior nasal passage may cause scarring and obliteration of ciliary streams; harm may be caused by excessive use of cauterization or ionization, or by the use of drugs with a paralysing effect, such as strong cocaine. Drying of the mucosa is fatal to ciliary action, whether by powerful vasoconstrictor drops or sprays, by prolonged

inhalation of constricting vapours, by excessive doses of atropine, or by drying of the air; in this respect humidification of inspired air is important, and those with tubes in the trachea should have the surrounding air moistened to a relative humidity of at least 50% until their mucosa has become adapted to changed conditions. Even then a shield should be worn over the tracheostome.

Potassium iodide in small doses is useful in encouraging liquefaction of viscid secretions; wetting agents are sometimes required in humidified air, as in acute tracheo-bronchitis. Since the larynx cannot exclude bacteria, reliance is placed on ciliary streams and on lysozymes in the mucoid blanket. With an efficient mucoid blanket and regular removal of viruses and bacteria there should be no invasion of the epithelium; it is only when this is penetrated from lack of moisture or from stagnation by obstruction that the second line of leucocytic defence is called into action. Some who are prone to recurrent colds in the nose may be helped by vaccine treatment. In cases of acute rhinitis it is important to obviate complete obstruction of the nose by the use of steam inhalations, or, in certain instances, a spray containing 0.5% ephedrine. Entrance of foreign bodies into the bronchi leads eventually to the formation of granulations, with interruption of ciliary streams; early removal of the intruding object is essential. Infection in the form of bronchitis, bronchiectasis, or bronchial or lung abscess may necessitate removal of secretions by direct or indirect aspiration, with treatment, in certain cases, of obstructing granulations by some astringent such as protargol or silver nitrate. A growth, whether simple or malignant, will cause obstruction and infection, and so will persisting bronchospasm, as in chronic asthma.

Cough

The majority of individuals never cough and yet suffer no disability in the respiratory tract, owing to the protective action of cilia. When a foreign body tends to enter the larynx or actually passes into the trachea cough may be valuable for ejecting the intruder; but the initial deep inspiration can be a danger by drawing the object, especially if it is light, still deeper into the bronchial tree. When a foreign body is lodged in a bronchus cough may have little or no effect, because the sudden expiratory effort is accompanied by contraction of bronchial musculature, which firmly grips the object; instrumental removal is required.

Plugs of viscid secretion can often be removed by cough, but here again partial or complete muscular closure of bronchi or bronchioles tends to prevent effective clearance, particularly in asthma.

Exchange of Fluid in the Respiratory Tract

It is impossible in a short communication to discuss this subject in detail, since it is one of considerable complexity.

It is essential for the exchange of oxygen and carbon dioxide between the air in the alveoli and the blood in the pulmonary capillaries that the air should be saturated with moisture at body temperature. Water vapour is derived in varying amounts from the external atmosphere according to the degree of relative humidity; part comes from transudate and some from serous and mucoid glands and goblet cells, but there are none of these in the atria, air-sacs, or alveoli, and in consequence

the moisture for these terminal cavities enters by diffusion of air. The amount from each source depends on hydrostatic pressure, on osmosis, and on the activity of secreting cells.

To the water covering the epithelium mucin is added in varying amount, and attention must be directed to this substance. Its presence is of help to ciliary action by giving viscosity to the mucoid blanket, as Proetz has described it; but, though cohesion assists the drag of cilia from a distance, mucin is not essential for ciliary action. Cilia, in fact, work more vigorously in a watery medium, as seen in animalcules such as *Paramecium*. But mucin, a polysaccharide insoluble in water, provides a waterproofing covering to the epithelial surface. This action may be observed in eels, whose thin and flexible skin is covered with viscid mucus. If this slimy covering is removed the fish will, if replaced in fresh or distilled water, take in so much fluid by osmosis as to swell up and die.

The lamellae in the olfactory recess of a fish such as the carp are provided with very many goblet cells, no doubt designed to waterproof the permeable olfactory epithelium (Special Plate Fig. 3).

In the respiratory tract the purpose of a mucoid blanket appears to be similar—to control outward transudation and inward osmosis through the thin and permeable epithelium, while at the same time allowing ciliary action to continue and also permitting passage of olfactory molecules.

In bronchitis excessive collection of fluid in the smaller bronchioles, probably as a result of irritating substances, interferes with respiratory exchanges; in association with this tendency to flooding there is considerable increase of goblet cells, apparently as a well-intentioned effort at protection. But increase of goblet cells entails reduction of ciliated cells, with the result that viscid secretion fills the lumen of the bronchioles and is not removed by ciliary streams.

In asthma a similar process occurs, no doubt dependent in part on increased permeability of cell membranes in an allergic state, with a coincidental rise of pH.

Permeability of cell membranes is influenced by the presence of calcium, a deficiency of which facilitates transudation and osmosis. Administration of calcium gluconate together with dilute hydrochloric acid may be of advantage, especially in cases of nasal allergy, in reducing oedema.

Conclusion

An attempt has been made to show how various evolutionary changes have affected the protection of the respiratory tract of man; these changes, though detrimental to the preservation of health, have been of great advantage in facilitating speech and thereby the development of man's intellect.

A plaque commemorating Dr. Marie Stopes has been unveiled at the entrance to Norbury Park Mansion, Dorking, once her home, on behalf of the Royal Society of Literature. The plaque says: "This house, built for William Lock in 1774, was for many years a favourite resort of Fanny Burney and was from 1934 to 1958 the home of Marie Stopes, scientist and author. By her it was bequeathed to the Royal Society of Literature. The society, of which she was a fellow from 1913 to 1958, has caused this plaque to be erected in her memory and in token of gratitude."

INTERFERENCE PATTERNS ENCOUNTERED WHEN USING ATTENUATED POLIOVIRUS VACCINES

BY

J. H. HALE, O.B.E., M.D., M.R.C.P.

L. H. LEE, M.B., B.S.

AND

P. S. GARDNER,* M.D., Dip.Bact.

Regional Public Health Laboratory, Newcastle upon Tyne, and the Department of Bacteriology, University of Malaya

An account has already been given (Hale *et al.*, 1959; Knowelden *et al.*, 1961) of a trial of Sabin live poliovirus vaccine type 2 in Singapore during the type 1 epidemic of 1958. It was shown that the number of patients who developed paralytic poliomyelitis less than eight days after vaccination was close to that expected from the rates in unvaccinated children, whereas the number of cases which occurred later was lower than the expected number. During the trial we took the opportunity of studying the interference effects between the vaccine strain and the epidemic strain and also between the vaccine strain and other existing enteroviruses.

Recently a further study of Sabin attenuated vaccine types 1, 2, and 3 took place under the auspices of the Medical Research Council. Children from Newcastle upon Tyne co-operated in this study, and we were presented with a further opportunity of observing interference effects.

SINGAPORE TRIAL

The schedules of vaccination and laboratory procedure have previously been described (Hale *et al.*, 1959).

Effect of Type 2 Vaccine Strain on Type 1 Epidemic Strain

Fig. 1 shows the days after vaccination, when clinical cases of poliomyelitis due to the type 1 virus occurred among children who had received the type 2 vaccine. Eight cases occurred within eight days of vaccination, and, as is shown later, these were in children already incubating type 1 infection when given the vaccine. There were two other cases in vaccinated children, but these cannot be included in the analysis, since vaccine was given to one who was actually paralysed and to the other after the onset of illness.

No cases occurred in children between 8 and 19 days after vaccination; it is during this period that multiplication of the vaccine strain in susceptible cells of the alimentary tract takes place and maximum interference effect would be expected. Paralytic cases that occurred later were probably instances of type 1 infection breaking through the heterologous type 2 protection acquired as a result of previous infection induced either naturally or by vaccine.

Effect of Type 1 Epidemic Strain on Type 2 Vaccine Strain

The eight cases of type 1 clinical disease that occurred within eight days of vaccination are presented in Table I.

*Now at the Department of Bacteriology, Royal Victoria Infirmary, Newcastle upon Tyne.