Connell, Miss M. Simpson, Mr. W. G. Lavers, and Mr. J. W. Michieli for help with the diagrams and tabulations, and Mr. G. Knight, of the Statistical Unit at the Wellcome Research Laboratories.

REFERENCES

- Enders, J. F., Katz, S. L., Milovanović, M. V., and Holloway, A. (1960). *New Engl. J. Med.*, **263**, 153.
 Hilliard, L. T., and Kirman, B. H. (1957). *Mental Deficiency*, pp. 122, 398. Churchill, London.
 Hoekenga, M. T., Schwarz, A. J. F., Carrizo Palma, H., and Boyer, P. A. (1960). *J. Amer. med. Ass.*, **173**, 868.
 Katz, S. L., Enders, J. F., and Holloway, A. (1960a). *New Engl. J. Med.*, **263**, 159.
 ———, Kempe, C. H., Black, F. L., Lepow, M. L., Krugman, S., Haggerty, R. J., and Enders, J. F. (1960b). *Ibid.*, **263**, 180.
 ———, Medearis, D. N., jun., and Enders, J. F. (1958). *Amer. J. Dis. Child.*, **96**, 430.
 Krugman, S., Giles, J. P., and Jacobs, A. M. (1960). *New Engl. J. Med.*, **263**, 174.
 Schwarz, A. J. F., Boyer, P. A., Zirbel, L. W., and York, C. J. (1960). *J. Amer. med. Ass.*, **173**, 861.
 Stokes, J., jun., Reilly, C. M., Hilleman, M. R., and Buynak, E. B. (1960). *New Engl. J. Med.*, **263**, 230.

MAGNESIUM METABOLISM IN PARATHYROID DISEASE

BY

SAMIR HANNA, Ph.D., M.B., D.Med.Sci.
United Arab Republic Government Scholar

K. A. K. NORTH, D.Phil., M.B., M.R.C.P.
Registrar in Medicine

I. MacINTYRE, Ph.D., M.B.
Lecturer in Chemical Pathology

AND

RUSSELL FRASER, M.D., F.R.C.P.
Professor of Clinical Endocrinology

From the Departments of Chemical Pathology and Medicine,
Postgraduate Medical School of London, Hammersmith

A number of reports suggest that the parathyroid glands may influence magnesium as well as calcium metabolism. Greenwald and Cross (1925) claimed that prolonged administration of parathyroid extract increased the faecal excretion of magnesium. Greenberg and Mackey (1932) observed an elevation of serum magnesium in dogs receiving injections of parathyroid extract. The first account in man is that of Bulger and Gausmann (1933). They described a case of hyperparathyroidism in which a negative pre-operative balance of both calcium and magnesium became positive after operation. Bassett and Van Alstine (1935) found a small change in magnesium balance after parathyroidectomy, and Tibbetts and Aub (1937) and Barnes, Krane, and Cope (1957) confirmed the findings of Bulger and Gausmann. Potts and Roberts (1958) have drawn attention to the dangers of hypomagnesaemia occurring after parathyroidectomy. We here present pre-operative and post-operative data obtained on the last seven cases of parathyroidectomy performed in the Hammersmith Hospital.

Material and Methods

Six cases of primary hyperparathyroidism and one case of pluriglandular syndrome are presented. The former were found at operation to have enlargement of one parathyroid gland, and in all but Case 5 the histological appearance of the gland was that of a chief-cell adenoma. In Case 5 the cells were mostly oxyphil. The case of pluriglandular syndrome (No. 3)

had, in contrast, enlargement of all parathyroid glands. Three of these were removed and part of the fourth was excised. Previously he had undergone subtotal adrenal-ectomy for Cushing's syndrome, and because of enlargement of the pituitary fossa he was on large doses of dexamethasone. He was also hypertensive and was receiving treatment with chlorothiazide and mecamlamine throughout. Biochemical details of the seven cases are given in Table I.

TABLE I.—Biochemical Data in the Seven Cases

Case	Sex and Age	Serum Ca (mEq/l.)	Serum Mg (mEq/l.)	Serum Inorganic P (mg./100 ml.)	Serum Alk. Phosphatase (K.-A. Units/100 ml.)	Blood Urea (mg./100 ml.)
1	F 61	7.0	1.5	2.3	74	75
2	F 58	7.8	1.9	2.2	135	55
3	M 33	6.5	1.6	2.1	11	74
4	F 72	9.0	1.7	3.8	21	72
5	F 32	9.0	0.5	2.0	54	40
6	F 52	6.6	1.7	2.1	78	38
7	F 40	7.7	1.4	2.4	10	63

The balance techniques and methods of analysis have been described (Hanna *et al.*, 1960). Muscle biopsy specimens from Cases 1, 2, and 7 were analysed for their magnesium and calcium content by the method described by MacIntyre and Davidsson (1958).

Illustrative Cases

Three cases are described in some detail to illustrate the evolution of changes after parathyroidectomy. The first of these received no magnesium supplements, the second required acute magnesium therapy, while the third was maintained on oral magnesium post-operatively.

Case 4.—A woman aged 72 complained of pain and swelling in her knees for seven months. She had suffered from thirst, with polyuria and constipation, for one year. A nodule was palpable in the right lobe of the thyroid, and effusions were present in both knees. X-ray examination

FIG. 1.—Calcium and magnesium metabolic data in Case 4. In the balance studies in Figs. 1-3 stippling=urinary output and hatching=faecal output.

showed generalized rarefaction with subperiosteal resorption in the phalanges and patellae. Pre-operatively she was in negative calcium and magnesium balance with a urinary loss of 16 mEq of calcium and 5 mEq of magnesium a day (Fig. 1). After operation she remained on the same diet and calcium supplements as before (2 mEq/kg./day=65 mEq/day), but her balance was now positive for both calcium and magnesium. The urinary excretion of these elements had fallen to 2 mEq of each per day. Serum calcium fell to a minimum of 4.5 mEq/l. on the 13th post-operative day. Serum magnesium also fell post-operatively, reaching a minimum of 1.1 mEq/l. on the ninth post-operative day (normal 1.5–1.8 mEq/l.).

Case 2.—A 58-year-old woman complained of weakness of her legs for four years. Backache had been present for 18 months and there had been loss of height, loss of weight, anorexia, thirst, and polyuria. Examination revealed a dorsal kyphosis, swollen knees, and a nodule in the lower pole of the left lobe of her thyroid. X-ray examination showed generalized rarefaction with widespread subperiosteal erosions and nephrocalcinosis. Pre-operatively, on an intake of calcium of 16 mEq/day her faecal loss was 21 mEq/day and her urinary loss 9.3 mEq/day—a net balance of minus 14 mEq/day (Fig. 2). The corresponding magnesium figures were 10.5, 11.5, and 10.9 mEq/day. After parathyroidectomy the calcium intake was

FIG. 2.—Calcium and magnesium metabolic data in Case 2.

increased to 88 mEq/day (=2 mEq/kg./day); nevertheless, the serum calcium fell to 4.5 mEq/l., tetany appeared, and she required daily calcium infusions. Serum magnesium also fell, and on the ninth post-operative day reached 0.9 mEq/l. She became hostile and resentful, but these symptoms cleared completely when magnesium (100 mEq) was added to the daily infusion. In the two days after this infusion only 35 mEq of magnesium was excreted. Subsequent balances for calcium and magnesium showed net positive balances of 27.5 and 6.5 mEq/day respectively.

Case 7.—A 40-year-old woman presented with a four-months history of lethargy and backache. She had been asthmatic for many years, had sustained a haematemesis 10

years previously, and for six years had been troubled by gout. The only relevant finding on examination was a dorsal kyphosis. X-ray examination showed mild generalized rarefaction with slight wedging of the dorsal vertebrae. There was extensive nephrocalcinosis and a duodenal ulcer was shown by barium meal examination. As with the previous cases, negative preoperative calcium and magnesium balances became positive after removal of the parathyroid tumour. The significant finding was the prevention of a post-operative fall in the serum magnesium by oral supplements of magnesium (Fig. 3).

FIG. 3.—Calcium and magnesium metabolic data in Case 7.

Collected Data

Serum Calcium and Magnesium.—Pre-operatively the serum calcium was raised in all cases, ranging from 6.3 to 9.2 mEq/l. After parathyroidectomy it fell progressively to reach subnormal figures (Fig. 4). In Cases 1, 2, and 6 it was below 4 mEq/l. and there was associated tetany. It seems significant that these three cases had the highest levels of serum alkaline phosphatase (Table I). Pre-operatively the serum magnesium levels were close to the normal range except in Case 5 (Fig. 4); this patient had been vomiting excessively. There was no correlation between the levels of serum magnesium and those of serum calcium. The four patients who received no magnesium supplement after operation showed a distinct fall in their serum magnesium. Case 5, in contrast, on a supplement of 1 mEq/kg./day showed a progressive rise, and Case 7 (1.5 mEq/kg./day) showed no change. Case 6 on a similar supplement showed a slight fall of serum magnesium, but only to the lower limit of the normal range.

External Balances of Calcium and Magnesium.—All cases were in negative calcium balance before operation, even Cases 4 and 7, who were on large supplements of

FIG. 4.—Pre-operative and post-operative serum Mg and Ca in the seven cases.

FIG. 5.—Pre-operative and post-operative calcium data in the seven cases. In Figs. 5 and 6 the distance between case numbers and the base-line shows intestinal absorption, the distance between the short horizontal lines and the base-line shows urinary output. The intakes and net balances are also charted.

oral calcium (Fig. 5). Net absorption of calcium was poor and urinary losses were moderately high. After parathyroidectomy all, except Case 3, were in distinct positive balance. This was due chiefly to reduced urinary loss, but Cases 4 and 7, who were on the same calcium intake before and after operation, indicated that there was also increased net absorption. The poor net absorption in Case 3 was presumably due to concurrent steroid therapy. Magnesium balances before operation were in every case negative (Fig. 6). After parathyroidectomy they were all positive, due mainly to reduced urinary loss.

FIG. 6.—Pre-operative and post-operative magnesium data in the seven cases. In Figs. 5 and 6 the distance between case numbers and the base-line shows intestinal absorption, the distance between the short horizontal lines and the base-line shows urinary output.

Muscle Magnesium.—Table II shows the pre-operative muscle magnesium in Cases 1, 2, and 7 compared with three normal subjects. In Cases 1 and 2 the muscle magnesium was near the normal range; in Case 7 it was low. Muscle calcium was high in the low-magnesium sample.

TABLE II.—Muscle Magnesium and Calcium in Three Cases

Case No.	Mg	Ca
	mEq kg. Dry Fat-free Solids	
1	74	8.2
2	71	13.7
7	61	19.4
N.1	75	6.3
N.2	78	7.5
N.3	74	11.2

Discussion

From these results there seems to be no doubt that the removal of a parathyroid tumour leads to a retention of magnesium. This confirms the reports of a

number of previous workers mentioned above. Two questions immediately arise—Where has the retained magnesium collected? And how has this retention been brought about?

Magnesium does not accumulate in the extracellular fluid, for there is no rise in the serum magnesium. On the contrary, serum magnesium tends to fall, indicating a movement of magnesium out of the extracellular space. It could move into the cells or, alternatively, be deposited in the bones. Tibbetts and Aub (1937) found that the ratio of magnesium retention to calcium retention was greater than the ratio of magnesium to calcium in bone and concluded that some of the retained magnesium had been taken up by the soft tissues. This is a valid inference only if bone were of constant composition, which it is not. New bone crystals are smaller and consequently contain relatively larger hydration shells than old bone crystals. Moreover, it is thought (Neuman and Neuman, 1957, 1958) that much of the bone magnesium is held in these hydration shells. It follows that the Mg:Ca ratio of new bone would be expected to be higher than that of old bone. This explanation is supported by the finding of Shohl (1939) that the Mg:Ca ratio of bone rises after parathyroidectomy. Our own muscle biopsies are insufficient to offer direct evidence, but two of the cases had normal muscle magnesium before operation—refuting, in these cases, any suggestion of muscle depletion requiring restoration.

Prior to parathyroidectomy, bone resorption is occurring faster than bone replacement, with resulting thinning of bone. The magnesium liberated by this excessive resorption is in part used again to form new bone, but the surplus is lost from the body. Once the parathyroid tumour is removed, bone resorption slows up, while bone formation continues unabated. A fall in extracellular magnesium is a natural consequence.

Comparison of balances before and after operation indicate that the chief change which occurs is a fall in the urinary magnesium in the post-operative phase. This can be accounted for by the reduction in serum magnesium. The excessive losses pre-operatively, however, require further explanation. It should be stressed that this is by no means a constant finding; Cases 1 and 4 in the series reported here showed a very slight negative magnesium balance, and Bassett and Van Alstine (1935) described a case which was in a positive balance pre-operatively. The often-encountered negative balance can, however, be explained by the increased filtered load of calcium in the kidneys. This increase will lead to increased loss of magnesium because of competition for a common transport reabsorptive mechanism (Hanna *et al.*, 1960). The same conclusion can be drawn from studying the effect of magnesium infusion and magnesium supplement on urinary calcium in Cases 2 and 7 in this series. If the magnesium intake is not high enough, this obligatory increase in urinary loss will lead to a negative magnesium balance.

From a practical point of view it is important now to recognize that hypomagnesaemia with its associated dangers (Hanna *et al.*, 1960) is likely to occur after parathyroidectomy. Further, it seems from this limited series that oral supplements are sufficient to prevent a fall of serum magnesium and should therefore be given prophylactically in the post-operative period.

Summary

Primary hyperparathyroidism is sometimes associated with a negative magnesium balance because of increased urinary loss. This increase in urinary loss may be explained by an increase in the filtered load of calcium in the kidneys.

Parathyroidectomy in hyperparathyroid subjects produces: (a) lowering of plasma magnesium level, and (b) a positive magnesium balance.

These changes can be explained by the uptake of magnesium by bone.

The findings do not justify the proposition of a "primary" action of the parathyroid hormone on magnesium metabolism.

REFERENCES

- Barnes, B. A., Krahe, S. M., and Cope, O. (1957). *J. clin. Endocr.*, **17**, 1407.
 Bassett, S. H., and Van Alstine, H. E. (1935). *J. Nutr.*, **9**, 345.
 Bulger, H. A., and Gausmann, F. (1933). *J. clin. Invest.*, **12**, 1135.
 Greenberg, D. M., and Mackey, M. A. (1932). *J. biol. Chem.*, **98**, 765.
 Greenwald, I., and Cross, J. (1925). *Ibid.*, **66**, 217.
 Hanna, S., Harrison, M., MacIntyre, I., and Fraser, R. (1960). *Lancet*, **2**, 172.
 MacIntyre, I., and Davidsson, D. (1958). *Biochem. J.*, **70**, 456.
 Neuman, W. F., and Neuman, M. W. (1957). *Amer. J. Med.*, **22**, 123.
 — (1958). *The Chemical Dynamics of Bone Mineral*. Univ. of Chicago Press.
 Potts, J. T., jun., and Roberts, B. (1958). *Amer. J. med. Sci.*, **235**, 206.
 Shohl, A. T. (1939). *Mineral Metabolism*. Reinhold Publishing Corp., New York.
 Tibbetts, D. M., and Aub, J. C. (1937). *J. clin. Invest.*, **16**, 503.

INCIDENCE OF HYPOTHYROIDISM IN THE ELDERLY

BY

W. H. LLOYD, M.B., M.R.C.P.Ed.
Consultant Physician

AND

I. J. L. GOLDBERG,* B.Sc., M.R.C.S., L.R.C.P.
Late Assistant Pathologist

From the Geriatric Unit, Oldham and District General Hospital

In the past two and a half years we have been impressed by the high incidence of hypothyroidism seen in patients attending this geriatric department. The purpose of our paper is to illustrate this.

Material and Methods

During a period of two and a half years 93 patients attending the geriatric department were thought to be suffering from hypothyroidism. From these, 58 patients have been selected whom we regard as unequivocally hypothyroid. The material of this paper is derived from our observations on this group of 58.

Each case was carefully assessed clinically on the basis of a specially prepared form. Most cases were subjected to one or more laboratory tests. The accompanying Table shows the tests employed and the number of patients examined by each test. Apart from five patients

*Now at the Department of Pathology, University College of the West Indies, Mona, Kingston, Jamaica.