radically resected ampullary cancer. All patients underwent surgical resection for tumours of ampullary origin; only those with no macroscopic residual disease were considered suitable for the study. Pathological findings (tumour size and spread, and lymph node status) were obtained from the pathologists' original reports and were reassessed by our pathologists. Survival was defined as from the date of initial surgery to the date of death or the last contact. Follow-up data were available for all included patients.

The formalin-fixed, paraffin-embedded samples were sectioned at 5 µm and stained with H&E. The histological diagnoses were re-examined by two independent pathologists. In addition, the most representative blocks of each patient were selected to be cut into new 5 µm-thick sections for immunohistochemical studies. Immunohistochemical staining was performed by the streptavidin-biotin method. Expression of MUC2 and MUC5 was detected using two monoclonal antibodies: MUC1 (Dako) and MUC5 (Dako). Deparaffinised sections were incubated for 40 min with each primary monoclonal antibody. The expression of MUC2 and MUC5 was assessed according to the percentage of immunoreactive cells in a total of 1000 neoplastic cells (quantitative analysis). Immunoreactivity was graded as follows: positive, more than 5% of carcinoma cells stained; negative, no detectable staining or <5% of carcinoma cells stained. Known positive and negative tissue controls were included with each series. A univariate survival analysis for each prognostic variable on overall survival was estimated by the Kaplan-Meier method.⁸ The terminal event was death, attributable to cancer or non-cancer causes. The statistical significance of the differences in survival distribution among the prognostic groups was evaluated by the log-rank test⁹; $p < 0.05$ was regarded as significant in two-tailed tests. SPSS V.10 (SPSS, Chicago, IL, USA) was used for statistical analysis.

The cohort of patients consisted of 45 patients with pathological diagnosis of ampullary cancer (23 men and 22 women)

undergoing pancreaticoduodenectomy. Median age at diagnosis was 63.1 years (range 36–80). Seventeen patients were node positive and 28 were node negative. The median duration of follow-up after surgery was 46 months (range 13–102). Median overall survival was 44 months (range 6–100) and the 1, 3, and 5 year overall survival was 88.8%, 28.8% and 24.4% respectively. After a median follow-up of 46 months, 27 patients (60.0%) are still alive without evidence of disease, 2 patients (4.4%) are alive with recurrence of disease and 16 patients (35.5%) are dead (12 died because of ampullary carcinoma). Adjuvant radiotherapy and/or chemotherapy for ampullary cancer was not routinely offered in the hospitals involved in the study.

To determine the prognostic impact of MUC2 and MUC5 protein expression by univariate survival analysis, patients were stratified according to the dichotomised variables (criteria as stated above) into MUC2 and MUC5 negative versus MUC2 and MUC5 positive groups. By univariate analysis, overall survival was not influenced by MUC2 or MUC5 expression ($p = 0.2004$ and 0.7248 , respectively). In particular, the median survival time in patients with negative MUC2 expression was 53.55 months (95% CI 54.78 to 88.33 months) versus 69.00 months (95% CI 20.42 to 131.58 months) in patients with positive MUC2 expression. Median survival time in patients with negative MUC5 expression was 58.31 months (95% CI 38.87 to 77.75 months) versus 67.00 months (95% CI 62.90 to 71.10 months) in patients with positive MUC5 expression.

There is little substantial data reporting significant prognostic markers for ampullary cancer patients. An increasing interest in oncology research is now focused on the study of mucins in ampullary cancer, as molecular markers useful for a correct histological classification.^{5–6} For these reasons, for the first time in the literature, we have attempted to characterise the expression examining the possible prognostic significance of MUC2 and MUC5 in a homogeneous cohort of patients

with radically resected cancer of the ampulla of Vater. The present study reports the absence of a prognostic role of mucin expression in this type of cancer.

D Santini, A Baldi, B Vincenzi, P Mellone, M Campioni, A Antinori, D Borzomati, R Coppola, P Magistrelli, G Tonini

Università Campus Bio-Medico, Via E Longoni n° 83, 00155 Rome, Italy

Correspondence to: Dr Daniele Santini, Università Campus Bio-Medico, Via E Longoni n° 83, 00155 Rome, Italy; d.santini@unicampus.it

doi: 10.1136/jcp.2005.035832

Accepted 13 January 2006

Competing interests: None declared.

References

- 1 Gendler SJ, Spicer AP. Epithelial mucin genes. *Annu Rev Physiol* 1995;**57**:607–34.
- 2 Byrd JC, Bresalier RS. Mucins and mucin binding proteins in colorectal cancer. *Cancer Metastasis Rev* 2004;**23**:77–99.
- 3 Balague C, Gambus G, Carrato C, et al. Altered expression of MUC2, MUC4, and MUC5 mucin genes in pancreas tissues and cancer cell lines. *Gastroenterology* 1994;**106**:1054–61.
- 4 Yonezawa S, Sato E. Expression of mucin antigens in human cancers and its relationship with malignancy potential. *Pathol Int* 1997;**47**:813–30.
- 5 Hong SM, Cho H, Moskaluk CA, et al. CDX2 and MUC2 protein expression in extrahepatic bile duct carcinoma. *Am J Clin Pathol* 2005;**124**:361–70.
- 6 Fischer HP, Zhou H. Pathogenesis of carcinoma of the papilla of Vater. *J Hepatobiliary Pancreat Surg* 2004;**11**:301–9.
- 7 Zhou H, Schaefer N, Wolff M, et al. Carcinoma of the ampulla of Vater: comparative histologic/immunohistochemical classification and follow-up. *Am J Surg Pathol* 2004;**28**:875–82.
- 8 Kaplan EL, Meier P. Nonparametric estimation from incomplete observations. *J Am Stat Assoc* 1958;**53**:457–81.
- 9 Peto R, Pike MC, Armitage P, et al. Design and analysis of randomised clinical trials requiring prolonged observation of each patient. *Br J Cancer* 1977;**35**:1–39.

PostScript

LETTERS TO THE EDITOR

The JAK2 V617F mutation in Philadelphia-negative chronic myeloproliferative disorders

Buccal epithelial cells are occasionally used as a source of supposedly non-neoplastic DNA in patients suffering from haematological malignancies. Formerly, Kralovics *et al* found the JAK2 V617F mutation in DNA derived from buccal swabs in only 2.2% of patients with Philadelphia-negative chronic myeloproliferative disorders (Ph- CMPD).¹

We compared the JAK2 V617F mutational status in DNA derived from buccal swabs to that in DNA extracted from either bone marrow or peripheral blood in 35 Ph- CMPD

patients, including five cases of polycythaemia vera (PV), five cases of chronic idiopathic myelofibrosis (CIMF) and 25 cases of essential thrombocythaemia (ET). Genomic DNA was isolated from buccal swabs immediately on collection and from fresh-frozen bone marrow or peripheral blood, using Sherlock AX (A&A Biotechnology, Gdynia, Poland) and QIAamp DNA Mini Kit (Qiagen GmbH, Hilden, Germany) respectively. Allele-specific PCR for detecting the JAK2 V617F mutation was performed in duplicate samples with 100 ng DNA, as described previously.² PCR products were analysed by electrophoresis on 2% agarose gels containing ethidium bromide at 100 V for 1–2 h.

The mutation was detected in 26 samples derived from bone marrow or blood and in 24 of 26 (92.3%) samples derived from matching buccal swabs. In two ET cases the mutation

was noted in DNA derived from haematopoietic tissue only, and was absent in buccal swabs. All nine cases negative for the mutation in blood/marrow samples showed concordant negative results in the buccal swabs. The mutation-specific 203 bp band was occasionally weaker in DNA derived from swabs compared to DNA of a direct haematopoietic origin (fig 1).

The high rate of mutation detection in our material might have resulted from a high sensitivity of the allele-specific PCR. Also of some potential importance was the technique of buccal swab sampling, with generous bilateral collection of the material from patients' oral mucosa. Buccal swabs are most likely contaminated with leucocytes and removal of non-epithelial cells by mouth rinsing was suggested as a precaution in chimerism studies.³ A more intriguing

Figure 1 JAK2 V617F mutation in matched buccal swab (s) and bone marrow (m) samples from patients with (lanes 1–5) and without the mutation (lanes 6–8). The mutation is represented by a 203 bp band. The 364 bp band serves as an internal JAK2-specific control. PV, polycythaemia vera; CIMF, chronic idiopathic myelofibrosis; TE, essential thrombocythaemia.

hypothesis, explaining why a genetic event apparently specific for the somatic haematopoietic tissue, is detected in buccal swabs may be proposed based on recently described trans-differentiation of haematopoietic cells into epithelial cells in allogeneic bone marrow transplant recipients.^{4,5}

The exact origin of the mutant DNA notwithstanding, buccal swabs appear to be a surprisingly good source of material for JAK2 mutation studies, provided that a sensitive mutation detection system is used. Their collection is much less invasive and cheaper, even when compared to collection of peripheral blood. On the other hand, caution has to be exercised when DNA derived from buccal swabs is planned to be used as a source of supposedly “wild-type” genetic material for comparison with DNA derived from blood or marrow, in the studies of mutations expected to be specific for haematopoietic tissues.

Anastazja Stój, Zbigniew Rudzki,
Jerzy Stachura

Department of Pathomorphology, Collegium Medicum,
Jagiellonian University, Kraków, Poland

Tomasz Sacha, Sylwia Czekalska

Department of Haematology, Collegium Medicum,
Jagiellonian University, Kraków, Poland

Correspondence to: Dr Zbigniew Rudzki, Consultant
Histopathologist, Department of Histopathology,
Birmingham Heartlands Hospital, Bordesley Green East,
Bordesley B9 5SS, Birmingham, UK; zbigniew.rudzki@
heartofengland.nhs.uk

doi: 10.1136/jcp.2006.044644

Accepted 27 October 2006

Funding: Supported by the Committee for Scientific
Research, Republic of Poland, grant no 3 P05B 084 24.

Competing interests: None declared.

References

- Kralovics R, Passamonti F, Buser AS, *et al*. A gain-of-function mutation of JAK2 in myeloproliferative disorders. *N Engl J Med* 2005;**352**:1779–90.
- Baxter EJ, Scott LM, Campbell PJ, *et al*. Acquired mutation of the tyrosine kinase JAK2 in human myeloproliferative disorders. *Lancet* 2005;**365**:1054–61.
- Thiede C, Prange-Krex G, Freiberg-Richter J, *et al*. Buccal swabs but not mouthwash samples can be used to obtain pretransplant DNA fingerprints from recipient of allogeneic bone marrow transplants. *Bone Marrow Transplant* 2000;**25**:575–7.
- Metaxas Y, Zeiser R, Schmitt-Graeff A, *et al*. Human hematopoietic cell transplantation results in generation of donor-derived epithelial cells. *Leukemia* 2005;**19**:1287–9.
- Tran SD, Pillemer SR, Dutra A, *et al*. Differentiation of human bone marrow-derived cells into buccal

epithelial cells in vivo: a molecular analytical study. *Lancet* 2003;**361**:1084–8.

Mouse mammary tumour-like virus gene sequences and specific breast cancer morphology

We previously reported that the histological characteristics of some human breast tumours in which mouse mammary tumour-like virus (MMTV) gene sequences had been identified, were similar to mammary tumours from C3H experimental mice.¹ Such mammary tumours in mice are usually associated with MMTV.

We have now confirmed that the mouse mammary tumours were MMTV positive (as we had assumed but not demonstrated). Standard PCR analyses were conducted on DNA extracted from 10 (unselected) of the 44 archival formalin fixed mouse mammary tumour specimens previously described. The PCR methods used were identical to those used for the human breast tumour specimens, again as previously described. The analyses were repeated three times. The products of the PCR for three samples were sequenced.

All 10 mouse mammary tumour specimens were positive for MMTV gene sequences. All three PCR product samples that were sequenced gave an exact match with AF228552, the complete exogenous mouse mammary tumor virus C3H proviral genome.

These data confirm that MMTV gene sequences are present in mouse mammary tumours that occur in experimental C3H mice as used in our investigations.

Wendy K Glenn, James S Lawson,
Noel J Whitaker

School of Biotechnology and Biomolecular Science,
University of New South Wales, Sydney, Australia

Correspondence to: Professor James S Lawson, School
of Biotechnology and Biomolecular Science, University
of New South Wales, Sydney, Australia;
james.lawson@unsw.edu.au

doi: 10.1136/jcp.2006.044487

Accepted 17 November 2006

Published Online First 8 December 2006

Competing interests: None declared.

Reference

- Lawson JS, Tran DD, Carpenter E, *et al*. Presence of mouse mammary tumour-like virus gene sequences may be associated with morphology of specific human breast cancer. *J Clin Pathol* 2006;**59**:1287–92.

Helicobacter pylori-associated peptic ulcer and florid follicular gastritis mimicking a malignancy

In chronic follicular gastritis the lymphoid follicles are usually localised in the lamina propria.¹ In lymphoid hyperplasia associated with a chronic peptic ulcer, numerous lymphoid follicles are sometimes present in the ulcer base, in which the muscularis propria has usually been destroyed and replaced by scar tissue.² The histology of *Helicobacter*-associated chronic gastritis with numerous reactive lymphoid follicles and an indistinct poorly developed marginal zone may closely simulate lymphoma and pose a diagnostic challenge. We present an extreme example of florid follicular gastritis, which mimicked a malignant neoplasm clinically and pathologically. To our knowledge, such florid hyperplasia with reactive lymphoid follicles extending into the muscularis propria and beyond has never been reported.

A 65-year-old man presented with intermittent abdominal hunger pain for one month. Endoscopic biopsies revealed a Bormann type II antral ulcer with *H. pylori* infection. Endoscopic findings and image study strongly suggested gastric cancer; a subtotal gastrectomy and regional lymph node dissection were performed. On the lesser curvature side of the antrum, there was a tumour-like lesion, 4×3×1 cm with ulceration and slightly elevated margins. The cut surface was whitish and firm with effacement of the gastric muscle wall (fig 1). Microscopically, sections of stomach show ulcerated antral mucosa with an adjacent and underlying intense chronic inflammatory infiltrate comprising reactive germinal centres surrounded by a narrow mantle zone with some peripheral lymphocytes. Beneath the ulcer there was intense scarring with destruction of the muscularis propria, which was preserved lateral to the ulcer margins. The hyperplastic lymphoid follicles extended into the scar tissue, muscularis propria and serosa (fig 2). Bacilli morphologically typical of *H. pylori* were identified. There were no lymphoepithelial lesions or aggregates of centrocyte-like atypical lymphocytes to suggest mucosa-associated lymphoid tissue (MALT) lymphoma. The dissected regional nodes measured up to 2.2 cm in diameter and revealed marked follicular hyperplasia.

Immunohistochemistry showed that the germinal centre cells were positive for CD10 and bcl-6 but not bcl-2 and they expressed polytypic light chains. The surrounding mantle zone lymphocytes expressed immunoglobulin

Figure 1 Cut surface of the antral ulcerative lesion is whitish and firm with effacement of the gastric wall.