ORIGINAL ARTICLE

Rapid identification of *Candida* species in oral rinse solutions by PCR

Giorgio Liguori, Angela Lucariello, Giuseppe Colella, Antonio De Luca, Paolo Marinelli

J Clin Pathol 2007;60:1035–1039. doi: 10.1136/jcp.2006.040782

See end of article for authors' affiliations

Correspondence to: Professor Giorgio Liguori, Chair of Hygiene and Epidemiology, Faculty of Movement Sciences, University of Naples "Parthenope", Via Medina 40, 80133 Naples, Italy; giorgio.liguori@uniparthenope.it

Accepted 29 September 2006

Aims: To determine the sensitivity and specificity of a multiplex PCR assay for the contemporary identification of major species involved in oral candidiasis, without extraction and purification of DNA from the samples under investigation; and evaluation of this method in comparison with routine phenotypic culture identification.

Methods: 78 oral rinse solutions were collected. The concentrated oral rinse technique was used for a quantitative and qualitative study. Research and identification of *Candida* spp, with routine phenotypic culture identification (germ-tube test in serum at 37°C for 3 hours and sugar assimilation strip analysis), were performed. Each sample was analysed with multiplex PCR directly on oral rinse solution. Samples giving discrepant results between routine phenotypic and PCR identification methods were resubcultured on CHROMagar *Candida* plates. The fungus-specific primers ITS1, ITS2, CA3, and CA4 were used. For the identification of other species (*C kefyr*, *C famata* and *C dubliniensis*), ITS1F, ITS1K, and ITS2D primers were designed.

Results: Multiplex PCR correctly identified all samples, including those with single species, or with mixed species, negative samples and positive samples which appeared to be negative from routine phenotypic methods.

Conclusion: This multiplex PCR assay provides a rapid alternative to the conventional culture based technique for the identification and speciation of the most frequently isolated *Candida* species. The absence of an extraction method made identification of 10 species possible in a few hours.

Oral candidosis is one of the most frequently observed pathologies in everyday practice.¹ It is an opportunistic disease because of some promoting factors that alter the equilibrium in the oral cavity microbial ecosystem, helping the transformation of yeasts from commensals to pathogens.² Although the transition from commensalism to disease may be associated with the virulence characteristics of the organism, it is widely accepted that host factors (iatrogenic and infective factors) are of paramount importance in the development of the infection.³

A number of relevant factors have been associated with oral carriage of yeast (denture wearing, periodontal disease, leucoplasic lesions, etc). They contribute to form the so-called "mycotic count" and to cause the following clinical cases.⁴

Candida albicans has been by far the most common yeast species carried as a commensal by healthy individuals and the agent most frequently responsible for oral yeast infection. However, the emergence of non-*albicans* *Candida* species (*C parapsilosis*, *C tropicalis*, *C glabrata*, etc) has also been observed and associated with long-term treatment with amphotericin and fluconazole.^{5–14}

More recently, oral candidosis has been associated with *C dubliniensis* in HIV-infected individuals^{15–16} and patients with diabetes,¹⁷ and has been detected in the sputum of patients with cystic fibrosis.¹⁸

Since pathogenicity and antifungal susceptibility often vary among species, a rapid and accurate identification of the disease-causing species of *Candida* is crucial for both clinical treatment and epidemiological studies of oral candidiasis.¹⁹ The conventional identification of pathogenic fungi in the clinical microbiology laboratory involves the examination of colony and microscopic morphologies and the assessment of various biochemical reactions.^{20–21} It often requires three or more days, and may be inaccurate. Moreover, the presence of more than

one *Candida* species in the oral cavity of the same host is not infrequent.

In recent years, numerous DNA-based methods such as DNA–DNA reassociation,²² DNA fingerprinting,²³ and Southern hybridisation with appropriate DNA probes^{24–25} have been reported to recognise *Candida* species in culture or in clinical materials. However, these genotypic methods have the disadvantage of being laborious and time-consuming, and also require specialised equipment.

The aims of the present study were to determine the sensitivity and specificity of a multiplex PCR method applied to oral rinse solution obtained in our laboratory, for the identification of *Candida* species that are frequently isolated from the oral cavity, and to evaluate this method in comparison with routine phenotypic culture identification. The method is based on the size variability of the ITS1 region in different species and on the amplification of a specific DNA fragment of the ITS2 region of *C albicans*. The test has allowed us to identify *Candida* species using seven species-specific oligonucleotides in a single PCR tube.

Drawing from this strategy we studied the structure of other species of *Candida* and generated in the same region other primers that were able to amplify specific fragments. Indeed for the identification of other three species, *C kefyr*, *C famata* and *C dubliniensis* we designed the following primers: ITS1F (5'-CCA GCG CTT AAT TGC G-3'), ITS1K (5'-ATC GTC TGA ACA AGG CCT GC-3'), and ITS2D (5'-GAG AAC CAA GAG ATC CGT TGT TG-3'). Table 1 lists these primers and the generated PCR products.

MATERIALS AND METHODS

A total of 78 oral rinse solutions obtained from symptomatic patients with odontostomatological and maxillo-facial diseases, with or without signs, attending the outpatients clinic of the

Figure 1 Diagram of hybridisation sites of ITS1 and ITS2 primers.

Microbiology Laboratory of the Hygiene Unit of the Department of Public, Clinical and Preventive Medicine were collected during the period 2004 to 2005.

The concentrated oral rinse technique for the determination of bacterial and mycotic colony counts (a gargle with 9 ml of phosphate-buffered saline) was used for a quantitative and qualitative study as previously described.²⁶ From all positive cultures, *Candida* genus yeasts were identified presumptively (germ-tube test in serum at 37°C for 3 hours) and definitively with API 20C AUX (bioMérieux, Rome, Italy) sugar assimilation strip analysis. Each sample was analysed with multiplex PCR directly on oral rinse solution. Samples giving discrepant results between routine phenotypic and PCR identification methods were resubcultured on CHROMagar *Candida* (Alfa Wassermann, Diagnostics SpA, Milan, Italy) plates to identify possible mixed yeast culture. PCR was performed directly on each oral rinse solution (without preliminary DNA extraction).

Patients gargled with 5 ml of phosphate-buffered saline, and the resulting solution was centrifuged at 14000 rpm for 5 min. The supernatant was discarded. The residual pellet was resuspended with 100 µl sterile distilled water and heated at 90°C for 3 min; 20 µl samples were added to the mix for PCR.

The fungus-specific primers ITS1 (5'-TCC GTA GGT GAA CCT GCG G-3') and ITS2 (5'-GCT GCG TTC TTC ATC GAT CG-3')²⁷ were used to amplify a small conserved portion of the 18S rDNA region, the adjacent ITS1, and a small portion of the 28S rDNA region (fig 1), generating PCR products for *C. glabrata*, *C. guilliermondii*, *C. lusitanae*, *C. parapsilosis*, *C. tropicalis* and *C. krusei*. In addition, *C. albicans*-specific primers CA3 (5'-GGT TTG GAA AGA CGG TAG-3') and CA4 (5'-AGT TTG AAG ATA TAC GTG GTA G-3')²⁸ were also included in the PCR mixture to amplify a portion of the ITS2 region of *C. albicans*.

Multiplex PCR was performed in duplicate with 20 µl of oral rinse solution in a total reaction volume of 50 µl, consisting of 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl₂, 0.8 mM deoxyribonucleoside triphosphates (0.2 mM each), 3.2 µM primers, *Taq* DNA polymerase (1.25 U), and 50 µl of a mineral oil overlay. PCR was carried out with a Perkin Elmer thermal cycler under the following condition: initial denaturation, 92°C, 2 min; 35 cycles of denaturation (95°C, 1 min), annealing (50°C, 1 min), and extension (72°C, 1 min); and final extension, 72°C, 10 min. A negative control run was performed with each test run by replacing the samples with sterilised water in

the PCR mixture. A positive culture broth containing *C. albicans* was run in parallel with unknown samples, and this culture broth was used as a positive control. Gel electrophoresis was conducted in TBE buffer (0.1 M Tris, 0.09 M boric acid, 1 mM EDTA, pH 8.4) at 100 V for 1–2 h using gel composed of 2% (w/v) agarose (Sigma-Aldrich, Milan, Italy). Gel was stained with 0.5 µg ethidium bromide per ml distilled water.

RESULTS

A total of 78 oral rinse solutions were analysed by routine phenotypic methods for species identification. From these samples, 63 strains of yeasts were isolated; 15 oral rinse solutions were negative. The most frequently isolated species was *C. albicans* (48 strains, 76.2%), followed by *C. glabrata* (6 strains, 9.5%), *C. tropicalis* (5 strains, 7.9%), *C. krusei* (2 strains, 3.2%), *C. parapsilosis* (1 strain, 1.6%), and *C. famata* (1 strain, 1.6%) (table 2).

Multiplex PCR identification results matched phenotypic identification results for 45 of 64 positive samples (70.3%) (47 samples containing *C. albicans*, 73.4%; 3 *C. glabrata*, 4.7%; 2 *C. krusei*, 3.1%; 2 *C. dubliniensis*, 3.1%; 1 *C. tropicalis*, 1.6%; 1 *C. parapsilosis*, 1.6%; 1 *C. famata*, 1.6%; 1 *C. guilliermondii*, 1.6%; 1 *C. kefir*, 1.6%). Three samples that were found to contain *C. albicans* only by phenotypic methods were identified by PCR to contain both *C. albicans* and *C. glabrata*. The subculture of the oral rinse solution on CHROMagar *Candida* medium indicated that this specimen contained a mixture of *C. albicans* and *C. glabrata* (table 3).

Two samples that were found to contain *C. albicans* only by phenotypic methods were identified by PCR to contain both *C. glabrata* and *C. guilliermondii* in one case and *C. tropicalis* and *C. guilliermondii* in another. In these two cases of mixed species, *C. glabrata* and *C. guilliermondii* and *C. tropicalis* and *C. guilliermondii* were also recovered on CHROMagar *Candida* plates (table 3).

Four oral rinse solutions that were found to contain *C. tropicalis* only by phenotypic methods were identified by PCR to contain 3 *C. albicans* and 1 *C. guilliermondii*; 1 sample containing *C. parapsilosis* was identified by PCR to contain *C. albicans*; 3 samples containing *C. glabrata* were identified by molecular analysis to contain 3 *C. albicans*; and 5 oral rinse solutions containing *C. albicans* were identified by multiplex PCR to contain 1 *C. parapsilosis*, 1 *C. glabrata*, 1 *C. kefir*, and 2 *C. dubliniensis*. The subculture of all these oral rinse solutions on CHROMagar

Table 1 Lengths of the PCR products

	Primers									
Organism	ITS1	ITS1F	ITS1K	ITS2	ITS2D	CA3	CA4	PCR product (bp)		
<i>C. glabrata</i>	+	—	—	+	+	—	—	482–483	462–463	—
<i>C. guilliermondii</i>	+	—	—	+	+	—	—	248	228	—
<i>C. famata</i>	—	+	—	+	+	—	—	234	214	—
<i>C. kefir</i>	—	—	+	+	+	—	—	249	229	—
<i>C. parapsilosis</i>	+	—	—	+	+	—	—	229	209	—
<i>C. tropicalis</i>	+	—	—	+	+	—	—	218	199	—
<i>C. albicans</i>	+	—	—	+	+	+	+	218–219	198–199	110
<i>C. krusei</i>	+	—	—	+	+	—	—	182	166	—
<i>C. lusitanae</i>	+	—	—	+	+	—	—	148	128	—
<i>C. dubliniensis</i>	+	—	—	—	+	—	—	—	198	—

Table 2 Comparative identification of *Candida* spp in oral rinse solutions by routine phenotypic identification methods versus multiplex PCR

Routine phenotypic identification	No of oral rinse solutions analysed	Multiplex PCR identification	No of oral rinse solutions analysed
<i>C albicans</i>	48	<i>C albicans</i>	47
<i>C glabrata</i>	6	<i>C albicans</i> + <i>C glabrata</i>	3
<i>C tropicalis</i>	5	<i>C glabrata</i>	3
<i>C krusei</i>	2	<i>C glabrata</i> + <i>C guilliermondii</i>	1
<i>C parapsilosis</i>	1	<i>C tropicalis</i>	1
<i>C famata</i>	1	<i>C tropicalis</i> + <i>C guilliermondii</i>	1
<i>C guilliermondii</i>	0	<i>C krusei</i>	2
<i>C dubliniensis</i>	0	<i>C parapsilosis</i>	1
<i>C kefyr</i>	0	<i>C famata</i>	1
Negative	15	<i>C guilliermondii</i>	1
		<i>C dubliniensis</i>	2
		<i>C kefyr</i>	1
		Negative	14
Total	78	Total	78

Candida medium confirmed the identification of multiplex PCR except for *C dubliniensis* (table 3).

Finally, one of the oral rinse solutions which was negative with the phenotypic method was positive to the multiplex PCR (*C albicans*).

The time from the first oral rinse solution culture to species identification by routine phenotypic methods (subculture, germ tube and API 20CAUX) was 4 days (96 h). More slowly growing, individual isolates took as long as 6 days for growth and species identification by routine phenotypic methods (144 h, *C guilliermondii* isolate). For the multiplex PCR, on the other hand, no time was required to isolate and extract DNA from oral rinse solution, 3 h was required for PCR amplification and 1.5 h was required for agarose gel electrophoresis analysis. Therefore, species could be identified in as little as 5 h, including the time required to prepare PCR reagent mixture, in contrast to routine phenotypic methods which took several days.

The mycotic and bacterial colony count obtained from the oral rinse technique used for a quantitative study ranged from

10 CFU/ml to 5.46×10^6 CFU/ml for the mycotic count and from 9.8×10^5 CFU/ml to 15.3×10^7 CFU/ml for the bacterial colony count.

The limit of detection of the multiplex PCR was approximately 10 CFU/ml and was very close to that reported by Jaeger *et al.*²⁹

The high quantity of bacterial colony count did not interfere with the detection and identification of yeasts with multiplex PCR. In fact, coexisting bacteria in oral rinse solution specimens did not produce any detectable PCR products. Therefore, the multiplex PCR method could also detect mixed yeast cultures missed by routine subculturing methods, even when the oral rinse solution contained bacteria (fig 2).

DISCUSSION

In order to improve the quality of our diagnostic research, we have set up a multiplex PCR assay for the contemporary identification of major species involved in oral candidiasis to avoid any time delay due to the extraction and purification of DNA from the samples under investigation.

Table 3 Resolution of discrepancies between phenotypic and multiplex PCR methods for the identification of *Candida* species

Identification by the following methods:			
Patient no	Routine phenotypic method	Multiplex PCR	Final identification*
293	<i>C albicans</i>	<i>C albicans</i> + <i>C glabrata</i>	<i>C albicans</i> + <i>C glabrata</i>
301	<i>C glabrata</i>	<i>C albicans</i> + <i>C glabrata</i>	<i>C albicans</i> + <i>C glabrata</i>
315	<i>C tropicalis</i>	<i>C albicans</i>	<i>C albicans</i>
330	<i>C tropicalis</i>	<i>C albicans</i>	<i>C albicans</i>
333	<i>C tropicalis</i>	<i>C albicans</i>	<i>C albicans</i>
348	<i>C albicans</i>	<i>C albicans</i> + <i>C glabrata</i>	<i>C albicans</i> + <i>C glabrata</i>
358	<i>C parapsilosis</i>	<i>C albicans</i>	<i>C albicans</i>
362	<i>C albicans</i>	<i>C glabrata</i> + <i>C guilliermondii</i>	<i>C glabrata</i> + <i>C guilliermondii</i>
374	<i>C glabrata</i>	<i>C albicans</i>	<i>C albicans</i>
392	<i>C glabrata</i>	<i>C albicans</i>	<i>C albicans</i>
420	<i>C albicans</i>	<i>C parapsilosis</i>	<i>C parapsilosis</i>
421	<i>C albicans</i>	<i>C tropicalis</i> + <i>C guilliermondii</i>	<i>C tropicalis</i> + <i>C guilliermondii</i>
424	<i>C albicans</i>	<i>C glabrata</i>	<i>C glabrata</i>
422	<i>C tropicalis</i>	<i>C guilliermondii</i>	<i>C guilliermondii</i>
427	<i>C albicans</i>	<i>C dubliniensis</i>	<i>C albicans</i>
433	<i>C albicans</i>	<i>C kefyr</i>	<i>C kefyr</i>
441	<i>C albicans</i>	<i>C dubliniensis</i>	<i>C albicans</i>
451	<i>C glabrata</i>	<i>C albicans</i>	<i>C albicans</i>
Negative	Negative	<i>C albicans</i>	Negative

*Final species identification was corroborated by colony morphology and colour on CHROMagar *Candida* medium and by reidentification by routine phenotypic methods.

Figure 2 Identification of yeasts present in oral rinse solutions by the multiplex PCR. M, 50 bp DNA ladder. Lane 1, *C. kefyr* bp 249. Lane 2, *C. famata* bp 234. Lane 3, *C. dubliniensis* bp 198. Lane 4, *C. krusei* bp 182. Lane 5, *C. parapsilosis* bp 209. Lane 6, *C. guilliermondii* bp 248. Lane 7, *C. albicans* bp 110–218. Lane 8, *C. albicans* bp 110–218. Lane 9, *C. albicans* bp 110–218 and *C. glabrata* bp 462–482. Lane 10, *C. albicans* bp 110–218. Lane 11, *C. glabrata* bp 462–482. Lane 12, *C. tropicalis* bp 218. Lane 13, *C. glabrata* bp 462–482; Lane 14, *C. albicans* bp 110–218. Lane 15, *C. albicans* bp 110–218. Lane 16, *C. albicans* bp 110–218 and *C. glabrata* bp 462–482. Lane 17, *C. albicans* bp 110–218. Lane 18, *C. albicans* bp 110–218. Lane 19, *C. glabrata* bp 462–482.

To the best of our knowledge this is the first application of a PCR assay on oral rinse solutions obtained directly from the samples without DNA extraction.

A total of 78 oral rinse solutions were used for a PCR assay to simultaneously identify and type the *Candida* positivity directly from the clinical samples without any particular treatment. Multiplex PCR correctly identified all 78 samples, including 58 cultures from subjects with single species, 5 cultures of oral rinses from patients with colonisation of mixed species (identified to contain a single *Candida* species by routine phenotypic methods), 14 negative samples and 1 positive sample identified as negative from routine phenotypic methods.

The advantages of this method are as follows: it does not require use of expensive or toxic chemical substances such as proteinase K or phenol-chloroform; the total time from species identification is 5 h, compared to a mean of 5 days by routine phenotypic culture identification methods; identification of more than one species of yeast is possible in mixed cultures, with no cross-reaction or interference with bacteria and viruses likely to be present in oral rinse solutions; and no modifications or changes in routine practice were necessary for the clinicians or the laboratory technicians, who were able to use the same samples used for conventional morphological and metabolic examination for the PCR assay.

Finally, considering the continuous decrease of the cost of purchasing a thermal cycler and the reagents necessary to use it, the estimated cost for this assay is similar to that of routine phenotypic culture identification.

In our experience, at least 10 species of *Candida* are frequently isolated from oral rinse solutions in laboratories: *C. albicans*, *C. glabrata*, *C. guilliermondii*, *C. lusitanae*, *C. parapsilosis*, *C. tropicalis*, *C. krusei*, *C. kefyr*, *C. famata* and *C. dubliniensis*. In order to be able to detect the most common *Candida* species, we have found in the literature primers to detect *C. albicans*, *C. glabrata*, *C. guilliermondii*, *C. lusitanae*, *C. parapsilosis*, *C. tropicalis* and *C. krusei*.³⁰ Moreover, in order to increase the number of primers to include all the medically important species of *Candida*, drawing from data in the literature, we have designed other primers in the region between the conserved portion of the 18S rDNA region, the adjacent ITS1, and a small portion of the 28S rDNA region in order to detect *C. kefyr*, *C. famata* and *C. dubliniensis*.

Take-home messages

- A method for rapid identification by PCR of the presence of *Candida* spp has been developed (5 h versus a mean of 5 days by routine phenotypic culture identification methods).
- The PCR assay is performed on the oral rinse solution obtained directly from the samples without DNA extraction.
- The technique is accurate, and permits identification of more than one species of yeast in mixed cultures, with no cross-reaction or interference with bacteria and viruses likely to be present in oral rinse solutions.
- The method could be used routinely in clinical microbiology laboratories.
- The method permits rapid identification of *Candida* isolates from high-risk patients or patients with oral candidiasis, enabling prompt and appropriate treatment and thus reducing hospitalisation time.

The multiplex PCR is particularly suitable for use in a routine clinical microbiology laboratory because it can easily be automated; moreover it could be utilised for the rapid identification of *Candida* isolates from high-risk patients or patients with oral candidiasis, enabling prompt and appropriate treatment and thus reducing hospitalisation time.

ACKNOWLEDGEMENTS

We thank Dr Pia Furno for editing the manuscript.

Authors' affiliations

G Liguori, A Lucariello, Chair of Hygiene and Epidemiology, Faculty of Movement Sciences, University of Naples "Parthenope", Naples, Italy
G Colella, Department of Head and Neck Pathology, Second University of Naples, Naples, Italy
A De Luca, P Marinelli, Department of Public, Clinical and Preventive Medicine, Second University of Naples, Naples, Italy

Funding: This work was supported in part by University of Naples "Parthenope" and Second University of Naples.

Competing interests: None declared.

REFERENCES

- 1 Reichart PA, Samaranayake LP, Philipsen HP. Pathology and clinical correlates in oral candidiasis and its variants: a review. *Oral Dis* 2000;**6**:85–91.
- 2 Rogers TJ, Balish E. Immunity to *Candida albicans*. *Microbiol Rev* 1980;**44**:660–82.
- 3 Saltarelli CG. *Candida albicans. The pathogenic fungus*. New York: Hemisphere Publishing, 1989.
- 4 Odds FC. *Candida and candidosis*. London: Balliere Tindall, 1988.
- 5 Nguyen MH, Peacock JE Jr, Morris AJ, et al. The changing face of candidemia: emergence of non-*Candida albicans* species and antifungal resistance. *Am J Med* 1996;**100**:617–23.
- 6 Nielsen H, Stenderup J. Invasive *Candida norvegensis* infection in immunocompromised patients. *Scand J Infect Dis* 1996;**28**:311–12.
- 7 Verduyn Lunel FM, Meis JF, Voss A. Nosocomial fungal infections: candidemia. *Diagn Microbiol Infect Dis* 1999;**34**:213–20.
- 8 Viscoli C, Girmenia C, Marinus A, et al. Candidemia in cancer patients: a prospective, multicenter surveillance study by the Invasive Fungal Infection Group (IFIG) of the European Organization for Research and Treatment of Cancer (EORTC). *Clin Infect Dis* 1999;**28**:1071–9.
- 9 Newman SL, Flanagan TP, Fisher A, et al. Clinically significant mucosal candidiasis resistant to fluconazole treatment in patients with AIDS. *Clin Infect Dis* 1994;**19**:684–6.
- 10 Pfaller MA, Jones RN, Messer SA, et al. National surveillance of nosocomial blood stream infection due to *Candida albicans*: frequency of occurrence and antifungal susceptibility in the SCOPE Program. *Diagn Microbiol Infect Dis* 1998;**31**:327–32.
- 11 Price MF, LaRocco MT, Gentry LO. Fluconazole susceptibilities of *Candida* species and distribution of species recovered from blood cultures over a 5-year period. *Antimicrob Agents Chemother* 1994;**38**:1422–4.
- 12 Rex JH, Rinaldi MG, Pfaller MA. Resistance of *Candida* species to fluconazole. *Antimicrob Agents Chemother* 1995;**39**:1–8.
- 13 Ruhnke M, Eigler A, Tennagen I, et al. Emergence of fluconazole-resistant strains of *Candida albicans* in patients with recurrent oropharyngeal candidosis and human immunodeficiency virus infection. *J Clin Microbiol* 1994;**32**:2092–8.
- 14 Wingard JR, Merz WG, Rinaldi MG, et al. Increase in *Candida krusei* infection among patients with bone marrow transplantation and neutropenia treated prophylactically with fluconazole. *N Engl J Med* 1991;**325**:1274–7.
- 15 Coleman DD, Sullivan B, Harrington K, et al. Molecular and phenotypic analysis of *Candida dubliniensis*: a recently identified species linked with oral candidosis in HIV-infected and AIDS patients. *Oral Dis* 1997;**3**:S96–101.
- 16 Sullivan D, Coleman D. *Candida dubliniensis*: an emerging opportunistic pathogen. *Curr Top Med Mycol* 1997;**8**:15–25.
- 17 Manfredi M, McCullough MJ, Al-Karaawi ZM, et al. The isolation, identification and molecular analysis of *Candida* spp isolated from the oral cavities of patients with diabetes mellitus. *Oral Microbiol Immunol* 2002;**17**:181–5.
- 18 Peltroche-Llacsahuanga H, Dohmen H, Haase G. Recovery of *Candida dubliniensis* from sputum of cystic fibrosis patients. *Mycoses* 2002;**45**:15–18.
- 19 Williams DW, Lewis MA. Isolation and identification of *Candida* from the oral cavity. *Oral Dis* 2000;**6**:3–11.
- 20 Warren NG, Hazen KC. *Candida*, *Cryptococcus*, and other yeasts of medical importance. In: Murray PR, Baron EJ, Pfaller MA, Tenover FC, Tenover RH, eds. *Manual of clinical microbiology*, 6th edn. Washington, DC: American Society for Microbiology, 1995:723–37.
- 21 Wickes LB, Hicks JB, Merz WG, et al. The molecular analysis of synonymy among medical important yeasts within the genus *Candida*. *J Gen Microbiol* 1992;**138**:901–7.
- 22 Kurtzman CP, Phaff HJ, Meyer SA. Nucleic acid relatedness among yeasts. In: Spencer JFT, Spencer DM, Smith ARW, eds. *Yeast genetics: fundamental and applied aspects*. New York: Springer-Verlag, 1983:139–66.
- 23 Scherer S, Stevens DA. Application of DNA typing methods to epidemiology and taxonomy of *Candida* species. *J Clin Microbiol* 1987;**25**:675–9.
- 24 Magee BB, D'Souza TM, Magee PT. Strain and species identification by restriction fragment length polymorphisms in the ribosomal DNA repeat of *Candida* species. *J Bacteriol* 1987;**163**:9–43.
- 25 Mason MM, Lasker BA, Riggsby WS. Molecular probe for identification of medically important *Candida* species and *Torulopsis glabrata*. *J Clin Microbiol* 1987;**25**:563–6.
- 26 Samaranayake LP, MacFarlane TW, Lamey PJ, et al. A comparison of oral rinse and imprint sampling techniques for the detection of yeast, coliform and *Staphylococcus aureus* carriage in the oral cavity. *J Oral Pathol* 1986;**15**:386–8.
- 27 White TJ, Bruns T, Lee S, et al. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds. *PCR protocols: a guide to methods and applications*. San Diego, CA: Academic Press, 1990:315–22.
- 28 Huang CH. Specific identification of yeasts on the bases of PCR amplified ribosomal DNA immobilized by covalent bond on the piezoelectric quartz crystal. PhD thesis. National Taiwan University, Taipei, Taiwan, 1996.
- 29 Jaeger EEM, Carroll NM, Choudhury S, et al. Rapid detection and identification of *Candida*, *Aspergillus*, and *Fusarium* species in ocular samples using PCR. *J Clin Microbiol* 2000;**38**:2902–8.
- 30 Chang HC, Leaw SN, Huang AH, et al. Rapid identification of yeasts in positive blood cultures by a multiplex PCR method. *J Clin Microbiol* 2001;**39**:3466–71.

Take advantage of BMJ Journals' remarkable catalogue of titles with Related Collections

No busy professional has time to browse through all pertinent journals to find relevant articles, but with Related Collections you no longer have to. Follow the "Related Collections" link from any article and use the "Show Collections from other Journals" to expand your search across all BMJ Journals. Or simply follow the "Browse by topic" link on the home page. By setting up your own collections and receiving email alerts every time an article is added to your chosen area, you can build up your own significant body of knowledge.