found in the pericardium. The right auricle and ventricle were distended by a large, rounded, somewhat gelatinous tumour arising from the upper part of the medial wall of the auricle (see photograph). The tumour measured about $2\frac{1}{2}$ by $1\frac{1}{2}$ in. (6.4 by 3.8 cm.) and passed through the tricuspid valve into the right ventricle. Over the surface of the tricuspid valve and endocardium of the auricle was an area of roughening on which were numerous fibrin vegetations, apparently caused by friction between the tumour surface and the endocardium. The aorta, great vessels, and coronary arteries were normal.

A considerable quantity of slightly turbid yellow fluid was present in the abdomen.

The liver was enlarged and showed gross changes of chronic venous congestion. There was no thrombus in any of the hepatic veins. The inferior vena cava was obliterated by a fairly recently formed thrombus. This appeared to have originated at a point near the orifices of the hepatic veins, and it extended throughout the entire length of the inferior vena cava and iliac veins. The thrombus passed across the openings of the hepatic and renal veins, but did not extend into them. The portal and splenic veins contained no thrombus. No indication of the cause or origin of this thrombus other than venous stasis due to back pressure by the tumour could be detected.

Histology of the Tumour (see Special Plate)

A cellular tumour, the cells being spindle in type and showing pleomorphism and an occasional mitosis. No infiltration of surrounding tissues was noted in the sections examined. Special staining methods revealed no evidence of cross-striations to suggest a myomatous origin, but did reveal a moderate amount of intracellular collagen. The tumour therefore appeared to be of fibroblastic origin, and was considered by the pathologist to be most probably a low-grade sarcoma.

Discussion

In a review of tumours of the heart Prichard (1951) states that such tumours are rarely diagnosed before necropsy, largely because there is little knowledge of their natural history. The total number of sarcomata Prichard was able to find in the literature was 113, of which the majority were infiltrative or sessile, but some 20% were polypoidal. Whorton (1949), who also analysed the types of cardiac tumour, classified 41 as "spindle-cell neoplasms" of the type described in the present case. Hurst and Cooper (1955) state that primary sarcomata occur most frequently in the right auricle and right ventricle—a point also noted by Prichard. Of Yater's (1931) series of 48 sarcomata, 18 were in the right auricle—an extension of Mandelstamm's (1923) series of 11 primary malignant tumours in this site out of a total of 26.

There are several types of clinical picture arising from tumour of the right auricle. In a discussion of myxoma Bahnson and Newman (1953) mention four possible syndromes: (a) obstruction or regurgitation at the auriculo-ventricular valve, varying with time and the position of the patient; (b) syncope and reactional tachycardia with positional changes; (c) dyspnoea; and (d) a syndrome resembling subacute bacterial endocarditis. Weller (1940) emphasizes the symptoms of respiratory involvement and also describes two cases with superior vena caval block. Hurst and Cooper mention that inferior vena caval obstruction may be the presenting feature. Whorton also notes this and quotes four cases from the literature. In one of these (Raw, 1898), although the fibrosarcoma had infiltrated directly into the inferior vena cava causing thrombosis of the upper third, the clinical picture was rather that of dyspnoea due to haemorrhagic pleural effusion.

In the present case the original two episodes of collapse may well have been due to temporary impaction of the tumour in the A.V. ring. Subsequently intermittent obstruction of the inferior vena cava occurred, giving rise to acute hepatic congestion and pain. Eventually, as a result of this repeated obstruction, thrombosis of the inferior vena cava

occurred, producing oedema, albuminuria, and enlargement of the liver. The next stage would appear to have been impaction of the tumour in the tricuspid orifice, when the harsh to-and-fro murmur was discovered, the patient dying within three days. The fact that the liver was soft and tender in the presence of slight jaundice, normal liver-function tests, and the absence of fever might have suggested that it became congested only when the signs of inferior vena caval obstruction were noted. An acute obstruction as anatomically high as this should then lead to consideration of a tumour of the right auricle as a possible cause, particularly as collapse with a sense of impending death is so frequently a symptom of an intra-auricular mass.

In the present case, although the tumour is regarded as a fibrosarcoma, the polypoidal appearance and the absence of infiltration suggest that, if it had been diagnosed, surgical removal might have been feasible.

Summary

A case of fibrosarcoma of the right auricle in which the tumour gave rise to the clinical picture of acute obstruction of the inferior vena cava is described.

The relevant points in the literature are briefly reviewed.

It is suggested that this type of case, if diagnosed, may be amenable to surgical treatment.

I should like to thank Dr. Monica Hurst for her detailed notes on the patient's progress before entering hospital. The pathological reports were prepared by Dr. Charles St. Hill, and the slides by Mr. Elliot, senior laboratory technician. I am also grateful to Mr. R. Doyle and Dr. R. W. Brookfield for permission to publish this case.

REFERENCES

- Bahnson, H. T., and Newman, E. V. (1953). *Bull. Johns Hopk. Hosp.*, 93, 150.
Hurst, J. W., and Cooper, H. R. (1955). *Amer. Heart J.*, 50, 782.
Mandelstamm, M. (1923). *Virchows Arch. path. Anat.*, 245, 43.
Prichard, R. W. (1951). *Arch. Path. (Chicago)*, 51, 98.
Raw, N. (1898). *British Medical Journal*, 2, 1335.
Weller, G. L. (1940). *Ann. Intern. Med.*, 14, 314.
Whorton, C. M. (1949). *Cancer (N.Y.)*, 2, 245.
Yater, M. W. (1931). *Arch. Intern. Med.*, 48, 627.

ALEUKAEMIC MYELOBLASTIC LEUKAEMIA

EFFECT OF CORTISONE THERAPY

BY

R. B. THOMPSON, M.D., M.R.C.P.

Department of Medicine, Royal Victoria Infirmary and
King's College Medical School, Newcastle upon Tyne

AND

S. G. M. MACKAY, M.B., B.S.

General Practitioner

[WITH SPECIAL PLATE]

That cortisone may have an adverse effect on at least some cases of myeloblastic leukaemia is not generally accepted; the course of the illness described in this paper, however, provides evidence that such therapy may sometimes be disastrous. The case is also of interest in that it supports the suggestion that certain cases of "aleukaemic myelosis" and "myeloproliferative syndrome" bear a close relationship to frank myeloid leukaemia.

Case Report

The patient was a woman aged 51 years. Her initial complaints of lethargy, exhaustion, and shortness of breath for one month had by the time of her first clinical examination become extreme. Increasing pallor had been observed

for the same period; there was also loss of appetite and a fall in weight by over 8 lb. (3.629 kg.). Apart from increased frequency and amount of menstrual loss over the previous year there had been no history of bleeding. From childhood until the age of 32 she had been treated from time to time with iron preparations. Two years before the present illness she had probably suffered from methaemoglobinemia following sulphonamide therapy; at this time her blood count was normal. This was the only history of exposure to any potentially toxic substance. The patient's mother was known to have received regular treatment following a clinical diagnosis of pernicious anaemia. Physical examination revealed marked pallor without any hint of icterus. The liver, spleen, and lymph nodes were not enlarged; there were no skin lesions, no purpura, and no sternal tenderness. There were occasional spikes of fever up to 100° F. (37.8° C.).

The initial blood count showed: R.B.C. 2,890,000 per c.mm.; haemoglobin 7.6 g. per 100 ml. (51%); P.C.V. 22 c.cm. per 100 ml.; M.C.V. 76 c. μ ; M.C.H. 26.0 μ g.; M.C.H.C. 34.6%. The reticulocyte count averaged 0.4%. The white cell count averaged 3,000 per c.mm.; initial differential counts were neutrophil polymorphs 60%, lymphocytes 37%, monocytes 2.5%, and plasma cells 0.5%. Platelets appeared scanty on smears and counts averaged 80,000 per c.mm.; no abnormal forms were seen. Bilirubin was less than 0.1 mg., iron 152 μ g., and urea 47 mg. per 100 ml. of serum. Alkaline phosphatase was 7.8 units (Jenner and Kay). There was histamine-fast achlorhydria. Radiological appearances of chest and upper gastro-intestinal tract were normal. A very small amount of marrow was aspirated with difficulty from one iliac crest; the specimen was too poor for detailed differential counting, but erythropoiesis was clearly normoblastic and no abnormal cells were found. A rib biopsy showed normal bone; the marrow spaces, however, were largely filled by proliferating fibroblasts (Special Plate, a), and scattered among these were normal marrow cells, with a predominance of the granulocytic series; there were many reticulin fibrils (Special Plate, b). Repeated differential counts were made on the peripheral blood during the first ten weeks of observation. Occasional primitive myeloid cells were found and, more rarely still, normoblasts (see Graph). These findings were not considered surprising, in view of the fibrosis of the bone marrow. During this period the haemoglobin level was maintained by transfusion; the clinical picture did not change.

Because of the clinical findings and the diffuse fibroblastic proliferation in the marrow, together with a pancytopenia, treatment with cortisone was discussed. It was felt that such therapy might raise the haematological values and produce clinical improvement as well as possibly checking the fibroblastic proliferation. The possibility that the disorder might be related to myeloid leukaemia, and so might be made worse by cortisone therapy, was thought to be remote, and treatment was started on the 72nd day of observation. At first the effect was gratifying; the patient's clinical condition improved and she felt stronger than for some weeks. Within eight days a remarkable outpouring of primitive myeloid cells began, however, and the cortisone dosage was reduced. The initial clinical improvement was not maintained on this reduced dosage, however, and the patient began to complain of weakness and severe pain in the limbs. After much discussion the dosage of cortisone was raised, and there followed a return of well-being and disappearance of the limb pain. An even more remarkable rise in total white cell count then occurred with an increase in

production of primitive myeloid cells; indeed, the peripheral blood picture became frankly leukaemic. The spleen became palpable for the first time, the tip reaching two to three fingerbreadths below the costal margin. Cortisone was then withdrawn and a course of 6-mercaptopurine was given with a very satisfactory haematological response. The total white cell count fell and the leukaemic cells almost disappeared. Transfusion was required less frequently; indeed, the haemoglobin level began to rise spontaneously. Clinical improvement was not so satisfactory, and after a period of intense weakness the patient died suddenly on the 175th day of observation. Unfortunately, necropsy was not performed.

Discussion

There seems no doubt that the patient was suffering from an unusual variant of myeloid leukaemia. This type of case has been well discussed in recent years by Heller *et al.* (1947), Merskey (1949), Dameshek (1951), Hutt *et al.* (1953), and Black-Schaffer and Stoddard (1953). These authors have stressed the close relationships which these cases have to chronic myeloid leukaemia on the one hand and polycythaemia on the other. It has been suggested that proliferation of marrow cells is the common factor which in some cases leads to the production of excessive granulocytes and in others to excessive erythrocyte formation. Either condition may be accompanied by a thrombocytosis and by fibrosis of the marrow, or the latter may be the predominant feature. The true nature of the underlying process in cases showing mainly myelofibrosis is usually very difficult to determine. The course of the case here described would certainly support the contention that some at least of these cases are very closely allied to myeloid leukaemia. This concept of myelofibrosis excludes, of course, those cases secondary to granulomatous deposits in tuberculosis, sarcoid, and Hodgkin's disease, secondary to carcinomatous deposits, and resulting from various chemical and physical agents. The case described here differed from most published cases of "myeloproliferative disorder" or "aleukaemic myelosis" in having no splenomegaly until the terminal leukaemic phase. A similar case (Case 2) was recorded in the series reported by Hutt *et al.*

Graph showing haematological findings and treatment.

(1953); here, however, the fibrosis was more pronounced and there was no leukaemic phase comparable to that seen in the present instance.

We have been unable to find a record of any similar case treated with steroids. Gray and Taylor (1953) have, however, reported a patient with an acute systemic reticuloendotheliosis who developed a frank monocytic leukaemia while on A.C.T.H. therapy. That steroid therapy might be harmful in at least some cases of acute myeloblastic leukaemia, in contrast to the remarkable though temporary remissions often obtained in lymphoblastic leukaemia, has been suggested by several authors (Rosenthal *et al.*, 1951; Fessas *et al.*, 1954; Björkman, 1956). Very recently Joseph and Levin (1956) have described a very interesting case of monocytic leukaemia in which cortisone made the leukaemia worse. One of us (R.B.T.) has further unpublished evidence pointing to the same conclusion. On the other hand, steroid therapy is recommended for both lymphoblastic and myeloblastic leukaemias by Scott (1955), and by Hayhoe and Whitby (1955), although it is clear that the responses in myeloblastic leukaemia are poorer. It seems likely, therefore, that not all myeloblastic cases show acceleration of the leukaemic process under the influence of steroids. At present there is no method of distinguishing one type of case from the other except by clinical trial of steroids. It is, admittedly, often difficult to distinguish acute lymphoblastic from acute granulocytic leukaemia. With care, however, such a distinction is usually possible with the exception of occasional very acute cases showing no evidence of differentiation. In these a diagnosis of stem-cell leukaemia must be made; most of them seem to react as lymphoblastic leukaemia. In view of the disastrous effect of steroids in the case described above it does seem important to make as accurate a diagnosis of the type of leukaemia as possible and to bear in mind the possibility that an adverse effect may be obtained in the myeloblastic type.

Summary

An unusual case of aleukaemic myeloid leukaemia with fibroblastic proliferation in the marrow is presented which, on treatment with cortisone, became frankly leukaemic and ended fatally in spite of a haematological remission with 6-mercaptopurine therapy.

Acknowledgments are due to the Departments of Pathology and of Photography, King's College, Newcastle upon Tyne, for sections and photographs respectively.

REFERENCES

- Björkman, S. E. (1956). *Acta haemat. (Basel)*, 15, 175.
 Black-Schaffer, B., and Stoddard, L. D. (1953). *Amer. J. Path.*, 29, 413.
 Dameshek, W. (1951). *Blood*, 6, 372.
 Fessas, P., Wintrobe, M. M., Thompson, R. B., and Cartwright, G. E. (1954). *A.M.A. Arch. intern. Med.*, 94, 384.
 Gray, J. D., and Taylor, S. (1953). *Cancer*, 6, 333.
 Hayhoe, F. G. J., and Whitby, L. (1955). *Brit. J. Haematol.*, 1, 1.
 Heller, E. L., Lewisohn, M. G., and Palin, W. E. (1947). *Amer. J. Path.*, 23, 327.
 Hutt, M. S. R., Pinniger, J. L., and Wetherley-Mein, G. (1953). *Blood*, 8, 295.
 Joseph, M. C., and Levin, S. E. (1956). *British Medical Journal*, 1, 1328.
 Merskey, C. (1949). *Arch. intern. Med.*, 84, 277.
 Rosenthal, M. C., Saunders, R. H., Schwartz, L. J., Zannos, L., Satiago, E. P., and Dameshek, W. (1951). *Blood*, 6, 804.
 Scott, R. B. (1955). *British Medical Journal*, 2, 75.

The Central Council for Health Education has published a report of a conference held on January 24 at B.M.A. House on the "Education of the Public regarding Cancer." The report (price 1s.) includes sections on "Prevention and Treatment of Cancer," "Problems of Cancer Education," and "Cancer Education in Practice." Other recent publications of the Central Council for Health Education include "Look Inside," an illustrated leaflet for children on the care of teeth (price 2s. 6d. per dozen); "Safe Storage," a guide to food storage without refrigeration (price 2s. per dozen); "Poliomyelitis" (price 1s. per dozen); "Weaning and Mixed Feeding" (price 6s. per dozen). These publications may be obtained from the Central Council for Health Education, Tavistock House North, Tavistock Square, London, W.C.1.

BLEEDING HAEMANGIOMATOUS HAMARTOMA OF THE SMALL BOWEL

BY

A. F. ROBINSON, B.Sc., M.B., F.R.C.S.
Late Senior Surgical Registrar

I. A. SHORT, M.B., F.R.F.P.S., M.R.C.P.
Senior Medical Registrar

AND

J. P. SMITH, M.D.
Lecturer in Pathology
Manchester Royal Infirmary

[WITH SPECIAL PLATE]

Haemangioma of the gastro-intestinal tract is a very uncommon condition, rarely diagnosed without laparotomy. Like Guy and Cooper (1950), we believe that this warrants the reporting of all cases. The case we now describe is instructive in that the diagnosis was not suspected before operation, and the result of exploratory laparotomy and excision of the lesion was dramatically favourable.

Case Report

A man aged 46 was admitted to hospital in the early hours of April 29, 1955. On the previous evening, while driving his car, he suddenly felt faint and had a desire to defaecate. He managed to drive home—a few miles—and then passed a stool mixed with bright red blood. Six days previously he had had a brief spell of nausea, vomiting, and diarrhoea which he attributed to something he had eaten, but he had felt perfectly well on the day of onset of the rectal bleeding. He had always been a healthy man, though five years previously there had been a little rectal bleeding, said to be due to haemorrhoids. There had been no weight loss and no dyspepsia.

On admission he was not shocked, though his own doctor had found him to be in a state of shock about an hour before. Abdominal and rectal examination revealed no abnormality except the presence of fresh blood in the rectum. Next morning he was passing dark fluid blood; his haemoglobin was 74%. There was no evidence of any abnormal bleeding tendency.

Over the next seven days three attacks of melaena associated with shock occurred, and on each occasion blood transfusion was required. During this week the haemoglobin level was never above 60% in spite of the infusion of a total of 6 pints (3.4 l.) of whole blood and 2½ pints (1.4 l.) of packed red cells. On May 2, after the second attack of bleeding, he was seen by one of us (A. F. R.), and it was decided that any further bleeding would necessitate urgent operation. Thereafter his progress was observed jointly by the medical and surgical units concerned. Operation was finally performed on May 6 after the third attack of bleeding, which, like the others, led to a rapid deterioration in the patient's general condition. At no time had there been abdominal tenderness or any sign of a local lesion in the abdomen, and a definite pre-operative diagnosis was not made. The following possibilities were considered (in order of likelihood): ulcerated Meckel's diverticulum; proximal colonic neoplasm; duodenal ulceration.

Exploration through a right paramedian incision centred opposite the umbilicus revealed distension of the lowest 6 ft. (180 cm.) of the ileum and of most of the proximal colon by an accumulation of blood. Just below the upper limit of the distended length of ileum there was a dimpling of the antimesenteric border with an underlying hard,

M. G. P. STOKER: LATENT INFECTIONS WITH VIRUSES AND RICKETTSIAE

FIG. 1.—Culture of uninfected human carcinoma cells (strain HeLa). All preparations $\times 320$, stained haematoxylin.

FIG. 2.—Cytopathic change five days after infection with herpes virus. Note giant cell and cytoplasmic processes between cells.

FIG. 3.—Cytopathic change five days after infection with herpes virus; herpes antibody added to medium two hours after virus. Focal distribution of affected cells surrounded by apparently normal cells.

A. M. DESMOND AND B. F. SWYNNERTON: ADULT HYPERTROPHY OF PYLORUS

FIG. 1.—Elongation and narrowing of pyloric canal with surrounding soft-tissue shadow and duodenal bulbar impression. (Case 18.)

A. F. ROBINSON *ET AL.*:
HAMARTOMA OF BOWEL

FIG. 2.—Elongation and narrowing of pyloric canal with proximal and distal impressions. (Case 14.)

Photomicrograph of tumour. ($\times 5$.)

R. B. THOMPSON AND S. G. M. MacKAY: ALEUKAEMIC MYELOBLASTIC LEUKAEMIA

Sections of ribs to demonstrate (a) fibroblastic proliferation; (b) reticulin fibrils. ($\times 225$.)