BRITISH MEDICAL JOURNAL

LONDON SATURDAY MAY 4 1957

SURGICAL TREATMENT OF AORTIC STENOSIS

BY

Sir RUSSELL BROCK, M.S., F.R.C.S.

Thoracic Surgeon to Guy's Hospital; Surgeon to the Brompton Hospital

The surgical treatment of aortic stenosis has received little attention in the literature of this country, in spite of numerous articles from the United States. Logan and Turner (1954) reported on seven patients treated by operation, and Marquis and Logan (1955) described six cases of congenital aortic stenosis treated by operation, but were unenthusiastic about their results. More recently Baker and Campbell (1956) have presented the results in 16 of their patients operated upon at Guy's Hospital.

This article gives my views and experiences in operating on 120 cases of aortic stenosis; it necessarily touches on the clinical features of this disease, but is intended primarily as a surgical essay.

Historical

The history of the surgical treatment of aortic stenosis is inevitably short. It is usually stated that the first operation was done in 1913 by Tuffier, who attempted to dilate the stenosis by invaginating the aortic wall with his finger. This can scarcely be accepted as a serious contribution.

My own interest in valvotomy focused upon the aortic valve soon after the war, when, in common with a number of surgeons throughout the world, I was engaged in seeking ways to develop direct operations upon the heart valves as the next obvious and long-overdue surgical development.

Souttar (1925) had pointed the way to a simple technique to deal with mitral stenosis, but it proved difficult in 1946 to persuade cardiologists that herein lay a promising (and as it turned out, brilliant) therapeutic future; there was too much inborn reluctance to touch the heart. However, it seemed that aortic stenosis offered to the surgeon a route which avoided direct interference with the heart and therefore was less likely to evoke the superstition. The important fact seemed to be that the aortic valve could be approached with relative simplicity from above. The anatomical basis of the arterial route to the aortic valve and to the pulmonary valve has been described elsewhere (Brock, 1950); it suffices to say that an instrument introduced into the innominate artery either via its subclavian or common carotid branch, or directly into its trunk, can be passed straight down to the aortic valve cusps.

During 1946 I designed an operating cardioscope, or combined valvoscope and valvotome, and was able to use it on a case of aortic stenosis on March 27, 1947. The right subclavian artery was exposed through a small

incision in the neck, and the cardioscope was passed without difficulty to the aortic valve; a good view was obtained of the irregular calcified masses which distorted the diseased valve. It was realized that an attempt to cut this calcified valve semi-blindly with the small knife attached to the instrument was unlikely to be successful enough to justify the risk involved. The instrument was withdrawn; the patient suffered no harm nor derived any benefit.

About this time also Smithy (Smithy and Parker, 1947) was grappling with the problem until his own untimely death from the disease itself. Bailey *et al.* (1952) next reported that the technique of passing a cutting valvotome through the ventricular wall and then using an expanding dilator to split the valve open, although so satisfactory in pulmonary valve stenosis, was liable to cause severe regurgitation if used on the aortic valve. They suggested that instead of preliminary use of a valvotome the stenosis should be relieved by means of a powerful expanding dilator, and to this end designed a three-bladed one, based on the two-bladed dilator already used with success on pulmonary stenosis (Brock and Campbell, 1950). There is no doubt that Bailey's recommendation of the basic use of an expanding dilator was sound. In ordinary calcified aortic valvar stenosis the architecture of the fused cusps is so distorted that the blind use of a cutting instrument is dangerous and ineffective; dangerous because it is easy to miss the valve lumen and perforate the aortic wall or the tissue of a cusp; ineffective because it may encounter only calcified tissue it cannot cut. A properly designed expanding dilator finds its way through the valve lumen with less danger and can disrupt the fused cusps. When the valve is not calcified, as in children or young adults, there can still be a place for the preliminary use of a cutting valvotome before the valve is cautiously split open with a dilator, unless of course open valvotomy is used.

Bailey was soon able to report a large experience with this type of transventricular valvotomy (Bailey *et al.*, 1954), but after operating on some 65 patients he rejected the transventricular route for a transaortic one (Uricchio *et al.*, 1955; Bailey *et al.*, 1956). Harken (1955) has also accepted the aortic route to the valve, and, indeed, has stated dogmatically that it is the only route that should even be considered. A number of other American writers have also made the same recommendation, among them Fell (1955) and Likoff *et al.* (1955). In Great Britain, Logan has used the transventricular route (Logan and Turner, 1954).

After my initial case of retrograde approach to the valve in 1947 I used this route in two more patients. Instead of a small neck incision the upper part of the sternum was split and the innominate artery itself used for the passage of instruments. My experience with it was so bad that I abandoned it in 1952, with reluctance, for the transventricular route. After some initial troubles this route has proved so satisfactory for general use that I have continued to use it for most cases up to date. I cannot agree with the arguments put forward for the superiority of the transaortic route over the transventricular; the matter is discussed more fully below.

The use of the blind method in valvotomy, whatever the route, has much against it. Although one can feel content that in relieving aortic stenosis by dilatation something is being achieved where nothing was being done before, it is not possible to be really satisfied with it. Especially is this so in children or young adults in whom the valve is not calcified. Here it would seem essential for good surgery to achieve as near a perfect result as possible with the least chance of causing regurgitation. There is only one way to achieve this in a high proportion of cases, and that is by open direct-vision valvotomy. This is, of course, in keeping with the present inexorable trend of cardiac surgical technique, towards open-vision operating.

Open aortic valvotomy has been inevitable, and it is interesting and significant that the step was undertaken by two or three surgeons about the same time. My own first open aortic valvotomy was done on January 26, 1956 (Brock, 1956), and I have since done six cases. Lewis *et al.* (1956) have reported three cases of open aortic valvotomy under hypothermia, the first being operated upon on December 2, 1955. Lillehei *et al.* (1956) report a single case of open aortic valvotomy for calcified aortic stenosis using a complete by-pass and retrograde coronary perfusion; this operation was done on January 31, 1956, and since then Lillehei has done several more. In the discussion following Lewis's communication Swan mentions having used open aortic valvotomy twice (presumably under hypothermia) and Bailey also twice, using a pump oxygenator. It is as yet rather early to assess the true value of open aortic valvotomy, but it is certainly the most promising and logical method of treatment so far available.

Assessment of the results of closed valvotomy is easier and will be presented later.

Therapeutic Problem of Aortic Stenosis

As with many other heart diseases, there has been much loose and immature thought about aortic stenosis. Typical is the attitude that it is a relatively benign condition based on the observation that patients are encountered in the 50's or 60's with a short history. From this it is argued that all patients can live to 50 or 60! The same attitude was seen a few years ago with mitral stenosis: one patient aged 60 or even 70 can be so prominent in a clinician's mind that he may fix his attentions on him and conclude that any sufferer from valve stenosis can live so long. The countless thousands suffering grave or total disability or dying

Age Groups of 78 Patients With Aortic Stenosis

1-9	10-19	20-29	30-39	40-49	50-59	60-69
4	7	5	15	25	20	2

before 45 are ignored. The accompanying Table gives the age groups of the patients in my series; it should be realized that each and every one was referred for treatment because of severe symptoms. It is seen that 56 patients were below the age of 50 years and 31 below 40.

Aortic stenosis is a surgical disease and each case should be considered as possibly needing surgery; not, as is now usual, being accorded regular observation until some sinister or subterminal symptoms appear. This statement is bound to provoke protest and so must be justified. For this it is imperative to realize that one of the basic functions of the surgeon is to relieve obstruction to a hollow viscus or duct—a function which has become elaborated chiefly since the advent of modern surgery. The relief of obstruction to the oesophagus, the pylorus, the intestine, the urinary or biliary apparatus is accepted, without question, as a surgical matter. Happily, in recent years this acceptance is extending to the mitral valve and the pulmonary outflow tract, but its significance in relation to the aortic outflow is less appreciated. Neglect of, or a *laissez faire* attitude towards, chronic or subacute obstruction of the bladder or colon would be considered professional incompetence or neglect, and yet chronic or subacute obstruction of the left ventricle is still viewed with complacency.

The left ventricle is the most important muscular structure in the body; it is possible to live, and live well, without a bladder or a colon, but not without a left ventricle. To some it may seem wrong to think of the colon, the bladder, and the heart in the same terms. The heart still carries with it the atmosphere of nobility and an exalted place in the human economy which is exemplified in its being credited with the seat of the emotions. Noble it is in the same way as that other friend of man, the horse. The left ventricle in particular is the willing horse; it will go on, week after week, month after month, year after year, doing its job manfully under the severest burden and will make but little complaint. When it does at last begin to complain, when symptoms are insistent and can no longer be ignored, it is dangerously near its end. It will live but a few months or a year or two—once the features of failure have appeared; it is liable to stop suddenly, just as the horse may go on until it drops down dead. Even at the late stage of impending failure it may be restored to many years of useful work if its burden is relieved, if the aortic stenosis is corrected. It behoves us to recognize and respect left ventricular obstruction and to relieve it much earlier than is now done.

It is, of course, realized that until very recently there was no safe or satisfactory surgical treatment for aortic stenosis, and thus nothing was to be gained by reference to a surgeon. The object of this article is to try to show that surgery now offers a great deal to the sufferer from aortic stenosis, but that to do his best the surgeon should be consulted before signs of chronic invalidism and impending death have appeared; that only in this way can good results be obtained and confidence be engendered.

Clinical Assessment

This brings us to another fact that has to be discussed—the need for a more scientific assessment of the clinical state of the patient than is generally used. The physician may defend his policy and practice of not referring his patients for surgical intervention because he has concluded that their disability or disease does not warrant it; that the aortic stenosis is not severe enough. Without doubt there are many cases in which this is the correct and proper decision. The point to be made is that in most cases this decision is empiric; it is in the nature of a “hunch”; a clinical hunch that may be right, and may be totally wrong. In such a responsible matter more than clinical sense is needed, for to-day methods are available to help towards a more precise assessment.

The simplest, and unfortunately the most ingenuous, assessment used is the patient's symptoms. There is, for instance, the astonishing statement in an important textbook (Lewis, 1933), “there are no symptoms of aortic stenosis.” The most that can be said of this is that it is presumably meant to be subtle and it merely succeeds in being sinister. If the patient says he is not troubled or is conscious of only slight restriction on exertion, this is usually accepted as ruling out the need for operation. Clearly this may be so,

but we do not receive long training to become doctors, and longer still to become specialists, to apply a simple rule of thumb that could be applied by a layman. It is our duty to assess the situation in each patient so that he can be given sound and sober advice. This may well be reassurance and observation, but we must be able to recognize when the circumstances are not satisfactory even though they may appear to be so superficially. The left ventricle, in this matter, is its own worst enemy; it will go on doing its job with but little outward protest until it can continue no longer without grave complaint. What we need is to be able to assess the burden the left ventricle is carrying even though it carries this burden without noticeable complaint.

The first step in this is clinical assessment. Others have written of this—for example, Matthews *et al.* (1955) and Baker and Campbell (1956)—and so only a few comments are needed here, and it is best to start at the end. When the heart has grossly failed the patient may be too late for surgery and certainly too late for more than moderate amelioration. Yet how often is this the very time that the patient is first directed to the surgeon even though he may have been regularly observed for years. It is like guarding closely the sufferer from chronic prostatic obstruction until his bladder is up to his umbilicus, his urinary tract infected, and his kidneys barely functioning, and only then referring him for operation. Features of right-sided heart failure are usually features of inoperability, not indications for operation; established right-sided failure indicates a virtually totally inoperable state. Much the same applies to left-sided failure. The condition unrelieved or unrelievable by routine medical treatment indicates that the time for useful surgery has passed. It is useless to appeal that only an operation gives the patient a chance of survival; this is not surgery, it is the prostitution of surgery. Surgery is a valuable tool which should be used properly and with respect; it becomes damaged and blunted if misused. The bad results of surgery repeatedly invoked in such cases promote lack of confidence, impair morale, and hold up progress.

If left-sided failure has occurred but has responded satisfactorily to treatment, especially if it was precipitated by an acute infection or some such episode, then successful operation may still be possible. The same applies to the early features of left ventricular failure, such as a tendency to nocturnal dyspnoea; it is still not too late to save these patients. It must, however, be realized that these are not so much indications for operation as indications that if operation is not undertaken as an urgency it will be too late.

Angina pectoris is often considered as a contraindication to operation; this is not so. It should be regarded as a *cri du cœur* for relief. Anginal symptoms are usually completely relieved by a successful valvotomy.

Dyspnoea at night is a grave feature; dyspnoea on exertion is often brushed aside. This should not be so; it should always, however slight, be accepted as a significant symptom until full assessment has shown it not to be. Although it is a very difficult feature to assess, and though it may prove unimportant, nevertheless, if genuine, it indicates that under the conditions in which it occurs the heart is unable to do its work properly, its ability is limited, it is failing in its task. In other words, dyspnoea can be an important feature, an important warning of the need for help.

Except in emotion or on physical effort we are unconscious of the heart's action. In a patient with aortic stenosis simple awareness of the heart's action may be a feature that should lead to complete and critical assessment of the burden the left ventricle is carrying. Especially is this so in the child or young adult, in whom the left ventricle has a remarkable power of compensation for the demands placed upon it; a power that we should recognize, respect, and seek to relieve from burden rather than take advantage of to the extent of abuse.

How can this assessment be made in the absence of the obvious clinical stigmata of failure? Simple enlargement

of the ventricle may be only reasonable compensation. If considerable it indicates a considerable burden; in the absence of regurgitation much enlargement suggests a degree of actual failure. The electrocardiogram is indispensable in assessment, especially if over a period of observation it shows deterioration. The clinical examination of the pulse is also invaluable, particularly when extended by the more critical analysis of an arterial pulse tracing.

In many cases it is quite easy from the above findings to decide that operation is desirable or essential. In others more critical assessment is needed.

More Critical Assessment

Right-heart catheterization by itself can give little direct help; it may demonstrate a low cardiac output and poor response to exercise. The "pulmonary capillary venous pressure" may also indicate or confirm evidence of left ventricular limitation or failure. The information obtained is likely to be more useful if combined with a knowledge of the pressure in the left ventricle and aorta. This may be obtained by direct left atrial puncture according to the technique of Björk *et al.* (1953), but we have found that, although a left atrial pressure record can be secured, it is, disappointingly, frequently impossible to manipulate a catheter into the left ventricle in cases of aortic stenosis; it is much easier when the left atrium is dilated as in mitral valve disease. Moreover, left atrial puncture may be very distressing to the patient, and, if he is ill with aortic stenosis, may be frankly dangerous. However, we have been able to obtain valuable information in this way by simultaneous recording of the left ventricular pressure and the brachial or femoral artery pressure, thus obtaining the gradient across the valve.

In view of the uncertainty and drawbacks to left atrial puncture we have been using direct left ventricular puncture with greater satisfaction. This method is quick and simple, and so far has proved safe (Brock, Milstein, and Ross, 1956). It has been invaluable in deciding for or against operation in difficult cases. Thus, although clinical and other features may indicate the probable need for operation, direct left ventricular puncture has shown unequivocally a trivial gradient across the aortic valve; conversely, it has demonstrated in not dissimilar cases a very high left ventricular pressure and a very high gradient, indicating the urgent need for operation. In the 90 cases in which we have used this method it has been incomparably less disturbing than left atrial puncture; there has been no morbidity and no mortality. A more detailed report on the analysis of the information obtained from left ventricular puncture is given by Fleming and Gibson (1957), and I need not anticipate their observations.

Figs. 1 and 2 show the type of pressure record and information obtainable from the direct left ventricular puncture combined with a brachial artery pressure tracing. Fig. 2 illustrates that a direct left ventricular pressure record may exclude significant aortic stenosis.

The only time the gradient may be deceptively high is if significant regurgitation coexists; it should be possible to recognize such substantial regurgitation by other methods.

FIG. 1.—Left ventricular and brachial artery pressure records showing an aortic systolic gradient of 120 mm. Hg, thus demonstrating severe aortic stenosis. Left ventricular pressure recorded by direct percutaneous puncture of ventricle.

FIG. 2.—Brachial artery and left ventricular pressure records showing an aortic systolic gradient varying from 5 to 15 mm. Hg, thus excluding significant aortic stenosis. Left ventricular pressure recorded by direct percutaneous puncture of ventricle.

Usually the pulse pressure is too wide for dominant aortic stenosis and the actual gradient is rarely as high as 100 mm. Hg and more usually nearer 50 mm. Hg. Incidentally there is a tendency to be too ready to reject a patient for aortic valvotomy because of the presence of *some* regurgitation; a small amount may be quite unimportant and should not turn one away from relieving the dominant stenosis; in these cases study of the aortic gradient may be especially helpful. Valvotomy may even relieve the regurgitation as well as the stenosis if it restores greater mobility to the cusps.

In a short communication to the British Cardiac Society in 1954 I indicated the difficulty of diagnosing significant aortic stenosis from the physical signs alone in all cases; in particular the mere presence of a systolic thrill and murmur may be misleading. This was illustrated by pressure records taken at operation for aortic coarctation in a boy who had been diagnosed as also having dangerous aortic stenosis. It can be seen (Fig. 3) that there was no gradient across the valve. In another patient, observed for several years by a skilled cardiologist who requested operation as a result of his clinical observations and his knowledge of the patient's progress, pressures taken across the aortic valve at thoracotomy also showed no gradient. (Today this information would have been obtained by direct

FIG. 3.—Pressure records of aorta and left ventricle taken at thoracotomy; there is no gradient and no significant aortic stenosis.

left ventricular puncture.) These two cases emphasized that it is not enough to accept even a skilled clinical assessment in all cases and indicate that the surgeon must check the pressures across the valve at operation, if these have not been done before. Operation results and experiences in patients in whom this has not been routinely done may be, in my submission, valueless; they are certainly incomplete.

The ability to do percutaneous left ventricular puncture and simultaneous brachial artery or aortic records of pressure allows us to obtain this precise and valuable information as a part of routine pre-operative assessment.

So far we have found a reasonably close correlation between the pressures recorded before operation and those taken at operation. This is in direct disagreement with the statements made by others that observations at operation are unreliable (Matthews *et al.*, 1955; Bailey *et al.*, 1956).

Matthews *et al.* (1955) also state that it should not be necessary to know the gradient across the valve to assess the severity of the stenosis, and that, apart from the figures being unreliable, the requisite information can be obtained by other means. They give no figures to support this, nor disclose any experience obtained from cases studied before and at operation. In my experience it is not possible to obtain sufficient precise information on these cases without left as well as right-heart catheterization. It is difficult to imagine any pre-operative clinical assessment of cases much superior to that which I am privileged to enjoy from my very able cardiological colleagues, and yet we have found that the final essential information has on several occasions come only from left ventricular puncture with or without right-heart catheterization. This has enabled us to assess the severity of the lesion and thus has greatly helped the difficult task of advising operation in an anxious case, even though there was already a strong clinical indication for operation. Also it has enabled us, unequivocally, to rule out or postpone operation in others in whom operation had seemed indicated.

At present large numbers of patients with aortic stenosis are being observed at cardiac clinics throughout the world by virtually rule-of-thumb methods. Especially is this so with children. Their aortic stenosis may be mild and unimportant; it may be severe. With present practice there is little chance of the more severe cases being selected for operative relief until they begin to show signs of grave failure, by which time it may be too late.

If these patients are to be selected for operation at the proper time, when their left ventricle needs relief from its burden and before advanced secondary changes have occurred, this cannot be done by clinical "guessing," however shrewd this may be and however hard it is to accept that it often is guessing. If proper progress is to be made in the surgery of aortic stenosis then the approach to and management of these cases have got to be reorganized. Especially must this be so in children and young adults, for it is in them that the best results will be achieved before the valve has become so severely calcified that it cannot be made even remotely normal. Nadas and Ongley (1956), of Boston, reported a careful and complete study, including left- and right-heart catheterization, they had undertaken in a series of children with aortic stenosis. They realized the need for this after observing five sudden deaths in five years. Their findings revealed the need for urgent surgery in almost all their patients.

Clearly it would be unreasonable to expect patients with aortic stenosis to have been put forward for operation, or even necessarily critically examined with a view to operation, until quite recently. The surgery of aortic stenosis had not justified itself; the mortality was high and the results were poor. Clinicians have, quite properly, been hesitant and unconvinced. One of the objects of this article is to try to show that there is now every justification for confidence in advising patients to consider surgery and to undergo it if found to be suitable for operation.

Morbid Anatomy of Aortic Stenosis

The outstanding feature in aortic stenosis is the frequency with which the valve cusps are deformed by calcified deposits. On this rock-like substance the surgeon's efforts may founder, for he is often at the mercy of the anatomical accident of the disposition of these masses; what he can do to help may be decided largely by them. In extreme cases the whole valve and ring may be so calcified that little or nothing can be achieved; in others the commissures, or at any rate part of them, may be sufficiently free from calcium to allow a satisfactory result. The least favourable type is the bicuspid valve in which it may be impossible materially to increase the valve lumen without rupturing part of the cusps and thus causing severe regurgitation. In such a valve the cusp margins are straight lines, rigid and fixed; they are not curved like bucket-handles and so possessed of the power of swinging wide open as is the case when a stenosed but prominent valve cone is cut transversely by valvotomy.

Clearly it is desirable, whenever possible, to operate before calcification has developed. Unfortunately, only too often extensive calcification is already present when the patient first comes for advice. In my series in only one patient over the age of 30 was the valve not calcified; she was 32. Once calcification is present it has to be accepted and the efficacy of surgery is inevitably limited; cure, as such, is impossible. This is, however, no reason why operation should be rejected. Theory may be against it; practice proves it can give satisfactory results, as will be shown. At the very least the clock can be turned back several years, even though considerable stenosis remains; the patient may gain a larger valve orifice than, in fact, he has had for the greater part of his life.

The non-calcified valve has been studied at open operation in six cases; in five there were three fused cusps, in one the valve was bicuspid. The cusps were much more thickened and swollen than is seen in the stenosed pulmonary valve, the commissures in particular being swollen and difficult to define on their ventricular aspect. This may well be due to the higher pressures involved.

The patient who had a bicuspid valvar stenosis also had a severe subvalvar stenosis. This condition was also found as an isolated lesion in four other patients, and the five cases have been fully reported elsewhere (Brock and Fleming, 1956). It must suffice to state here that the pre-operative differentiation between valvar and subvalvar stenosis is difficult; the various clinical and semi-clinical differential features put forward are valueless. In four cases the diagnosis was made at operation by a withdrawal pressure record taken with a catheter passed into the aorta through the left ventricle. In the fifth case a withdrawal pressure record failed to show the subvalvar stenosis which was disclosed by open operation on a supposed valvar stenosis. It is our policy to make a withdrawal pressure record whenever there is any doubt about the lesion being valvar, especially in the absence of calcification. The possibility of the stenosis being subvalvar must be borne in mind and every effort made at a precise differentiation.

When the aortic stenosis is associated with mitral stenosis severe calcification is less frequent and consequently valvotomy offers more in isolated aortic stenosis. In 34 cases of combined aortic and mitral stenosis calcification was absent in 20 and was not severe in most of the others.

Surgical Technique

As already stated, the transventricular approach is preferred, except for open valvotomy. The heart is exposed by the standard left antero-lateral thoracotomy through the bed of the fifth rib, which is used for mitral valvotomy also. Although the skin incision extends backwards only some 2 inches (5 cm.) beyond the inferior angle of the scapula, the rib bed is opened for its whole length. This gives an excellent exposure and spares the big shoulder muscles. Occasionally, when the left ventricle is very large, it is

better to divide the sixth costal cartilage as well, and occasionally it may be more convenient to use the sixth rib bed.

Once the chest is opened the next step is dissection of the aortic branches to the head and neck and the passage of snares for their later control; it is essential to do this to help guard against calcified emboli. It is best to secure the vessels early in case emergency valvotomy is needed should the heart show distress.

The aorta is next palpated and the character of this thrill and pulsation is noted. The state of the valve and the valve ring is examined, especially for calcification. Pressures are then taken in the aorta and left ventricle by simultaneous needle puncture; in this way the gradient is assessed or confirmed. A catheter withdrawal tracing is not now used as a routine, because the manipulations of introducing it may be disturbing and may even produce ventricular fibrillation; it is used when there is a possibility of a subvalvar stenosis.

A favourable site on the left ventricle is selected for entry; this should be towards the apex and in a region as free as possible from coronary branches. A local anaesthetic solution is not injected into the myocardium; it does no good and may do harm. Formerly a heavy linen thread purse-string stitch was used and held in a Rumel tourniquet, but it was found that the strong pull that an assistant can exert may cause partial ischaemic atrophy within the constricted area. A light 00 silk stitch is now preferred, held by a short piece of soft rubber tubing; this is effective and not harmful. Sometimes no stitch at all is used.

Through a small incision a probe is inserted to feel the way into the ventricle; this is followed by a 0.4-cm. bougie. Neither of these instruments goes high enough to engage the valve lest calcium emboli are detached. The snares are now tightened on the aortic branches and the valve dilatation is done without loss of time. The 0.4-cm. bougie is advanced to the valve, and palpation will quickly give some idea of the size, shape, and disposition of the stoma. This is quickly followed by the 0.7-cm. bougie, and then the expanding dilator is introduced and opened in two or three directions. All these movements must be fully controlled. If there is any doubt at all of the true position of an instrument it should be withdrawn and a fresh start made. To push or to use force may be disastrous. Only rarely is the orifice so small that a bougie is really gripped tight, and then the advancing force must be steady and controlled; all movements of the instrument should be checked and guided by fingers placed outside the aorta. As the instruments are changed the assistant applies light pressure, if needed, to the purse-string tourniquet.

An observer calls out the duration of the time of occlusion of the head and neck vessels; first at 45 seconds, then at 60 seconds, and thereafter every 10 seconds. It is usually possible to complete the valve dilatation in 50-70 seconds (Fig. 4). A pause of some 5-10 heart beats is allowed to disperse any free calcium, and the snares are then loosened. Haemorrhage is usually slight and is controlled by finger pressure in addition to light tightening of the tourniquet

FIG. 4.—A continuous brachial artery pressure record taken before, during, and after aortic valvotomy. Three features are demonstrated. (1) The fall in blood pressure to almost zero marks the duration of occlusion of the aortic branches to the head and neck to prevent calcium emboli; it lasts some 65 seconds, and during this time the valvotomy was completed. (2) The blood pressure rapidly returns to the pre-valvotomy level. (3) On the left and the right are slow tracings to show the form of the pulse wave. Before valvotomy the slow upstroke shows a classical anacrotic notch; after valvotomy the upstroke is steep and has no notch.

If bleeding is heavy a swab can be held over the wound. At this stage it is better to wait quietly a few minutes to let the heart settle down; this will always lessen the bleeding and may allow it to stop altogether. Palpation of the aorta will usually reveal changes in the thrill, pulsation, and pressure to indicate the degree of improvement. When the heart's action is steady the improvement can be checked by taking pressures across the valve once more by needle puncture. If these are satisfactory it only remains to close the small wound in the ventricle with two or three stitches. If the gradient is still high further instrumentation is needed.

It is necessary to take great care in tying the stitches. If the muscle is at all friable and they are tied too tightly they will cut out, and control may then be difficult; the sutures should be bedded down just so much and no more. Much has been written about uncontrollable haemorrhage and unsuturable wounds, but these have never been so in any of the cases in this series. It may be that the use of a large, heavy type of expanding dilator is responsible for severe damage to the myocardium. The simple two-bladed dilator of the type used for pulmonary stenosis and opening to 3.5 cm. is less than 1 cm. in diameter. It has given no

FIG. 5.—Two-bladed expanding dilator.

trouble and is very powerful and effective (Fig. 5). The suggestion that a three-bladed expanding dilator can open the three commissures is unconvincing.

The heart may show distress during the early manipulations, and ventricular fibrillation may even begin. In these circumstances it is best to proceed at once to complete the valvotomy and then begin resuscitation.

If fibrillation begins during the valve dilatation this can be completed and then massage and defibrillation carried out. Several patients have made a perfectly normal recovery even after going through a period of many minutes of ventricular fibrillation (Milstein and Brock, 1954).

When aortic and mitral stenosis coexist the heart is exposed in the same way. In most cases the mitral stenosis is the more important lesion and is dealt with first; pressures are always taken across both valves to try to elucidate this. If the aortic stenosis should be the more important it is naturally treated first. After the mitral valvotomy the aortic gradient is again checked before the aortic valve is dealt with according to the technique already described, except that the aortic branches would not be snared in the absence of calcification. Aortic valvotomy is usually well tolerated in these cases because the left ventricle has, as it were, been protected by the mitral stenosis.

The technique for open aortic valvotomy is, of course, quite different. The chest is opened by the standard transverse, transsternal incision through the bed of the right third and left fourth or fifth rib. Pressure estimations are done as usual and then venovenous cooling is instituted accord-

ing to the technique described elsewhere (Brock and Ross, 1955; Brock, 1956), the temperature being lowered to about 29° C.

After interruption of the circulation by clamping the superior and inferior venae cavae the aorta is opened to expose the valve for visual valvotomy; little more than five minutes is needed to complete this. The incision in the aorta is closed with a clamp and can then be sutured at leisure after the circulation has been re-established.

Pressure Records

A certain amount has been said about pressure records across the aortic valve, but this is a convenient moment at which to elaborate certain features.

The criticism has already been mentioned that pressure records taken at operation are unreliable, although these statements are rarely supported by actual figures.

Pressure recordings have been made at operation on all the cases of aortic stenosis in this series and on at least another 50 cases in which it has been necessary to assess the gradient critically (usually when mitral stenosis coexists). Over a period of years, and from this not inconsiderable

FIG. 6.—Pressure records from left ventricle and aorta taken at operation before and after aortic valvotomy. The pre-operative gradient of 90 mm. Hg has been reduced to 10 mm. Hg; the systemic pressure has actually risen 10 mm. Hg. The form of the aortic pulse tracing has improved.

number of cases, the impression has been formed that these observations cannot be dismissed as unreliable or misleading. It is obvious that the conditions at operation may be different from those existing at other times, and proper allowance can be made for this. Thus the heart's action may be poor, the systemic pressure low, and the gradient correspondingly reduced. Clearly this must be taken into account after the manipulations of aortic valvotomy, when the ventricular pressure might be lower from the insult the heart has received. It is not difficult to recognize this state and either wait until the condition has improved or just note that the observations are not fully reliable under the conditions pertaining.

It has already been stated that our experience with direct left ventricular puncture has shown that pressure records made before operation can be exactly confirmed at operation. Furthermore, far from the heart being embarrassed by valvotomy, its action is often immediately and strikingly improved so that an even higher systemic pressure is recorded (Fig. 6). If this is so, or the systemic pressure is at any rate not lowered, it can be inferred that the drop in left ventricular pressure is the expression of relief of stenosis and not ventricular failure. Thus typical records are:

	Before valvotomy	After valvotomy
Aorta	100/56 mm. Hg	100/50 mm. Hg
Left ventricle	260/40 mm. Hg	140/40 mm. Hg

The aortic and brachial artery tracings will also usually confirm the efficacy of the procedure (Figs. 4 and 7).

If pressure records made at operation are to be rejected as unreliable it would be good to know what other method

of assessment is to take their place. Even with open aortic valvotomy there is the problem of secondary subvalvar stenosis.

The pressure gradient in isolated aortic stenosis coming to operation is usually much higher than in aortic stenosis combined with mitral stenosis. Thus the usual gradient is

FIG. 7.—Brachial artery tracings taken before and after aortic valvotomy to show conversion from a grossly anacrotic to an almost normal tracing.

about 100 mm. Hg—for example, aorta 100 mm. Hg, left ventricle 200 mm. Hg. In severe cases the left ventricular pressure may be nearly 300 mm. Hg. In one patient it was found to be 325 mm. Hg when measured by direct left ventricular puncture; the brachial artery pressure was 115 mm. Hg.

It is difficult to lay down any firm rule for a figure below which one would probably not advise operation in pure aortic stenosis; 50 mm. Hg suggests itself, especially if this were a figure arrived at by left ventricular puncture before operation. If found at operation full weight would have to be given to the evidence of any general circulatory depression, particularly an unduly low aortic pressure. Normally, also, all the other clinical features would have to be taken into account, including cardiac output.

It is expecting too much to be able to obliterate the gradient completely in a case of severe calcified aortic stenosis. Occasionally this can be nearly achieved, as is shown by Fig. 6, in which there has been a reduction from 90 mm. Hg to 10 mm. Hg. Fig. 8 also shows almost complete obliteration of the gradient.

FIG. 8.—Withdrawal pressure records made at operation for aortic stenosis. Valvotomy has achieved virtually complete obliteration of the aortic valve gradient.

Often one must be content with a residual gradient of 30–50 mm. Hg. Occasionally it is not possible to achieve any satisfactory reduction of the gradient. In such cases, however, it would be wrong to assume always that there has been no specific relief of the actual valve stenosis. For instance, in a recent case the pressures before aortic valvotomy were: aorta, 100 mm. Hg; left ventricle, 200 mm. Hg; gradient, 100 mm. Hg.

The valve orifice was felt to split well and the dilator could be opened freely. In spite of this the pressures were: aorta, 80 mm. Hg; left ventricle, 150 mm. Hg; gradient, 70 mm. Hg. The dilator was again passed, and it was confirmed that the valve opened easily and widely; it seems certain that residual valve stenosis was slight.

The probable explanation lies in the appearance of a secondary subvalvar muscular obstruction due to gross left ventricular hypertrophy analogous to what has been fully described as occurring in the right ventricle in certain cases of pulmonary valvar stenosis (Brock, 1956). A catheter withdrawal record seemed to give support to this. It can be reasonably expected, if this is the explanation, that improvement may accompany diminution in the gross muscular hypertrophy.

The value of pressure records made before, during, and after operation may be judged from the following notes of a case.

A youth aged 16 had congenital aortic stenosis; the electrocardiogram showed severe left ventricular strain pattern. Right heart catheterization and direct left ventricular puncture gave the following data. *Cardiac output*: At rest, 5 litres; during exercise, 7.1 litres; during L.V. puncture, 6.3 litres. *Pressures*: Brachial artery—at rest, 110/60 mm. Hg; during L.V. puncture, 95/60 mm. Hg; left ventricle, 177/17 mm. Hg; systolic pressure gradient, 82 mm. Hg. Estimated aortic valve area, 0.8 square cm.

When the heart was exposed at operation the pressure records were: aorta, 69/52 mm. Hg; left ventricle, 136/15 mm. Hg—systolic gradient, 67 mm. Hg.

After venovenous cooling to 29.5° C. the aorta was opened to expose a stenosed valve cone with a central aperture about 0.4–0.5 cm. in diameter; the three fused commissures were cut accurately and a finger was passed into the left ventricle.

Pressure records were again taken when the temperature was 32° C. These showed: aorta, 60/42 mm. Hg; left ventricle, 73/11 mm. Hg; systolic gradient, 13 mm. Hg.

Investigations were repeated six months after operation with the following findings; figures at the pre-operative catheterization are given in parentheses. *Cardiac output*: At rest, 16.7 litres (5 litres); during L.V. puncture, 10.6 litres (6.3 litres). *Pressures*: Brachial artery—during L.V. puncture, 115/67 mm. Hg (95/60 mm. Hg); left ventricle, 135/12 mm. Hg (177/17 mm. Hg); systolic gradient, 20 mm. Hg (82 mm. Hg). Estimated aortic valve area, 2.32 square cm. (0.8 square cm.).

The brachial artery pressure pulse was now normal instead of being grossly anacrotic.

It should be noted that in addition to proof of the effectiveness of the open aortic valvotomy in improving the cardiac output and converting the pressure gradient across the valve, there is very close agreement between the pre-operation records and the pre-valvotomy records at operation; also between the post-valvotomy records at operation and those made six months after operation.

When mitral stenosis exists with aortic stenosis the gradient is usually much less than in pure aortic stenosis. My first combined aortic and mitral valvotomy was done in January, 1953, but for at least three years before this the practice was made of taking pressure records across the aortic valve in cases of mitral stenosis with features of aortic stenosis and trying to obtain some information by a post-operative follow-up. As a result of this experience it is felt that a gradient of only 25–30 mm. Hg is an indication for aortic valvotomy when there is a coexistent mitral stenosis, and of course associated with clinical features of aortic stenosis. This is based on a follow-up of patients in whom the gradient was as small as this and who seemed to have a degree of disability due to unrelieved aortic stenosis. There have, of course, been patients in whom a gradient of 25 mm. Hg was found who have shown no signs of disability. Two patients in whom the gradient was 50 mm. Hg and 70 mm. Hg were left with their aortic stenosis untouched, and both had to have secondary aortic valvotomy; both then obtained complete relief. There can be no doubt that a gradient of 50 mm. Hg in a patient with combined stenosis must not be left unrelieved.

In the presence of significant mitral stenosis the gradient must inevitably be smaller because of the diminished filling of the left ventricle and hence the restricted output. As a corollary, relief of the mitral stenosis will improve the filling and the output, and hence the gradient across the aortic valve stenosis will increase.

In these combined cases it is much easier to obliterate the pressure gradient completely (Fig. 9).

The aortic valve manipulations are so well tolerated in these patients that in any doubtful case it is now thought preferable to introduce a bougie and try to gauge the size

FIG. 9.—Pressure records from left ventricle and aorta in a case of combined aortic and mitral stenosis before and after combined aortic and mitral valvotomy. The small pre-valvotomy gradient of 18 mm. Hg has been reduced to 3 mm. Hg.

of the valve orifice, or even to insert an expanding dilator to try to assess this more closely, and, if need be, open it widely and split the valve.

There were three deaths among the 34 cases of combined aortic and mitral valvotomy, but in all three death was due to the mitral stenosis and not to the aortic valvotomy.

Results

The greater tolerance of aortic valvotomy when combined with mitral valvotomy makes it desirable to separate this group from pure aortic stenosis.

As stated above, combined aortic and mitral valvotomy was done in 34 patients with three deaths. In 12 patients the clinical result was excellent or good; it is in this group of combined aortic and mitral valvotomy that some of the very best results can be obtained. In only seven patients was the result classed as fair and in only one as poor.

Four patients have since died; in these the initial result had been judged as good in one, fair in one, poor in two.

In two cases aortic stenosis was associated with coarctation of the aorta, both lesions being severe.

There remain 84 cases of isolated aortic stenosis; from these should be deducted the first, 1947, case in which no valvotomy was done; 83 remain. Among these are five cases of subvalvar stenosis, and it would appear better not to consider them with the pure valvar stenosis group; 78 cases remain for consideration.

Among these 78, there occurred 14 deaths due to the operation—an overall mortality of 18%. It must, however, be pointed out that five of these deaths occurred in the first eight patients, all of whom were very bad risks; four were over the age of 50, one over 60; some were in advanced right- or left-sided failure.

This early experience was so disturbing that operations on aortic stenosis were nearly abandoned, but after consideration it was decided that it was wrong to select such desperate cases; thereafter the results improved vastly. It must not be supposed that only mild cases were chosen; it was simply that moribund patients were rejected.

If, for the moment, these first eight cases are excluded we find the mortality is 9 in 70 (12.8%). Actually there have been only three deaths in the last 48 cases of pure aortic stenosis; and in one of these the patient had been in left-sided failure and had grossly impaired lung function; one was a case of open aortic valvotomy under hypo-

thermia; the third patient died 10 days after operation from a heavy wound infection resistant to antibiotics. Included in these 48 are four cases of open aortic valvotomy.

In the last 63 cases of aortic valvotomy, including those with mitral stenosis, there have been only four deaths; all except one were very bad risks.

In an average-risk patient with isolated aortic stenosis the mortality from operation should be low; it is only when features of advanced secondary changes are present that the mortality rises. This means that in certain types of case the risk of operation may lie between 25 and 50%; for average cases the risk is probably no more than 5%.

Baker and Campbell (1956) made a detailed clinical survey of the results of the first 16 patients operated on at Guy's Hospital. How gravely ill were the patients in this series is seen by the fact that six died and in only five was the result good; even among these five one died after 18 months. Baker and Campbell, indicating how these patients were chosen for operation at a late stage of their disease, reveal that 9 of the 16 had left ventricular failure and three had progressed to right-sided failure as well; all had electrocardiographic signs of severe left ventricular strain. In this larger series of 78 cases 21 had left ventricular failure and a further 11 had progressed to right-sided failure as well; 68 had electrocardiographic features of severe left ventricular strain.

The clinical result in the 64 survivors in this series of 78 can be presented simply as follows:

Died since	..	..	..	..	..	..	9
Still alive	..	..	..	..	..	..	55
Excellent result	..	..	..	..	12	..	42
Good result	..	..	..	..	30	..	
Fair result	..	..	..	..	..	13	

The nine patients who have died since require more detailed analysis. In these the result had been classed as

Excellent	1	(died 13 months later of lung cancer)
Good ..	1	(died 6 months)
Fair ..	4	(died 2 years, 20 months, 15 months, 3 months)
Poor ..	3	(died 1 year, 9 months, 4 months)

Post-operative Aortic Regurgitation

A criticism of blind aortic valve dilatation has been that it is followed by a high incidence of severe regurgitation. In this series some degree of regurgitation was produced in 14 patients and was largely responsible for the death of four of these. In the others it varied from mild to moderate in degree, the result being classed as good in five and fair in five. To balance this, pre-operative regurgitation has been lessened or completely relieved in several patients. Regurgitation is a hazard, but only a small one, and certainly not large enough to weigh against operation.

Transventricular or Transaortic Approach?

The transaortic route for valvotomy is the one now most generally advocated, chiefly as a result of the strong recommendation of American surgeons. Unquestionably this must be the correct route for open aortic valvotomy, but we are concerned now with the "blind" procedure. For this I have been unable to accept that the transventricular route is inferior. These claims must now be examined.

As mentioned earlier, I first used a supravascular approach nearly 10 years ago; in two later cases the aorta was exposed by splitting the upper part of the sternum and instruments were inserted through the innominate artery. The results were so poor that this route was abandoned for the transventricular; this has since been used exclusively without reverting to the transaortic approach (except for open surgery).

The unfavourable features claimed for the transventricular route have not been manifest. These have been stated to be: undesirable interference with the left ventricle, liability to ventricular fibrillation, excessive or uncontrollable haemorrhage, inability to suture the ventricle, inability to do a satisfactory valvotomy, and higher mortality.

No one will question that it is better to avoid direct interference with the left ventricle if this can be avoided. However, one of the chief reasons why I abandoned the supra-auricular approach was that it did not seem to protect the ventricle, seeing that ventricular fibrillation occurred in two cases, proving immediately fatal in one; the second patient died a few hours later. It is true that these were bad risks and just the type in which incision in the left ventricle would have been badly tolerated; the point is that collapse was not avoided by sparing the ventricle. From reports and records this is also the experience of others. Among the 109 cases in which the ventricle has been transversed (75 cases of pure aortic stenosis; 34 cases with combined mitral stenosis) ventricular fibrillation due to this occurred in nine. Four of these patients died at operation or soon afterwards; the remaining five were successfully converted to normal rhythm. Of the nine deaths apart from these, in none can the incision in the ventricle be directly blamed.

Ventricular fibrillation occurred in two out of three supra-auricular cases in this series and in 9 out of 109 transventricular cases. In four of these, ventricular fibrillation began before the ventricle was incised, and was, in fact, a token of the serious heart condition. One great advantage of the transventricular route then becomes apparent in that the heart can be quickly massaged, immediate valvotomy can be done, and electric defibrillation achieved without delay. The transaortic operation can be a long and much more complicated procedure, a fact against using it in a very poor risk case in which a rapid, effective valvotomy is needed. There is no evidence that ventricular fibrillation is a more serious menace in the transventricular operation. Whether ventricular fibrillation occurs or not can be determined by the way the ventricle is handled: the utmost gentleness must be used; the slightest extra stimulation may precipitate distress or disaster. In addition to the surgeon, the assistants also carry a great responsibility, especially in regard to care in use of a sucker within the pericardium. This can easily cause ventricular tachycardia and fibrillation.

Although haemorrhage is claimed to be a danger in the transventricular procedure it would appear unlikely that the transaortic approach is itself free from the dangers of uncontrolled or severe bleeding, and, indeed, this is borne out from reports. Inability to control haemorrhage from the left ventricle has never occurred in this series. In a few cases with a huge left ventricle haemorrhage has been excessive, but no more. Perhaps this difference in experience lies in the use of a smaller, kinder instrument which inflicts little or no damage on the muscle; a large instrument can be very traumatic. In this also may lie the cause of the greater incidence of ventricular fibrillation complained of, and perhaps also the other objection of inability to close the wound.

Inability to suture safely has not been met with in any of our cases. When the disease is not grossly advanced there is no difficulty at all in muscle suture. When the ventricle is big and unhealthy the muscle may hold stitches badly and great care must be taken in tying sutures.

Deaths have certainly been reported from trouble with closure of the aortic incision in the supra-auricular approach; these include delayed rupture and haemorrhage and also aneurysm formation. On balance there seems little to favour the aorta over the ventricle as safer tissue for suture.

It is maintained that operation from above allows better relief of the stenosis from ability to use the finger as well as instruments. Blind transventricular dilatation is a rather crude and surgically unsatisfactory technique, but I am unconvinced that any better result can be achieved with the finger from above. Certainly the finger can feel the lesion, but the severe and vicarious calcification of the valve tissue prevents effective digital splitting; the pathological anatomy is in no way comparable to that of mitral stenosis, in which the finger can be so effective. It may well be that in occasional cases rather more can be done than from below, but

too rarely to form a significant argument in favour of a blind approach from above. A severely calcified valve is hopelessly unfavourable by any method.

The proponents of the supra-auricular approach state that the mortality is lower in their cases.

Thus Likoff *et al.* (1955) mention a mortality of 28% in 79 patients with the transventricular route and, following this, a mortality of 10% in 50 cases of the transaortic operation.

Bailey *et al.* (1956) state that since March, 1950, they have operated on 287 patients, and their mortality figures are: transventricular, 68 cases, 19 deaths (28%); combined mitral and aortic, 87 cases, 16 deaths (18.4%); transaortic, 47 cases, 7 deaths (14.7%). In my own small series the comparable figures are: transventricular, 76 cases, 12 deaths (16%); combined mitral and aortic, 34 cases, 3 deaths (9%); Transaortic, 3 cases, 2 deaths (66%).

It should, however, be noted that the cases in the three groups of Bailey *et al.* total 202, leaving 85 cases unaccounted for in their total series. It is difficult from their report to know which cases have been excluded, although by inference it is a convenient number of the earlier ones who were operated on in an earlier technical phase. If this is so it is difficult to know how exactly to appraise the results. In my own series, if the first seven cases in which technique and selection of cases was less mature are omitted the results are: transventricular 69 cases, 9 deaths (13%).

Direct comparison of mortality figures between different and relatively small series of cases is patently weak and unsatisfactory. The above comparison does, however, show that the mortality of the transventricular operation is not essentially greater than that of the transaortic.

My experience and practice is not presented as a plea for the transventricular operation; it is merely intended to show that there are no overwhelming advantages in favour of the transaortic approach such as have been claimed.

In actual fact neither blind method can carry any strong claim, because the basic morbid anatomical changes in the valve favour neither.

The answer to obtaining better technical results in aortic valvotomy must lie in the development of open direct-vision surgery. This has already been begun, and it will soon be possible to report on a large enough number of cases to enable a fair assessment of its value to be made. At present this must be awaited, and many surgeons may still prefer to continue to use the palliative procedure of transventricular dilatation of calcified aortic stenosis rather than to submit such patients routinely to the unassessed hazards of open valvotomy under hypothermia or with total heart-lung by-pass.

Summary

The historical development of the surgery of aortic stenosis is presented and also the principles governing the need for operation. It is pointed out that aortic stenosis is fundamentally a state of obstruction of the left ventricle and the same principles should be followed in its management as are already well established in obstruction of other hollow muscular viscera, such as the colon, stomach, or bladder.

Surgery can now offer sufficient relief with a relatively low mortality (10–15%) to make it thoroughly justifiable to refer patients for operation.

The importance of critical clinical assessment is emphasized, aided by the more objective methods of right and left heart catheterization. Percutaneous puncture of the left ventricle is a simple, safe, and quick way of determining the pressure gradient across the valve.

The various operative approaches are discussed. For calcified aortic stenosis the transventricular route is preferred. For non-calcified cases, open aortic valvotomy under direct vision and with hypothermia is used.

Operation was done in 130 cases, 34 being examples of combined mitral and aortic stenosis.

The mortality of operation need be no more than 10-15%. A good result was obtained in about 70% of survivors. The mortality will always be high if operation is deferred until features of heart failure have appeared.

The patients included in this study were almost all operated on in collaboration with Drs. Maurice Campbell, Paul Wood, Charles Baker, and Ronald Gibson; I am indebted to them all. I am also grateful for the invaluable help of various colleagues in the investigations before, during, and after operation. Fig. 8 is from records made by Dr. P. Fleming, and Figs. 1, 2, 3, 4, 6, and 9 are from records made by Dr. R. Gibson. I am grateful to Dr. Lawson McDonald for the arterial pulse tracing in Fig. 7.

REFERENCES

- Bailey, C. P., Bolton, H. E., Jamison, W. L., and Nichols, H. T. (1954). *Circulation*, **9**, 22.
 — Nichols, H. T., Jamison, W. L., and Litwak, R. C. (1956). *J. thorac. Surg.*, **31**, 375.
 — Ramirez, H. P. R., and Larzelere, H. B. (1952). *J. Amer. med. Ass.*, **150**, 1647.
 Baker, C. G., and Campbell, M. (1956). *Lancet*, **1**, 171.
 Björk, V. O., Malmström, G., and Uggla, L. G. (1953). *Ann. Surg.*, **138**, 718.
 Brock, R. C. (1950). *Guy's Hosp. Rep.*, **99**, 236.
 — (1956). *Proc. roy. Soc. Med.*, **49**, 347.
 — and Campbell, M. (1950). *Brit. Heart J.*, **12**, 377.
 — and Fleming, P. R. (1956). *Guy's Hosp. Rep.*, **105**, 391.
 — Milstein, B. B., and Ross, D. N. (1956). *Thorax*, **11**, 163.
 — and Ross, D. N. (1955). *Guy's Hosp. Rep.*, **104**, 99.
 Fell, E. H. (1955). In *International Symposium on Cardiovascular Surgery*. Detroit, p. 318. Philadelphia.
 Fleming, P. R., and Gibson, R. V. (1957). *Thorax*, **12**, 37.
 Harken, D. E. (1955). In *International Symposium on Cardiovascular Surgery*. Detroit, p. 315. Philadelphia.
 Lewis, F. J., Shumway, N. E., Niazi, S. A., and Benjamin, R. B. (1956). *J. thorac. Surg.*, **32**, 481.
 Lewis, T. (1933). *Diseases of the Heart*. London.
 Likoff, W., Berkowitz, D., Denton, C., Goldberg, H., and Reale, A. (1955). *J. Amer. med. Ass.*, **157**, 1367.
 Lillehei, C. W., DeWall, R. A., Gott, V. L., and Varco, R. L. (1956). *Dis. Chest*, **30**, 123.
 Logan, A., and Turner, R. (1954). *Lancet*, **1**, 1091.
 Marquis, R. M., and Logan, A. (1955). *Brit. Heart J.*, **17**, 373.
 Matthews, M. B., Medd, W. E., and Gorling, R. (1955). *British Medical Journal*, **2**, 759.
 Milstein, B. B., and Brock, R. C. (1954). *Guy's Hosp. Rep.*, **103**, 213.
 Nadas, A., and Ongley, D. A. (1956). Second European Cardiology Congress, Stockholm.
 Smithy, H. G., and Parker, E. F. (1947). *Surg. Gynec. Obstet.*, **84**, 625.
 Souttar, H. S. (1925). *British Medical Journal*, **2**, 603.
 Tuffier, T. (1913). *Trans. Int. Congr. Med.* (London, 1914), sect. vii, surgery, pt. 2, p. 249.
 Uricchio, J. F., Litwak, R., and Bailey, C. P. (1955). *Circulation*, **12**, 785.

MENTAL PERFORMANCE IN CHRONIC ANOXIA*

BY

RAYMOND GREENE, M.A., D.M., F.R.C.P.

Formerly Vice-President of the Alpine Club

No one has ever had the opportunity to observe the unalloyed effects of prolonged severe oxygen deficiency in man. In physical illness there are always impurities present: in anaemia, the cause of the anaemia; in pulmonary disease, the toxic effects of the causative organism or of the secondary invaders; even in cardiac disease, such confusing factors as the presence of cerebral arteriosclerosis. At great altitudes the complicating factors are numerous. The most important is probably the psychological make-up of the climber. A full discussion of the problem would lead the inquirer into such private fields as his subject's reason for going to the mountains at all. The Walter Mitty complex, so well described by that excellent psychologist James Thurber, is probably the common thread which runs through all our tangled motives for being mountaineers, but in many of us some of the motives are less worthy of

respect, and in consequence psychological breakdowns may occur and may have effects upon the other members of a party which may wrongly be ascribed to what on Everest in 1933 was vulgarly described as "alt."

Such considerations urge caution in attacking the question of the mental effects of chronic anoxia along the usual anecdotal lines. A description of the manic behaviour of A on Everest or the depressive interlude experienced by B on Kangchenjunga must be sceptically examined for possible psychological, as opposed to physiological, causes. Unfortunately the very proper loyalty which censors the reports of every returning mountaineer makes it impossible to obtain a sufficiently detailed description of the mental atmosphere of the party as a whole or the pattern of behaviour of individual members.

Confused though the problem is by psychological considerations and by such additional complications as wind and weather, nutrition, and technical mountaineering difficulties, we are not entirely without guidance in our search for essentials. We can look for the common factors in the conditions of chronic anoxia due to physical disease and chronic anoxia due to altitude. Bearing in mind that the clinical picture is modified by different impurities, we may justifiably assume as a working hypothesis that common factors are the result of chronic anoxia pure and simple. We can administer a further test by studying acute anoxia, whether it be anoxic, anaemic, stagnant, or histotoxic. Broadly speaking, a chronic condition produces signs and symptoms similar to those of an acute condition, modified by their being milder, more prolonged, and in consequence admixed with the syndrome produced by the adaptive reactions of the body, which, in the acute condition, have not had the necessary time to develop.

Chronic Anoxia

It is convenient first briefly to consider the effects upon the mind of chronic anoxia studied at moderate altitudes in health and at sea-level in other forms of anoxic anoxia (as in pulmonary disease); in anaemic anoxia (as in pernicious or Addisonian anaemia); and in stagnant anoxia (as in cardiac failure).

Perhaps the most reliable information bearing on our present purpose is that which has been obtained by the observation of man during prolonged residence at moderate altitudes. When I was in Darjeeling, which lies at about 7,000 feet (2,130 metres), I found a fairly general impression that those who stayed there for a year or more without a break at lower levels, often became somewhat irritable, and I have heard the same story from doctors practising in Johannesburg, which lies at a similar level. If we go twice as high, to 14,000 feet (4,270 metres), we have the observations of Barcroft (1925) at Cerro de Pasco. Mental tests disclosed nothing; and Barcroft wrote:

"Yet I am not prepared to admit that our mental capacity was the same at Cerro as it is here. All the party will agree, I believe, that the reason why the mental tests failed to reveal the difference was because our mental disability was of too subtle a character to be revealed by somewhat simple tests than because the fitness of our minds was unimpaired. It is possible to force oneself to do good work even though the test measures the quality of work done; but were it ideal it would also measure the degree of concentration brought to bear upon the task." [One is reminded of the interesting experiment conducted many years ago by some young gentlemen of a famous medical school who by constant practice trained themselves to perform perfectly, even in a state of severe alcoholic intoxication, all the tests employed by the police for drunkenness.] "Apart from the actual processes of addition, subtraction, etc., there seemed to be an unwonted difficulty in stating a sum at Cerro if it were not of a quite simple character. The Van Slyke method of gas analysis

*Read at a Symposium at the Royal Society, Burlington House, London, on January 28, 1954.