amounts in cells in a relatively isolated position argues strongly in favour of an intracellular origin rather than histiocytic phagocytosis.

Radiological Features

The radiological appearances show many features resembling rheumatoid arthritis, but differ in a number of ways: the bone is destroyed rapidly, whereas the destruction of articular cartilage proceeds more slowly; involvement of the terminal interphalangeal joints is extensive; some unusual joints, such as the posterior articulations of the lumbar spine and the costotransverse joints, are affected; some bony sites other than joints are involved—for example, the connective-tissue attachments on the femoral trochanters, ischial tuberosities, anterior angles of vertebral bodies, and the shafts of long bones such as the radius and ulna.

Relationship to Allied Conditions

Although cases similar to those here presented cannot be found in the literature of the xanthomatous disorders, certain features, notably the multiple xanthelasma palpebrarum and presence of giant cells laden with sudanophilic material, indicate a connexion. There is a histological resemblance to the giant-cell synovium of tendons and joints, and, although clinically the conditions are not clearly related, it is to be noted that five of the cases in the series developed tendon sheath tumours. Granular-cell myoblastoma also has a similar histological appearance. Such tumours are usually solitary, although a few cases of multiple myoblastomata have been described (Bloom and Ginzler, 1947; Powell, 1946). None of these has a close clinical resemblance to the cases included here, and no similar bone and joint changes are described. It is interesting that Pearse (1950) has shown that the cytoplasm of the granular-cell myoblastoma contains a lipid probably in combination with protein and sugars or amino-alcohols. Subcutaneous nodules in rheumatoid arthritis may be widespread, but never approach the dissemination and the papular appearance of the cases described here, nor is the histology similar. Fletcher (1946) reported a case of severe rheumatoid arthritis in which after 4½ years extensive subcutaneous nodules developed. These showed an appearance more like the necrobiotic nodule of rheumatoid arthritis, but numerous "foam cells" and many multinucleated giant cells were present. Presumably this case is closely related to the series under discussion.

It is suggested that the condition is due to an abnormality in connective-tissue metabolism probably akin to that which occurs in rheumatoid arthritis. However, in the series of cases now described the histochemistry of this change and the resulting histology are different.

Recently cases of this condition have been reported under the heading of reticulohistiocytoma or reticulohistiocytosis. This title does not seem adequate to describe the clinical condition or the underlying disordered connective-tissue cell biochemistry. The term "lipoid dermato-arthritis" is preferred.

Summary

Four patients having an unusual eruption and arthritis are described. Twelve cases with a similar clinical and histological picture have been recorded in the literature under various titles. The eruption consists of sheets of yellow-brown papules over the backs of the hands and fingers, the face, forearms, and other areas, being widespread in some cases. Purple nodules are also present. Xanthelasma palpebrarum and tendon-sheath swellings may occur. The joint condition resembles a severe rheumatoid arthritis, with a number of minor differences and marked bone destruction. The histology of all involved tissue shows the same type of giant cell with an abundant granular cytoplasm which probably contains a lipid-polypeptide complex.

We should like to thank the numerous clinicians, pathologists, and radiologists who have been concerned in the management

of these cases. In particular we would like to thank Dr. J. T. Ingram and Dr. E. Ritter for permission to include Case III, and Dr. J. T. Ingram to include Case IV. Professor C. B. Perry has very kindly given us helpful advice in the preparation of the paper. Our thanks are also due to Mr. J. E. Hancock for the photomicrographs.

REFERENCES

- Allington, H. V. (1950). *Arch. Derm. Syph. (Chicago)*, **62**, 452.
 Bloom, D., and Ginzler, A. M. (1947). *Ibid.*, **56**, 648.
 Caro, M. R., and Seneac, F. E. (1952). *A.M.A. Arch. Derm. Syph.*, **65**, 701.
 Davies, B. T., and Wood, S. R. (1955). *Brit. J. Derm.*, **67**, 205.
 Fletcher, E. T. D. (1946). *Ann. rheum. Dis.*, **5**, 88.
 Goltz, R. W., and Laymon, C. W. (1954). *A.M.A. Arch. Derm. Syph.*, **69**, 717.
 Graham, G., and Stansfield, A. G. (1946). *J. Path. Bact.*, **58**, 545.
 Laymon, C. W. (1952). *A.M.A. Arch. Derm. Syph.*, **66**, 647.
 Montgomery, H. (1952). *Ibid.*, **65**, 708.
 Parkes Weber, F. (1943). *Brit. J. Derm.*, **55**, 1.
 — (1944). *Ann. rheum. Dis.*, **4**, 3.
 — (1948). In McKenna, R. M. B.: *Modern Trends in Dermatology*. Butterworth, London.
 — (1955). *Lancet*, **2**, 395.
 — and Freudenthal, W. (1937). *Proc. roy. Soc. Med.*, **30**, 522.
 Pearse, A. G. E. (1950). *J. Path. Bact.*, **62**, 351.
 Portugal, H., Fialho, F., and Miliano, A. (1944). *Rev. argent. Dermatosis*, **28**, 121.
 Powell, E. B. (1946). *Arch. Path. (Chicago)*, **42**, 517.
 Targett, J. H. (1897). *Trans. path. Soc. Lond.*, **48**, 230.

EXFOLIATIVE CYTOLOGY OF THE COLON AND RECTUM

PRELIMINARY REPORT ON THE RECTAL WASHING TECHNIQUE

BY

D. J. OAKLAND, M.B., Ch.B., F.R.C.S.

(From the Surgical Research Laboratories, Queen Elizabeth Hospital, Birmingham)

[WITH SPECIAL PLATE]

Earlier reports on the cellular content of rectal discharges were made by those concerned with the diagnosis of specific inflammatory disease of the colon—that is, bacterial and amoebic dysentery (Dutcher, 1903; Jurgens, 1907; Anderson, 1921; Callender, 1925). Their interest was necessarily confined to the bacterial and parasitic content of the rectum and colon. In recent years there have been several papers in the American literature on the cytological content of the rectum and colon and the recognition of exfoliated malignant cells in cases of neoplastic disease of the large bowel. One of the problems for the cytologist working in this field is the difficulty of obtaining suitable specimens from such a large organ as the colon. A variety of techniques have been described, but none is entirely satisfactory, particularly where the lesion is situated in the proximal colon.

Previous Investigations

Bercovitz (1941) examined the cellular content of bowel discharges in a very large number of patients with normal and diseased large bowel and described the cellular flora, but did not report the presence of malignant cells in his cases of large-bowel tumour. In 1947 Wisseman *et al.* made a brief report on the diagnosis of carcinoma of the colon at various locations from the splenic flexure down to the rectum by the examination of smears prepared from the rectal contents obtained during proctoscopy. Wisseman *et al.* (1949) later reported their experiences with smears taken during sigmoidoscopy as well as proctoscopy with reference to the diagnosis of malignant epithelial tumours. The accuracy of cytological diagnosis of malignancy in the latter series was 76%, as compared with 79% by x-ray examination and 67% by sigmoidoscopic examination. Ayre (1950) described one case in which carcinoma of the rectum had been successfully diagnosed by cytology with a smear

technique. Blank and Steinberg (1951), also using the smear technique, examined a series of patients, including 45 with malignant disease of the rectum and sigmoid colon. They found malignant cells in 65% of cases when the smear was taken from the surface of the growth, but in only one out of 19 cases when it was taken from the bowel wall below the site of the lesion. They concluded that cytological examination of rectal and sigmoidal discharges had no advantage over the classic methods of investigation and diagnosis of carcinoma of the rectum and sigmoid colon.

Loeb and Scapier (1951) described a new method of obtaining specimens of exfoliated cells from the rectum and colon during sigmoidoscopy. Bader and Papanicolaou (1952), using Loeb's rectal washing apparatus, studied a series of 200 cases with symptoms suggestive of malignant disease of the lower alimentary tract. Of these cases 19 were proved to be malignant by surgery and biopsy, and 18 of these had positive or suspicious smears. Rubin *et al.* (1953) reviewed the value of cytology in gastroenterological diagnosis. They examined 33 patients with proved malignant disease of the colon by collecting the returned fluid after an enema of Ringer's solution, and found malignant cells in 27 cases. Of this same group, a radiological diagnosis of carcinoma had been established in only 18 cases.

Most recently Galambos and Klayman (1956) reported a series of 68 patients, 26 of whom had malignant lesions confirmed at operation. The correct diagnosis was established in 16 of these cases by exfoliative cytology. They concluded that colonic exfoliative cytology in experienced hands was a valuable technique in the diagnosis of colonic lesions.

Present Investigation

The purpose of this report is to describe my experience and findings with exfoliative cytology of the colon and rectum at the Surgical Research Laboratories, Queen Elizabeth Hospital, Birmingham. The 86 patients examined were composed of three groups: (A) patients with normal lower alimentary tracts; (B) patients with known carcinoma of the rectum; and (C) patients with suspected malignant disease of the colon. The rectal washing technique of Loeb and Scapier (1951) was used to obtain specimens in all cases.

Technique

In the series of patients without disease of the alimentary tract (Group A) the apparatus described by Loeb and Scapier (1951) was used, sigmoidoscopy being performed in the theatre just before or after the induction of anaesthesia. All patients in this series had an enema the day before sigmoidoscopy. Rectal washings from the patients with known carcinoma of the rectum (Group B) and suspected carcinoma of the colon (Group C) were collected by means of a modification of Loeb and Scapier's apparatus. This consisted of a similar long double-lumen brass tube, the proximal ends of which were led through and soldered to small metal disks which fitted beneath the screw-on caps of small glass specimen jars, a watertight joint being ensured by a rubber washer. Two further brass elbows were soldered to the metal disks, the atomizer bulb being attached to the one and the pump to the other. This apparatus (illustrated herewith) proved more rigid and easier to handle during sigmoidoscopy.

About 20 ml. of normal saline was used to wash the bowel wall, and an equal quantity of 90% alcohol was added to the aspirated specimen to delay autolysis of the cellular content. This fluid (together with a few drops of

Modified rectal washing apparatus.

Meyer's egg albumen) was then centrifuged at 2,000 r.p.m. for ten minutes. Smears were prepared from the deposit, fixed in an ether-alcohol mixture, and stained with the Papanicolaou stain EA 65. The smears were scanned on a moving stage of a microscope, using the high-power objective. Groups of cells of interest were marked with a diamond object-marker, and the position was further noted by reference to the Vernier scale of the moving stage.

Group A

Rectal washings were obtained from 50 patients admitted to hospital for surgical treatment of a condition unrelated to the large bowel. So far as could be ascertained by clinical examination none of them had malignant disease of the alimentary tract; however, x-ray studies were not performed to confirm this. Of the 50 cases, 34 had a non-malignant condition unrelated to the alimentary tract, such as thyroid adenoma or varicose veins; 11 had malignant disease other than in the alimentary tract, and most of these cases were of carcinoma of the breast; and a small group of five cases had chronic peptic ulcer. One washing was taken in each case either just before or just after the induction of anaesthesia.

Results.—Of the 50 smears prepared in this series, three were acellular, but the remainder contained a small variety of cells which varied in number, but were uniform in appearance throughout the series.

Squamous cells were seen in 64% of cases; they were the largest cell observed and had an average diameter of 32 μ . The nuclei were small and pyknotic in some of these cells, and in others they were larger and well defined. The cytoplasm, which was always plentiful, was occasionally laminated, and clear rings or halos were observed around some of the nuclei (Special Plate, Fig. 1).

Columnar epithelial cells and goblet-shaped mucous cells originating from the mucous membrane lining the large bowel were seen in 24% of cases. Clumps of columnar epithelial cells had a typical palisade appearance, but more often these cells were seen end on as a mosaic of regular round cells with centrally placed nucleus (Special Plate, Figs. 2 and 3).

Degenerate and unidentifiable cells were seen in 26% of cases. Many of these cells had the appearance of distended columnar epithelial cells and the nucleus was undergoing karyorrhexis (Special Plate, Fig. 4).

Macrophages were present occasionally, usually containing numerous dark-staining droplets. Polymorphonuclear leucocytes were seen in only 12% of cases, and in only one smear were they present in any number. Red blood cells were never seen, probably owing to the occurrence of haemolysis when alcohol was added to the aspirated specimen.

There appear to be three types of exfoliated cell in the normal colon and rectum: (1) Squamous cells; (2) columnar epithelial cells and mucous cells; and (3) degenerate cells of unknown origin. White blood cells and macrophages are also seen occasionally.

Group B

Rectal washings were taken from 19 patients with carcinoma of the rectum, the diagnosis being confirmed by either biopsy or operation. The object in including this series was to gain experience in the recognition of exfoliated malignant cells and compare them with the individual cells of the actual carcinoma after removal or biopsy. All the washings were taken from a point immediately below the growth.

Results.—Exfoliated malignant cells were observed in 79% of the smears. The cellular content of the smears varied considerably, and some normal cells were seen in all of them. Squamous cells formed the majority of the normal cells in this series, and polymorphonuclear leucocytes were always present but varied greatly in numbers. The exfoliated malignant cells were quite different in appearance from any cell seen in Group A. They had large,

dark-staining nuclei with thickened nuclear membranes and scanty cytoplasm which was difficult to see and stain. There was great variation in size and shape of the nuclei, which had two or more nucleoli (Special Plate, Figs. 5 and 6). The appearance was similar to that of the individual tumour cells.

Group C

Rectal washings were collected from 27 patients with signs and symptoms suggestive of malignant disease of the large bowel. In 12 cases in this series the diagnosis of carcinoma of the colon was confirmed at operation. In two further cases the diagnosis could not be established even after laparotomy and biopsy, one of these having a fixed mass in the ileo-caecal region and the other carcinomatosis peritonei. Of those 12 patients with proved carcinoma of the colon, two had carcinoma of the caecum, two carcinoma of the transverse colon, and eight carcinoma of the sigmoid colon.

Results (see Table).—The correct diagnosis was established in nine of the cases by cytological examination, and in one further case malignant cells were identified, but, although laparotomy revealed a fixed mass in the ileo-caecal

Analysis of Cytological and Radiological Findings in Suspect Cases of Carcinoma of the Colon

	No. of Cases	No. of Cases
Total suspected of having carcinoma of colon examined	27	
Diagnosis of carcinoma confirmed	12	
Correct cytological diagnosis	9 (75%)	
Site of lesion:		
Caecum	1	
Transverse colon	2	
Sigmoid	8	
False positive cytological diagnosis	1	
Unconfirmed diagnosis of carcinoma of colon	2	
Barium enema studies performed	9	
Correct radiological diagnosis	7 (78%)	
False negative radiological diagnosis	2	

region with the macroscopic features of carcinoma, biopsy did not confirm the diagnosis. In one case a false diagnosis was made, malignant cells being identified in the smears, but two barium enemas, sigmoidoscopy, and examination under an anaesthetic revealed procto-sigmoiditis with no evidence of malignant disease. In the nine cases where the cytological diagnosis was correct the lesion was in the caecum in one, splenic flexure in one, and sigmoid colon in seven cases.

Barium enema studies were carried out in nine of the 12 patients with proved carcinoma of the colon, giving a correct diagnosis in seven cases (see Table). A diagnosis of carcinoma was queried by the radiologist in three further cases, but in two of these laparotomy refuted the diagnosis, and in one, in which an ileo-caecal mass was present, the diagnosis was not proved. In three cases the results of the barium enema studies of the colon were reported as normal, but laparotomy revealed carcinoma of the sigmoid in two and carcinomatosis abdomini in the other.

In this series carcinoma of the colon was successfully diagnosed by cytology alone in one case, and the history is reported below.

Case Report.—Female aged 70 with one month's history of bleeding per rectum but no alteration in bowel habit. No abnormalities detected on clinical examination. Sigmoidoscopy showed a mild degree of granular proctitis and a small adenoma of the rectum. At 7 in. (18 cm.) from the anus there was an area of stenosis through which the sigmoidoscope could not be passed. A rectal washing was collected at this site and a specimen taken for biopsy. Two smears were prepared from the rectal washings; several groups of malignant cells were seen in both of them. The biopsy material was reported as showing chronic inflammatory changes with no evidence of malignancy. Barium enema investigation showed many diverticula in the sigmoid colon with a narrowed segment at the junction of the descending and sigmoid segments, but no characteristic features of malignancy were seen radiologically. Laparotomy revealed a carcinoma at the junction of the sigmoid and descending colons, and the pathologist reported adenocarcinoma of the colon with no invasion of the regional lymph nodes.

In this series abnormal columnar epithelial cells were frequently seen, particularly in those patients with proctitis,

colitis, or diverticulitis. The columnar epithelial cells had enlarged nuclei with an increase in chromatin content, and the amount of cytoplasm was decreased (Special Plate, Fig. 7). These changes are similar to those reported by Boddington and Truelove (1956) in ulcerative colitis. In one patient cells of this type were interpreted as malignant cells, but subsequent investigation by barium enema, contrast enema, and sigmoidoscopy under anaesthesia revealed only proctitis. This patient is being carefully followed up, but up to now, after ten months, has had no further symptoms.

Conclusions

The apparatus for collecting rectal washings was found to be satisfactory in use and good specimens were obtained by this technique. It was found that adequate cleansing of the bowel was important, since any excess of faecal material in the aspirated specimen resulted in a smear too thick to interpret microscopically. Cleansing was effected by soap enema, and purgatives were not given by mouth. Although in this investigation malignant cells were observed in one case of carcinoma of the caecum, the saline enema technique described by Rubin *et al.* (1953) and Galambos, Massey, Klayman, and Kirsner (1956) is more likely to collect cells exfoliated from lesions in the right side of the colon.

The mucosa of the large bowel is lined by columnar epithelium with many goblet-shaped mucous cells. Only two types of cell can be exfoliated from this surface which, together with the squames of uncertain origin and degenerate cells, form a constant cytological content of the normal rectum and colon. The findings in the patients with normal alimentary tracts were in accord with those of Bader and Papanicolaou (1952) and of Wisseman *et al.* (1949). However, Bercovitz (1941), examining 2,158 specimens obtained from 1,123 patients with normal alimentary tracts, found cells in only 11 cases, most of them polymorphonuclear leucocytes. He claimed that the discharges from the normal healthy bowel did not contain cells, and that a marked cellular content was one of the earliest evidences of pathological change in the lower bowel.

Carcinoma of the rectum was successfully diagnosed after one washing in 79% of cases. Three of the five false negative results occurred early in the series and were due to gross faecal contamination of the smear, making cellular identification impossible. The cytological diagnosis of malignant lesions within reach of the sigmoidoscope appears to be a practical proposition, but it is doubtful if this technique has any advantage over biopsy.

Carcinoma of the colon was diagnosed correctly by cytology in 75% of cases. Although the series is small, the results are encouraging, and a larger series of cases are being investigated by both the rectal washing and enema techniques. The one false positive case in the series draws attention to the problem discussed by Galambos and Klayman (1956) and by Boddington and Truelove (1956) of differentiating exfoliated malignant cells from cells exfoliated from an inflamed colonic mucous membrane. In the present series abnormal cells were seen in all the smears prepared from patients with inflammatory disease of the colon. Many of the cells bore some resemblance to exfoliated malignant cells in that the nuclei were enlarged and heavily stained, but they did not show wide variation in contour and the cytoplasm was usually easily recognized.

As McAmmond (1955) points out, cytological investigation requires team-work. The scanning of smears is time-consuming and is not a practical proposition for the clinician. Any cytological service for the diagnosis of malignancy of the colon and rectum would require a trained technician to prepare and scan the smears and a cytologist to give an opinion on the nature of the cells identified.

Summary

In a study of the cytology of the colon and rectum the rectal washing technique of Loeb and Scapier was used on a group of 86 patients, and a modification of

their apparatus is described. The cytological content of the normal lower alimentary tract was found to be constant and composed of three groups of cells: (1) squamous cells; (2) columnar epithelial and mucous cells; and (3) degenerate cells together with occasional white blood cells and macrophages. Exfoliated malignant cells were found in 15 of 19 patients with carcinoma of the rectum, but it is felt that this technique has no diagnostic advantages over biopsy.

Of 27 patients with suspect malignant disease of the colon, a correct diagnosis was obtained cytologically in nine, and 12 were subsequently proved to have carcinoma. The rectal washing technique appears to be most successful in the diagnosis of carcinoma of the descending and sigmoid colons.

I wish to thank Professor F. A. R. Stammers and Mr. B. N. Brooke for advice and encouragement in the preparation of this paper. My thanks are due to Mr. S. H. Sharpe, histology technician, who devised the modification of Loeb and Scapier's rectal washing apparatus and who prepared all the smears and many of the photographs. I am also indebted to Mr. J. F. Dee for some of the photographs, and Mrs. I. Denton for the typescript.

REFERENCES

- Anderson, J. (1921). *Lancet*, 2, 998.
 Ayre, J. E. (1950). *Amer. J. Surg.*, 80, 316.
 Bader, G. M., and Papanicolaou, G. N. (1952). *Cancer*, 5, 307.
 Bercovitz, Z. (1941). *Ann. Intern. Med.*, 14, 1323.
 Blank, W. A., and Steinberg, A. H. (1951). *Amer. J. Surg.*, 81, 127.
 Boddington, M. M., and Truelove, S. C. (1956). *British Medical Journal*, 1, 1318.
 Callender, G. R. (1925). *Milit. Surg.*, 56, 686.
 Dutcher, T. B. (1903). *J. Amer. med. Ass.*, 41, 480.
 Galambos, J. T., and Klayman, M. I. (1956). *Surg. Gynec. Obstet.*, 101, 673.
 — Massey, B. W., Klayman, M. I., and Kirsner, J. B. (1956). *Cancer*, 9, 152.
 Jurgens (1907). *Z. exp. Path. Ther.*, 4, 769.
 Loeb, R. A., and Scapier, J. (1951). *Amer. J. Surg.*, 81, 298.
 McAmmond, E. N. C. (1955). *Proc. roy. Soc. Med.*, 48, 919.
 Rubin, C. E., Massey, B. W., Kirsner, J. B., and Palmer, W. L., and Stonecypher, D. D. (1953). *Gastroenterology*, 25, 119.
 Wisseman, C. L., Lemon, H. M., and Lawrence, K. B. (1947). *Bull. Amer. Coll. Surg.*, 32, 243.
 — — — (1949). *Surg. Gynec. Obstet.*, 89, 24.

CONGENITAL LOBAR EMPHYSEMA

BY

D. G. COTTOM, M.C., B.M., M.R.C.P.

AND

N. A. MYERS, F.R.C.S., F.R.A.C.S.

From the Thoracic Unit, the Hospital for Sick Children,
Great Ormond Street, London

[WITH SPECIAL PLATE]

Breathlessness in babies may provide much diagnostic difficulty. Frequently it is due to respiratory infection, occasionally to cardiac failure, and rarely to other conditions, one of which is congenital emphysema of the lung. The purpose of this paper is to report six cases and to draw attention to a syndrome which, if recognized, allows of surgical correction. Details of the six cases are set out in the accompanying Table. All the patients were under the age of 4 months, all were dyspnoeic, and, in addition, four had failed to thrive. The left upper lobe was most frequently affected, and in four cases a congenital anomaly of the heart accompanied the pulmonary lesion. Other common symptoms were cough, stridor, and difficulty in feeding due to breathlessness, but cyanosis was observed in only two cases and in neither was it present continuously.

All cases were not of equal severity, and in two the condition became extremely urgent. In the first of these

pulmonary valvotomy had been performed, but 24 hours after operation the baby's condition deteriorated and a chest radiograph (Special Plate, Fig. 1) showed gross emphysema of the left upper lobe with displacement of the mediastinum to the right. A left upper lobectomy was performed, but the baby died shortly after the end of the operation. In the other case a patent ductus arteriosus was ligated in a female baby aged 7 weeks. At this operation the left upper lobe was noticed to be emphysematous, but its removal was not thought to be necessary. Four hours after operation she became extremely dyspnoeic and appeared likely to die. A second thoracotomy was undertaken urgently and the left upper lobe, now grossly emphysematous, was removed. Following this she was much better, and she subsequently made an uneventful recovery. In both of these cases emphysema of the left upper lobe was known to coexist with the congenital heart lesion, and they illustrate the fact that correction of the cardiovascular defect is insufficient to relieve breathlessness in the presence of congenital emphysema.

Clinical Features

All the babies in this series were born at full term after a normal pregnancy, and in only one case was resuscitation difficult. Tachypnoea amounting to dyspnoea is the cardinal symptom, and it is most noticeable when the baby is being fed. A dry cough is often present and stridor is sometimes a prominent feature. Cyanosis is rare and is probably an indication for urgent treatment. Mediastinal displacement is usually present, and over the involved area of lung the chest wall is prominent and relatively immobile. The breath sounds are diminished and typically the percussion note is hyperresonant, but there are no adventitious sounds. Over the rest of the lungs there may be no abnormal physical findings or there may be evidence of collapse or infection.

X-ray Findings

The diagnosis of congenital emphysema may be suspected on clinical grounds, but it can be established only by radiography. The radiograph will show an area of increased translucency with bronchovascular markings still present. There will be downward displacement of the diaphragm on the same side, and there may be a herniation of the over-distended lung across the midline (Special Plate, Fig. 2). The films should be supplemented by screening, and this will show some or all of the following features: (1) relative immobility of some part of the lung fields; (2) lack of variation in the degree of translucency of the emphysematous part; (3) mediastinal displacement with phasic respiratory activity—namely, shift to the affected side with inspiration and the reverse with expiration.

Diagnosis

The possibility of congenital emphysema of the lung should be considered when a previously well infant begins to have attacks of difficulty in breathing. It is primarily necessary to distinguish it from bronchopneumonia or cardiac failure, either of which may coexist with congenital emphysema. In neither of those conditions, although there may be cyanosis together with a raised pulse and respiratory rate, is it likely that the mediastinum will be displaced. The dyspnoea in cardiac failure may be episodic, coming on only with the exertion of feeding; there may be a murmur which directs attention to the heart; and the liver may be enlarged and tender. This last sign, while frequently indicative of cardiac failure, may also occur when the right hemidiaphragm is displaced downwards, and is therefore suggestive but not diagnostic. The presence of altered breath sounds associated with hyperresonance is found in emphysema, while in pneumonia the affected side tends to be dull to percussion.

D. J. OAKLAND: EXFOLIATIVE CYTOLOGY OF COLON AND RECTUM

FIG. 1.—Squamous cells with halos around nuclei. ($\times 840$.)

FIG. 2.—Columnar epithelial cells. ($\times 480$.)

FIG. 3.—Columnar epithelial cells viewed end on. ($\times 240$.)

FIG. 4.—Degenerate cells. ($\times 225$.)

FIG. 5.—Exfoliated malignant cells (carcinoma of rectum). ($\times 1050$.)

FIG. 6.—Exfoliated malignant cells (carcinoma of sigmoid). ($\times 300$.)

FIG. 7.—Abnormal columnar epithelial cells (case of proctitis). ($\times 300$.)

D. G. COTTOM AND N. A. MYERS: CONGENITAL LOBAR EMPHYSEMA

FIG. 1.—Congenital emphysema of left upper lobe (Case 1).

FIG. 3.—Congenital emphysema of left upper lobe associated with patent ductus arteriosus (Case 5): (A) pre-operative; (B) post-operative.

FIG. 2.—Herniation of emphysematous left upper lobe (Case 2).

FIG. 4.—Congenital emphysema of right middle lobe showing considerable tension: (A) pre-operative; (B) after right middle lobectomy.