during 1950-2, showed that 19 (50%) were improved, 15 (40%) were unimproved, and 4 (11%) died during their stay in the sanatorium. Little can be drawn in the way of conclusions from these reports. The bad prognosis given in the literature is no longer justified and should be regarded as largely of historical interest only.

In view of the high mortality previously reported the single death in the present series of 84 patients is particularly noteworthy. Mortality figures alone, however, give an incomplete picture of the effects of treatment, and it is important to know the clinical condition of those who survived and the activity of their tuberculosis. At the end of a year 70 of the patients have quiescent disease, and of the 13 with active disease improvement has occurred in 11. A total of 47 (56%) of the patients had collapse therapy or resection as part of their treatment, and these played an important part in helping to control the disease. Major thoracic surgery has proved perfectly safe in the well-controlled tuberculous diabetic.

Results at one year can only be regarded as a short-term evaluation, but the attainment of quiescent disease was closely linked to the immediate clinical and radiographic response to treatment. Even patients with features usually indicative of acute or extensive disease made striking progress. Quiescent disease was attained in the presence of such individual factors as an initial pyrexia of over 100° F. (37.8° C.) (in 77% of 13 patients), a high E.S.R. (in 78% of 27 patients), involvement of five or six zones of the lungs on a standard radiograph (in 75% of 20 patients), and even in the presence of all of these features in five patients. In these cases most of the radiological improvement occurred in the first six months with chemotherapy. It can be said that the prognosis depends on the early response to treatment, and where the clinical and radiological response in the first six months with chemotherapy is good the outlook may be regarded as favourable. Acute bilateral pulmonary tuberculosis in a diabetic is a serious complication to be regarded as a medical emergency. It calls for immediate treatment in a special hospital unit with an effective dosage of combined chemotherapy for the tuberculosis and stabilization of the diabetes as a matter of great urgency. The start of chemotherapy should not be delayed until the patient is admitted to hospital.

In conclusion, the immediate response to treatment in this group of cases as judged by the results at one year has been very satisfactory. The progress of the pulmonary tuberculosis—despite the diabetes—compares very favourably with the progress in the non-diabetic. As with the non-diabetic, however, it is necessary to follow these cases for several years before the final picture will be clear.

Summary

A one-year follow-up of 84 tuberculous diabetics is presented. Clinically most of the patients were ill. The pulmonary tuberculosis was bilateral in 48 patients, and 57 had involvement of three or more zones; 27 had acute disease, and cavitation was present in 73.

By one year there was only one death and this was non-tuberculous. Quiescence of the tuberculosis was attained in 70 patients. The average gain in weight per patient was 25.6 lb. (11.6 kg.) at a year. At least moderate clinical and radiological improvement occurred in over 90% of patients. Cavitation had either disappeared in 83% or become less evident in a further 15%. Only 5 of 83 patients still had a positive sputum on direct examination.

Rest in bed and chemotherapy, with an effective combination of anti-tuberculosis antibiotics in full dosage, was the basis of treatment for the tuberculosis in nearly all cases, and 47 had collapse therapy or resection, major surgery being carried out in 18 patients. In addition the diabetes was carefully controlled. Three diets were used, but the majority of patients had a 2,500-

calorie diet. The commonest combination of insulin used was a mixture of S.I. plus P.Z.I. in the mornings and a further dose of S.I. in the evenings.

I am grateful to numerous colleagues for kindly referring patients to me; Dr. Wallace Fox for valuable advice and criticism in the preparation of this paper; Dr. M. H. Zirk for help with the care of the patients; Dr. Jack Aspin for the radiographic assessments; Mr. S. J. MacHale for operating on these patients personally; Dr. Hector Ross for help with the pre- and post-operative care of the patients; Sister B. Crocombe and the nursing staff for their invaluable and devoted attention to the patients in the diabetic wards; Miss Joanna Tilley, the therapeutic dietitian; and Mrs. B. I. Roe for secretarial assistance.

REFERENCES

- Banyal, A. L. and Cadden, A. V. (1944). *Arch. intern. Med.*, **74**, 445.
 Central Health Services Council (1952). Annual Report.
 Ferrara, M. A. (1952). *New Engl. J. Med.*, **246**, 55.
 Fitz, R. (1930). *Amer. J. med. Sci.*, **180**, 192.
 Gais, E. S. (1953). *N.Y. St. J. Med.*, **53**, 1844.
 Gauld, W. R., and Lyall, A. (1947). *British Medical Journal*, **1**, 677.
 Himsworth, H. P. (1938). *Quart. J. Med.*, **31**, 373.
 Jacobs, S., Leonard, A. J., and Yager, I. (1953). *J. med. Ass. Ga.*, **42**, 519.
 Kennedy, W. R. (1933). *Canad. med. Ass. J.*, **29**, 482.
 Lawrence, R. D. (1948). *Tubercle, Lond.*, **29**, 255.
 — and Whittaker, H. (1948). *Ibid.*, **29**, 256.
 Lorenzen, J. N. (1931). *Acta tuberc. scand.*, **5**, 265.
 Luntz, G. R. W. N. (1947). *British Medical Journal*, **1**, 943.
 — (1952). *Tuberculosis in the Commonwealth*, p. 270. N.A.P.T., London.
 — (1954). *Lancet*, **1**, 973.
 Mark, M. F., Mosenthal, H. O., and Liu, F. (1942). *Amer. J. med. Sci.*, **203**, 490.
 Ministry of Health (1953). Circular No. 62.
 Munkner, T. (1953). *Acta tuberc. scand.*, **28**, 335.
 Myers, G. B., and McKean, R. M. (1935). *Amer. Rev. Tuberc.*, **32**, 651.
 Oakley, W. (1947). *British Medical Journal*, **1**, 780.
 Poulsen, J. E. (1946). *Acta med. scand.*, **123**, 403.
 Reaud, A. (1953). *Sth. med. J.*, **46**, 248.
 Root, H. F. (1934). *New Engl. J. Med.*, **210**, 192.
 Thosteson, G. C., and Tibbits, R. S. (1953). *J. Mich. med. Soc.*, **52**, 1088.
 Wiener, J. J., and Kavec, J. (1936). *Amer. Rev. Tuberc.*, **34**, 179.

PERIODIC DISEASE: PERIODIC FEVER REPORT OF A CASE

BY

KRIS SOMERS, M.B., M.R.C.P.

From the Department of Medicine, Postgraduate Medical School of London, Ducane Road, London, W.12

Disease conditions manifesting rhythmic recurrence of symptomatology of diagnosable organic origin are well known in medicine—notably epilepsy, Hodgkin's disease, malaria, and brucellosis, to mention a few. The natural history of such illnesses is known and some of them are curable.

However, a distinct group of conditions with periodic symptoms and signs which elude exact diagnosis has been reviewed by Reimann (1951) and called periodic disease. These conditions may occur in childhood or adult life and recur regularly over several years, sometimes for a decade. Various systems may be involved, and symptoms are so regularly cyclic as often to be predictable. The general health is unaffected and patients have been fully investigated without a correct diagnosis or cure being made. They have sought advice from numbers of physicians and have been known at times to seek the help of unorthodox practitioners.

Reimann has included such conditions as periodic fever, periodic abdominal pain, periodic arthralgia, and perhaps periodic (cyclic) vomiting and Behcet's syndrome (recurrent ulceration of oral and genital mucosa with hypopyon) in his description of periodic disease. Several cases of periodically occurring fever closely resembling Reimann's description of periodic fever are to be found in the literature (Wolf and Wolff, 1942; Albahary and de Villeroy, 1951; Hitzig and Fanconi,

1953). Reimann (1947, 1948, 1951, 1953) has followed up a number of cases himself.

The following is presented as a case of periodic fever because of regularly occurring pyrexia for over five years with no convincing evidence of aetiology and, so far, no palliation.

Case Report

A housewife, aged 51, has had undiagnosed pyrexia for five years. Her symptoms date from 1951, about a year after her menopause, at the age of 45. About this time she developed diarrhoea and vomiting lasting several days. She improved, but was left with persistent fever and headache. She was treated without success by her own doctor, who then referred her to a local hospital, where she was admitted for observation three weeks after the onset of her illness.

She had migraine-like headaches at the age of 17. The headache used to be right-sided, lasted most of the day, and was accompanied by nausea and occasionally sickness. At the age of 21 tuberculous lymph nodes were excised on the right side of her neck. She was reported to have had "dry pleurisy" for a few days at the age of 26. At 38 she had shingles on the right side of her chest. She has never been outside England, has never kept pets—bird or animal—and has never been a drinker of raw milk.

The patient has always presented a picture of good nutrition and fitness, and, in particular, has not lost weight. There is an operation scar on the right side of her neck. The ears, throat, and teeth have always been healthy and no abnormal physical signs have been noted in the systems at any of the several careful examinations documented on admission to various hospitals. The spleen has never been palpable, and examinations of the central nervous system have revealed no abnormal signs. Her blood pressure has been in the region of 135/70 mm. Hg.

Her temperature chart has been characteristically periodic, rising over a day or two to 102–103° F. (38.9–39.4° C.), accompanied by a concomitant rise of pulse rate, every third or fourth day, falling again over a day or two to normal levels (see Chart). Associated with the pyrexial bouts she has headaches, anorexia, and malaise. Rectal temperatures taken in hospital have always confirmed simultaneous oral readings.

Investigations

A multitude of investigations have been done and many have been repeated on several occasions.

Haematological.—Routine blood counts made when the temperature was normal and during pyrexial periods showed: Hb, 80–100%; W.B.C., 5,000–8,000 per c.mm. (neutrophils 50–60%, eosinophils 4–5%, lymphocytes 32–43%, monocytes 0.2%). Films—normochromia, normocytosis. E.S.R. 5–12 mm. in one hour (Westergren): normal throughout illness. Platelet count, 262,000 per c.mm. Bleeding-time 2½ minutes. Thick and thin smears negative for malarial parasites. Sternal marrow: no evidence of any specific infection, no reticulosis.

Bacteriological.—Blood cultures sterile (specimens taken at height of pyrexia). No agglutinations for *Salm. typhi* H and O; *Salm. paratyphi* A H, A O, B H, B O, L O; composite salmonellae; *Brucella abortus*; or *Brucella melitensis*. Paul-Bunnell test negative. Leptospiral agglutinations negative with *L. icterohaemorrhagiae*, *L. canicola*, *L. sejroe*, *L. grippityphosa*, *L. bataviae*, and *L. pomona*.

Complement-fixation Tests.—Lymphocytic choriomeningitis virus, negative 1/2; mumps virus, negative (Virus Reference Laboratory, Colindale); influenza virus tests negative;

influenza sol. A <1/8, sol. B <1/8; Q fever, <1/4; psittacosis L.G.V., <1/8; agglutination streptococcus M.G., ±1/10. All negative results (Virus Reference Laboratory, Colindale).

Tests for Toxoplasmosis.—Dye test negative; C.F.T. negative (Hospital for Sick Children, Great Ormond Street).

Miscellaneous.—Mantoux, positive 1/100. Intradermal brucellin test negative. Plasma proteins: total 7.1 g. per 100 ml. (albumin 4.3 g., globulin 2.8 g.). Normal electrophoretic pattern.

Urine Examination.—No albuminuria, sugar, or acetone. No abnormalities on direct examination of deposit on numerous occasions. Culture sterile. Tubercle bacilli not seen on direct examination of deposit or culture on four occasions: 24-hour collections.

Cerebrospinal Fluid.—Lumbar puncture was first done in 1951 after the onset of the pyrexial illness. Pleocytosis was the only abnormal finding that has been inconstantly noted at subsequent examinations (see Table). C.S.F.

Cerebrospinal Fluid Findings

Date	Cells	Protein	Chloride	Sugar	Miscellaneous
		mg. per 100 ml.			
2/2/52	44 red. 2 white	10			Wassermann negative
27/2/52	36 white (70% lymphs, 30% polys.)	20			
5/3/52	50 lymphs	10	770	63	No organisms on direct exam. Negative agglutinations for <i>Br. abortus</i> . Culture sterile. <i>Myc. tuberculosis</i> not grown
18/4/52	27 lymphs	30		57	Lange—no change. No yeasts found. <i>Myc. tuberculosis</i> not isolated on direct exam. or culture
30/5/52	29 white (90% lymphs, 8% polys.)	20	740	53	No organism seen. Torula not isolated on culture
21/5/54	5 red. 11 lymphs	30	720		No organisms on direct exam. No <i>Myc. tuberculosis</i> . Culture sterile. Lange negative

pressure has always been within normal limits and no block has been demonstrated; fluid clear and colourless. A suggestion that lymphocytes in the C.S.F. might, in fact, be yeast cells was unconfirmed. Torulosis was also excluded.

Radiological.—Chest: calcification in both hilar glands; lung fields clear; heart shadow normal. Skeletal: films

Temperature and pulse chart. Rectal temperature indicated by interrupted lines.

of sacro-iliac joints, lumbar spine, and hips showed mild osteoarthritic changes; no other abnormality. Skull: no abnormality shown; pineal body calcified but not displaced. Sinuses: clear. Electroencephalogram: no significant asymmetry or focal abnormality suggestive of a localized lesion.

Treatment

Courses of various antibiotics were tried empirically—sulphonamide, penicillin, chloramphenicol, and chlortetracycline (aureomycin). All were without effect. A course of streptomycin and isoniazid was given, as it was thought that a deep-seated tuberculoma might account for her symptoms and also her persistently abnormal C.S.F. It had no beneficial effect.

On a second occasion she was given a course of cortisone, together with streptomycin, isoniazid, and P.A.S., for about nine weeks. Again her symptoms did not improve at all.

Quinine was tried for a week without any change in the periodicity of her pyrexia. Arsenicals, antihistamines, including chlorpromazine, thyroid, and codeine phosphate were all ineffective.

She did mention that she had been having severe hot flushes, and herself suggested that her symptoms might be related to the menopause. Oestrogens and also testosterone was tried, but no benefit accrued.

Finally, she had a course of mild hypnotherapy in the hope that it might enable her to accept her symptoms rationally, but her state remained unchanged.

Discussion

The whole condition of the patient is puzzling. The outstanding feature is a regularly occurring pyrexia at intervals of three to four days for as long a duration as five years. Careful checks on the temperature ensure that it is not an artifact.

Physical examination has been quite negative apart from the scar of an operation for a tuberculous lymph node twenty years ago. There are no abnormalities in the central nervous system. All relevant investigations have drawn a blank. Interestingly enough, the first lumbar puncture was negative, and it may be just possible that the slight pleocytosis could be a consequence of repeated lumbar puncture.

The patient has been under observation at regular intervals and no change has been noted. The condition is apparently benign. Her general health and nutrition remain unaltered, except that over the years she has become nervous and introspective, which seems a reasonable reaction to such a prolonged and tedious organic disease.

Various specialists—medical, surgical, and gynaecological—have seen this patient and are unable to arrive at an aetiological diagnosis.

An infective cause appears unlikely. The pattern has been one of regular, uniform, recurrent pyrexia over many years, not suggestive of a known condition. It seems reasonable to assume that there is no good evidence of tuberculous activity. Empirical treatment with sulphonamide, penicillin, broad-spectrum antibiotics, and anti-tuberculosis drugs (with the possibility that she might have a tuberculoma) have been of no avail.

Atypical epilepsy has been suggested as a diagnosis in patients with periodic abdominal pain in whom abnormal electroencephalographic patterns are demonstrated, relief being obtained with anticonvulsants (Moore, 1944, 1948). Our patient, however, shows no suggestion of epilepsy on electroencephalography.

An obvious reticulosis has been excluded on clinical and haematological grounds.

Periodic fever may be a hypothalamic sign of craniopharyngioma (Müller and Wohlfart, 1950). In the absence of pituitary involvement, radiological calcification of the

tumour, and signs of raised intracranial tension with such a prolonged history, that diagnosis seems untenable.

Reimann's Case 9 (1949) developed periodic fever, similarly after the menopause, which may or may not have served as a trigger mechanism to set it off. It is difficult to postulate a definite endocrine basis. In some patients episodes lessen in intensity or cease after the menopause. Menstruation does not seem to influence periodic disease (Reiman, 1956, personal communication).

Allergy might be suggested as a cause of periodic disease; but in the absence of associated allergic phenomena, such as asthma, urticaria, and hay-fever, family history and lack of response to antihistamines and corticotrophin and to cortisone given empirically in adequate doses, it seems hardly tenable here.

In Wolf and Wolff's case (1942) fever was related to emotional disturbances. No one, however, has been able to entertain a psychogenic cause for this patient's illness. If anything, her nervous symptoms are secondary to her prolonged periodic fever. She herself suggested that her condition might be related to the menopause, and, indeed, as one physician remarked, "who are we to say that it is not?" She has reached the stage where she most adamantly resists the suggestion of any further investigation, such as lumbar puncture, cerebral angiography, or ventriculography unless it can be guaranteed to produce the answer to her illness.

It is difficult to know what management is advisable. Amidopyrine slightly reduced the temperature in one of Reimann's (1951) cases, but the cycles of periodic fever were unaltered. Another case appeared to respond temporarily on cortisone (Reimann, 1953). We have not been able to demonstrate any change with antipyretics or cortisone.

Dr. Reimann has been consulted about this case and has made these comments: "I feel quite sure that this patient has periodic disease, or, more specifically, periodic fever. From all the evidence, positive and negative, I don't know what else it could be. You have excluded most other possibilities, but, from the old history, the spectre of tuberculosis remains despite the absence of evidence. Even if tuberculosis were present, the periodicity of fever still would be of interest. She is lucky that (as yet) none of the more distressing symptoms of severe abdominal and thoracic pain or of arthralgia occur. This should be of some consolation to her. Unless you find some cure the best she can hope for is that the intervals may lengthen and/or the episodes may lessen in severity as time passes."

With all the therapeutic failures it seems necessary to regard her illness as periodic fever, if only to save her from further medical and surgical measures.

Summary

A case of periodically recurring pyrexia for five years is detailed. No satisfactory explanation is possible. It would seem a sufficiently clear-cut case to merit inclusion in Reimann's well-described condition—periodic disease.

Thanks are due to Professor J. McMichael for his constant encouragement, to Dr. H. A. Reimann, of New York, for his kind help and advice, and to Dr. Horace Joules, Central Middlesex Hospital, London, for access to notes made whilst the patient was under his care.

REFERENCES

- Albahary, C., and de Villerooy. H. (1951). *Bull. Soc. méd. Hôp. Paris*, 67, 1391.
- Hitzig, W. H., and Fanconi, G. (1953). *Helv. paediat. Acta*, 8, 326.
- Moore, M. T. (1944). *J. Amer. med. Ass.*, 124, 561.
- (1948). *Rev. Gastroent.*, 15, 381.
- Müller, R., and Wohlfart, G. (1950). *Acta med. scand.*, 138, 121.
- Reimann, H. A. (1947). *Ibid.*, Suppl. 196, p. 666.
- (1948). *J. Amer. med. Ass.*, 136, 239.
- (1949). *Ibid.*, 141, 175.
- (1951). *Medicine (Baltimore)*, 30, 219.
- Wolf, S., and Wolff, H. G. (1942). *Ibid.*, 70, 293.
- (1953). *A.M.A. Arch. intern. Med.*, 92, 494.