In 1953, before vaccination was offered more widely, nine unvaccinated workers from an adjoining laboratory were sharing a refrigerator in the Q fever laboratory, which they repeatedly entered. None developed either clinical or serological evidence of infection. On the other hand, Patient B entered the laboratory for a few seconds and later succumbed to the disease.

This effect might be due to chance or individual variation in susceptibility. It is possible, however, that small repeated infections produce an immunity which is not detectable by conventional tests for antibody.

The resistance of *R. burneti* to adverse environmental conditions has allowed its transfer to new hosts by varied and often devious routes. The two infections, which resulted from contamination of the drain-pipe, suggest that the possibilities are not exhausted, but it is fortunate that the circumstances of this particular incident are unlikely to recur frequently.

I am grateful to Professor H. R. Dean for permission to publish this paper.

REFERENCES

- Beeman, E. A. (1950). *Publ. Hlth Rep. (Wash.)*, 65, 88.
 Malloch, R. A., and Stoker, M. G. P. (1952). *J. Hyg. (Camb.)*, 50, 502.
 Meiklejohn, G., and Lennette, E. H. (1950). *Amer. J. Hyg.*, 52, 54.
 Oliphant, J. W., Gordon, D. A., Meis, A., and Parker, R. R. (1949). *Ibid.*, 49, 76.
 Smadel, J. E., Snyder, M. J., and Robbins, F. C. (1948). *Ibid.*, 47, 71.

A SURVEY OF DIABETES IN WEST CORNWALL

BY

C. T. ANDREWS, M.D., F.R.C.P.

Physician, West Cornwall Clinical Area

The objects of this survey were (1) to determine the incidence of diabetes in the West Cornwall clinical area, which is an isolated geographical entity with a population of a quarter of a million, largely shut off from the rest of England by Bodmin Moor; and (2) to find out to what extent the diabetic patient constitutes a load on the community. In assessing the second object four main agencies were considered: the home, the district nurse, the general practitioner, and the hospital service.

Incidence

Approximately one-third of the population has been surveyed through general practitioners, the doctors taking part being chosen so as to obtain an even geographical distribution and a balance between town and country practices.

In the spring of 1953 preliminary invitations were sent to 55 general practitioners, and a subsequent letter invited each one who agreed to co-operate to survey his list and complete a short questionnaire (see Appendix). In this way figures were obtained covering the practices of 48 doctors distributed from near Land's End to Camelford and Bodmin. These varied in size from just over 500 to one partnership of nearly 13,000. The results are given in Table I. The rate per 1,000 is seen to be 5.6. The incidence varied within the practices surveyed from 3.2 to 8.4 per 1,000.

Of the 593 patients accepted as diabetics, the diagnosis had been confirmed in 579 by attendance at a diabetic or general medical clinic, by blood-sugar investigation, or by previous investigation as a hospital in-patient. The

remainder were accepted after careful scrutiny, and, in some cases, after personal discussion with the doctor making the return. No significant difference was found in the predominantly rural and predominantly urban practices. A cross-check on these findings was made by counting the number attending general medical and

TABLE I.—Incidence of Diabetes in West Cornwall

Total population of West Cornwall	266,389
No. of list patients surveyed	104,827
Percentage of total population surveyed	39.92
Total number of diabetics found	593
Number of diabetics per 100,000 population	565.7
Estimated number of diabetics in West Cornwall	1,507
Percentage attending diabetic or general medical clinics	77.4
Estimated total number attending diabetic or general medical clinics	1,167
Actual number found attending	1,141

diabetic clinics, and the number (1,141) was found to accord closely with the estimated figure obtained from the general-practitioner returns (1,167).

Very little information is available on the incidence of diabetes in this country. During the second world war the necessity to obtain a medical certificate for special rations afforded an opportunity of arriving at an approximate figure. In 1947 it was thought that nearly all diabetics were availing themselves of the extra allowances. The incidence calculated from medical certificates in this year was 231 per 100,000 for England and Wales. It is possible, however, that many of the milder diabetics in rural areas did not avail themselves of the special rations.

The advent of the National Health Service has made it possible to enlist the help of the general practitioner in investigations of this type. In a recently published survey of the work of eleven practices covering the years April, 1952, to March, 1954, it was found that the annual rate of patients having consultations for diabetes varied from 2 to 7 per 1,000 population within the practices investigated. The average annual rate for all the practices was 3.6.

A partial explanation of the high incidence of the disease in Cornwall is provided by a study of the population structure. In this area approximately 19% of the population is aged 60 or over, as compared with 15% for the rest of the country. As diabetes predominates in the older age groups a higher incidence would be expected. A more powerful reason is probably provided by the isolation of the area, with consequent inbreeding in the past. Isolation in West Cornwall has meant a great deal more than lack of intercourse with the rest of England. The small fishing village and rural township have kept themselves apart to a degree rarely seen in the rest of England. Local geography has fostered this isolationism, for many of the coastal villages even to-day have no more than one bus a day or even one a week.

Sex Distribution

The distribution between the sexes is given in Table II, and is compared with that derived from a count of special ration certificates for the whole of the country in 1947. It will be seen that the sex distribution in Cornwall closely follows that of England and Wales as a whole. The

TABLE II.—Sex Distribution

	Males		Females	
	No.	%	No.	%
General-practitioner survey	234	39.5	359	60.5
Old clinic cases	173	39.3	267	60.7
Present Truro Clinic	178	36.9	304	63.1
England, 1947 (special ration certificates)		38.0		62.0

slightly greater predominance of female patients at the clinics as compared with the general-practitioner survey may be explained by the fact that the principal clinic in the area is held at 10 a.m., so that men in jobs may have some difficulty in attending.

Age at Onset

In relation to the load on the community it is necessary to consider the age at which the disease first produces symptoms and the manner of its presentation.

The average age of onset as determined from the case records of 901 clinic patients was 52 years. The age of onset in relation to sex shows a striking difference between male and female diabetics (see Chart). The female curve rises

Chart showing age at onset of diabetes in males and females.

sharply to a peak in the sixth and seventh decades while the male curve is flatter. Of greater interest perhaps is the relative preponderance of males in the first four decades and the tendency for this preponderance to reappear in extreme old age. The effect of this on the incidence of the disease in the over-70 age groups is accentuated by the tendency in West Cornwall for the female diabetic to die at an earlier age than the male (see below). The striking excess of females in the sixth and seventh decades is well known and accounts for their overall preponderance in the survey (60.5% in a total of 693 cases).

Mode of Onset

The material used for this part of the investigation was the case records of 573 clinic patients, in nearly all of which the history had been taken with this investigation in mind. It will be seen from Table III that the classical symptoms of thirst, polyuria, and weight loss constituted the reason for seeking advice in only 41.4% of the cases. Incidental illness and routine examination together account for the first attendance of 13.5% of all the cases attending. Some of these were picked up by shrewd general practitioners looking after minor illness such as influenza at home. Most, however, were diagnosed on admission to hospital for illness in no way associated with diabetes. Routine medical examination for life assurance, for the Services, or for employment led to the diagnosis of 3.7%.

TABLE III.—Mode of Onset in 573 Cases

Classical (thirst, polyuria, weight loss) ..	237 (41.4%)
Incidental illness ..	56 (9.8%)
Eye complications ..	38 (6.6%)
Pruritus and eczema ..	53 (9.2%)
Skin sepsis (including foot sepsis) ..	46 (8.0%)
Weight loss alone ..	30 (5.2%)
Routine examination ..	21 (3.7%)
Lassitude alone ..	12 (2.0%)
Neuritis ..	9 (1.6%)
Coma or pre-coma ..	20 (3.5%)
Gangrene ..	20 (3.5%)
Other ..	31 (5.4%)

A consideration of these groups raises the question of whether there are not in the community to-day many diabetic patients more or less symptom-free and therefore undiagnosed. While it is true that many of this group on direct questioning admitted having symptoms of diabetes such as polyuria, they were not such as to bring them to medical attention. It seems likely, indeed, that routine urine testing of the whole community would lead to a further substantial addition to the figure of incidence in this area at least.

The figure for skin sepsis includes also cases presenting with foot sepsis. All but one of the cases of gangrene were of the lower limb. Several of the 20 cases presenting in coma or pre-coma appear to have had no symptoms suggestive of the disease till a few days before the onset of coma.

Use of Clinics

Information was obtained on this subject from the answers to the questionnaire. They show that 77.4% of all patients attend either a diabetic or a general medical clinic (Table IV). Some were attending more than one clinic at

TABLE IV.—Attendance at Clinics

Total number of diabetics found (G.P. survey) ..	593
Information re clinic attendance not given ..	26
Remainder ..	567
Attending clinics ..	445 (78.5%)
" diabetic clinic ..	418 (73.7%)
" general medical clinic ..	21 (3.7%)
" other clinics ..	11 (2.0%)
Hospital in-patients ..	26 (4.6%)

the same time—for example, diabetic and chest. Some were regarded as still attending the clinics though in-patients at the time of the survey.

The extent to which the clinics were used varied widely, and this variation appeared to bear no relation to geographical accessibility. For example, in one large practice returning 63 patients all but three of these attended the diabetic clinic four miles away. In the same town the attendance rate from another doctor's practice was 18%. It was clear that a small proportion of the general practitioners regarded themselves as competent to look after their own diabetics and referred to the clinics only those presenting special problems.

It might be thought, particularly in a sparsely populated rural area, that age would exert a considerable influence on attendance, but this is apparently not so. The average age of those attending the clinics is 64, of those not attending 65. Of the 120 patients shown as not attending the clinics in the general-practitioner returns, 50 (41.7%) were males and 70 (58.3%) females. These figures show a slightly greater attendance rate for females than for males in proportion to their total numbers.

Problems of the Home

These are mainly those constituted by special diet, insulin administration, and intercurrent illness. Of these, the dietetic problem gives rise to least anxiety in the minds of diabetic patients. Of 501 patients investigated at the Truro Clinic, 100 (20%) were adhering to a strict dietetic regime (Lawrence line ration scheme); 363 (72.4%) were observing some dietetic restriction short of this; and 38 (7.6%) might be regarded as observing no dietetic restrictions.

It is probably true to say that the problem of diet constitutes a handicap to no more than 20% who adhere to a strict regime. The problem of insulin administration, however, is of a different order. Of the clinic patients, 18% are found to require no insulin, 37% use one type, and 45% two types (generally soluble and protamine zinc) daily, while 5% of all patients attending require more than one injection of insulin daily. Thus insulin administration constitutes a problem in 82% of all cases, and amongst this number the problem is of several grades from the point of view of domestic responsibility. The patient who requires one injection daily of one type of insulin is much less of a problem than the patient who has to manipulate two separate phials, while the patient who requires two daily injections may be much restricted in activity if she is not able to cope with the problem herself.

Table V shows the home arrangement for insulin administration in 383 patients attending the Truro Clinic. It shows that the problem confronting the female patient is solved through a wider variety of agencies than that of the male. The number of patients regularly giving their own injections is 80.4% of the males and 62.5% of the females. For those, however, who are unable to give their own injections the wife proves more valuable to the helpless

TABLE V.—*Insulin Administration in 383 Cases*

	Males		Females	
	No.	%	No.	%
Self	111	80.4	153	62.5
Wife	18	13.0		
Husband			26	10.6
Father	1	0.8	1	0.4
Mother			1	0.4
Daughter	3	2.2	27	11.0
Son	2	1.4	8	3.2
Neighbour	1	0.8	9	3.7
Nurse	2	1.4	20	8.2
Total	138		245	

Nurse gives 5.7% of all insulin injections. 69% of all patients give their own insulin.

husband (13%) than the husband to a dependent wife (10.6%). The helpless female patient in fact chooses from a wide range of assistance of which daughter, husband, and district nurse are the most important. The helpless husband tends to rely exclusively on his wife, and usually only when a widower does he resort to help from other agencies.

The figure of 5.7% of all injections given by the district nurse is probably too low. In country districts she plays an important part in the training of the patient or a relative in insulin administration. This applies particularly to the milder diabetic who is stabilized as an out-patient. This phase ended, she generally ceases to attend except for the small percentage who are dependent on her for daily insulin. There are some, however, who require a supervisory visit once a week or oftener, and on these occasions the nurse usually gives the insulin. In some cases where measurement of the correct dose constitutes the main problem she may fill the syringe and leave it for the patient herself to administer the next day. In some cases of defective eyesight a member of the household measures the dose and the patient gives the injection.

The commonest reasons given for inability to administer their own insulin were defective eyesight, nervousness, and general frailty of old age. Though defective eyesight was the commonest reason given it often seemed to be mainly an excuse to cover lack of courage. The relationship to age was very definite, as shown by Table VI. From this

TABLE VI.—*Age and Inability to Give Own Insulin*

Age Group:	0-9	10-	20-	30-	40-	50-	60-	70+	Total
Total on insulin	4	6	8	25	42	90	115	93	383
Dependent on others	3	1	0	2	4	18	33	52	113
% " "	75	16.6	0	8	9.5	20	29	56	29.5

it is seen that dependence on others for insulin administration increases numerically in each decade from the third and that more than half of those over 70 are dependent on others for this service.

Intercurrent Illness

How much illness does the diabetic patient have? In an attempt to answer this question 483 patients passing through the clinic at Truro were questioned at each quarterly attendance during 1954 regarding illness in the previous three months. The investigation has now been continued, and has been completed for the year 1955, during which 663

TABLE VII.—*Days in Bed 1954 and 1955*

	Males		Females	
	1954	1955	1954	1955
None	139 (77.7%)	216 (82.4%)	214 (70.4%)	317 (79%)
1-7 days ..	14 (7.8%)	15 (5.7%)	27 (8.9%)	24 (6%)
8-35	19 (10.6%)	25 (9.5%)	55 (18.1%)	50 (12.5%)
Over 35 days	7 (3.9%)	6 (2.3%)	8 (2.6%)	10 (2.5%)
Total	179	262	304	401

patients were questioned. The results are set out in Table VII. At the onset it became clear that in dealing with the older age groups it was extremely difficult or impossible to obtain a true measure of morbidity. Many of the patients are frail or infirm. Standards based on fitness for work or confinement to the house have little meaning in a group in which many are over working age or are housewives doing a limited job of work in their own homes. In the end it was decided to inquire about days spent in bed through illness. Two main reasons for this inquiry were (1) such illness constitutes a load on the home to a much greater extent than that which does not require rest in bed, and (2) it is a measurable entity which the patient can easily remember.

From this investigation the fact emerges that 73% of all patients had no illness requiring rest in bed in the year 1954 and that the proportion was rather higher for 1955, a year in which less respiratory illness and fewer fractures of the lower limb occurred. These figures might not be so good if they included diabetics not attending the clinics. This point is discussed below.

The figures also include time spent by clinic patients in hospital. Owing to the large number of hospitals involved it has not been possible to determine this figure accurately for clinic patients, but the hospital requirements of the total diabetic population of the area are discussed in a later section.

It will be seen that illnesses of one to five weeks predominated in the females in both years reviewed. This was due almost entirely to the greater tendency to fracture of the lower limb and abdominal operations in this group.

TABLE VIII.—*Principal Causes of Incapacity During 1954 and 1955*

	1954	1955
No. of patients questioned	483	663
Total No. of bed days, all patients ..	2,385	2,727
Average for all patients attending the clinic ..	4.9	4.1
laid up	18.2	21
Total No. of patients confined to bed ..	131	130
Respiratory infections; colds, influenza, bronchitis, sinusitis, pneumonia ..	666 days	632 days
Percentage of all bed days	27.9	23.2
No. of patients affected	42	43
Recognized diabetic complications: coma, pre-coma, hypoglycaemia, cataract extraction, skin sepsis, gangrene ..	370 days	377 days
Percentage of all bed days	15.5	13.8
No. of patients affected	26	28
Abdominal operations	307 days	276 days
Percentage of all bed days	12.9	10.1
No. of patients affected	12	10
Fractures and other injuries	249 days	82 days
Percentage of all bed days	10.4	3.0
No. of patients affected	7	4
Cardiovascular disease	171 days	262 days
Percentage of all bed days	7.2	9.6
No. of patients affected	5	9
Rheumatism, fibrositis, backache ..	55 days	58 days
Percentage of all bed days	2.3	2.1
No. of patients affected	6	5
Infectious fevers	149 days	118 days
Percentage of all bed days	6.3	4.3
No. of patients affected	7	6
Pulmonary tuberculosis	0	310 days
Percentage of all bed days	0	11.3
No. of patients affected	0	2

Table VIII gives the time spent in bed with the principal groups of illness. It amplifies Table VII and shows that the average number of days during which a clinic patient was laid up was 4.9 and 4.1 respectively for the years 1954 and 1955. There was a slight predominance of illness in the females: this was more definite in the year 1955, in which the average duration of bed illness was 3.6 days for males and 4.5 for females. In this respect the findings agree with those of the general-practitioner survey recently published by the General Register Office (1956).

An analysis of the illnesses shows that those which predominate in the lives of diabetics are not the recognized complications of diabetes but the sort of illnesses prevalent in the general community of corresponding age, with some diabetic complications added. These latter account for only about one-seventh of all bed days.

The predominance of respiratory ailments in a group which includes a large percentage over 60 is not surprising.

Respiratory ailments also account for much of the minor illness in this group of patients. All the fractures noted in this group are of the lower limb. The total disablement time to which they give rise may be very great in proportion to their number (only 7 in 483 cases in 1954). There were also a number of fractures of the upper limb which, though not necessitating rest in bed, gave rise to a considerable amount of disability.

It is probable that the figure for cardiovascular disease would be higher if this section included those patients who do not attend the clinics. There is some reason to believe that this is one of the common reasons for failure in attendance. The patient who, for example, develops congestive heart failure or who has had an attack of coronary thrombosis decides, very often with his general practitioner's agreement, that his cardiac condition no longer permits attendance at the clinic.

It may too easily be assumed from a consideration of Table IX that the recognized diabetic complications are a nearly negligible cause of morbidity in the lives of diabetic

TABLE IX.—Complications in 738 Clinic Patients (Average Period of Observation 6.9 Years)

Diabetic Complications		Cases
Cataract	74 (10%)	
Retinopathy	45 (6.1%)	
Skin sepsis	72 (9.8%)	
Severe pruritus	41 (5.6%)	
Dermatitis (generalized)	14 (2.0%)	
Gangrene	33 (4.5%)	
Symptomatic neuritis	36 (4.9%)	
Diabetic coma	17 (2.3%)	
Non-diabetic Complications		
Hypertension (systolic 180+ or diastolic 110+) ..	190 (25.7%)	
Other forms of cardiovascular disease ..	72 (9.8%)	
Congestive failure	28	
Coronary insufficiency	9	
Auricular fibrillation	8	
Intermittent claudication	10	
Cerebrovascular disease	17	
Respiratory ailments	38 (5.1%)	
Bronchitis, chronic	15	
Asthma	9	
Pulmonary tuberculosis	8	
Pleurisy, ? tuberculous	4	
Pulmonary fibrosis	2	
Thyroid disease	21 (2.8%)	
Thyrotoxicosis	9	
Simple goitre	10	
Myxoedema	2	
Rheumatoid and osteoarthritis	15 (2.0%)	
Neuroses and psychoses	7 (1.0%)	
Carcinoma	8 (1.1%)	
Gall-bladder disease	20 (2.7%)	

patients. Such is far from being the case. While troublesome, most complications of diabetes are compatible with a reasonable degree of activity and do not preclude attendance at the clinics.

Table IX sets out the frequency with which recognized diabetic complications have been observed in a group of 738 clinic patients observed over an average period of nearly seven years and places them in comparison with illness of other types. Eye diseases, which head the list, cause prolonged and often permanent disability, though in many of the 119 cases observed the degree of visual impairment is not great. Skin sepsis causes a large number of minor illnesses from which full recovery is the rule. By and large, it is only infections of the foot which are likely to lead to permanent disablement or to endanger life.

The small figure for pulmonary tuberculosis emphasizes the comparative infrequency of this disease in the controlled diabetic to-day. The incidence of gall-bladder disease is certainly much higher than the figure given under this heading suggests. It could be assessed accurately only by radiological examination of a large number of diabetic patients. No figure is given for the incidence of hypoglycaemic reactions. These are very common, and it is probable that in the course of seven years nearly every patient has experienced hypoglycaemic symptoms. In an educated diabetic community, however, these seldom amount to much except in the brittle unstable diabetic, who constitutes a problem in every diabetic clinic.

Work of the District Nurse for Diabetics

The main figures showing the work of the district nurse for diabetics are set out in Table X. The principal information that emerges from this investigation is that the district

TABLE X.—Work of the District Nurse for Diabetics

Total No. of district nurses involved ..	98
No. of diabetic patients in the area (estimated) ..	1,507
" " " " visited in 1954 ..	228
Percentage of all diabetics visited ..	15.1
Total No. of general nursing visits to all sick patients in the area in 1954 ..	135,468
Total No. of visits paid to diabetics ..	25,068
Percentage of total visits paid to diabetics ..	18.5
No. of visits per diabetic patient visited ..	110
Average No. of visits per diabetic patient in the area ..	16.6
No. of diabetics requiring regular visits ..	125
Percentage of all diabetics in the area ..	8.3
No. dependent on district nurse for daily insulin ..	96
Percentage of all diabetics in the area ..	6.4

nurse is concerned with only a small proportion (15.1%) of these patients, yet these entail a large amount of work because so many are dependent on her for daily insulin. How many are so dependent? From a study of Table V, which was derived from an interrogation of clinic patients, it would appear that only 5.7% of all patients having insulin require the daily help of the district nurse. Here the figure is 6.4% of all diabetic patients (based on an estimate of 1,507 diabetics in the area). This would mean 8% of all patients having insulin. This apparent discrepancy may be explained partly by the fact that the first figure relates to clinic patients only, while Table X is concerned with all diabetics in the area. It would appear likely that the patient who has his diabetes managed exclusively at home is more dependent on the district nurse than the one who regularly attends the clinic. Secondly, Table V is based on patients' answers; the figures in Table X come from the district nursing service.

As already suggested, some of those whom the district nurse regards as dependent are not in fact given the daily injections by her. She may visit the patient on one day, give the injection, and leave a further dose ready for administration the next day by a member of the family or by the patient. The number of patients in whom insulin administration is a combined operation between nurse, patient, and family would appear to be considerable.

Load on the General Practitioner

The amount of work done for diabetics in the area by general practitioners is set out in Table XI. A large number of general practitioners failed to answer this section of the

TABLE XI.—Home Visits and Surgery Attendances in 1954

Home Visits		
Total No. of diabetics	593	
No information	131	
Remainder	462	
No. visits	236 (51.1%)	
1-5 visits	124 (26.8%)	
6-10	40 (8.7%)	
Over 10 visits	62 (13.4%)	
Total No. of visits paid to 462 patients ..	1,721	
Average per diabetic patient ..	3.7	
" " " " visited ..	7.6	
Surgery Attendances		
No. of attendances by 462 diabetic patients ..	2,815	
Average per diabetic patient ..	6.1	

	No.	Visits	Average	Surgery Attendances	Average
Males ..	187	454	2.4	1,459	7.8
Females ..	275	1,267	4.6	1,356	4.9

questionnaire as no records were kept of home visits or surgery attendances. Information was, however, obtained on the detailed care of 462 patients.

There was wide variation in the amount of visiting and surgery attendance from patient to patient and from practice to practice. For example, one patient was returned as

paying 365 visits to the doctor's surgery in a year (presumably to have his daily insulin given). On the other hand, there was the doctor who found it necessary to pay only three visits to a total of 15 diabetics in a year. Surgery attendance seemed to be the general rule in some practices, in others the doctor seemed to prefer to see the patient at home. There was no significant difference between the urban and rural groups of practices in regard to the ratio of surgery attendance to home visiting.

It will be seen that about 50% of all diabetics require some attention in their own homes by the doctor during the year. This is in sharp contrast to the corresponding information on the work of the district nurse, whose visiting is concentrated on a much smaller section of the diabetic population (15%). It would appear, however, that, owing to the nurse's responsibility for daily insulin administration for a small number, the total visits paid and the average per diabetic patient in the area is much higher than that of the doctors.

The addition of surgery attendances, however, alters the picture considerably. *In toto*, the doctor sees the average diabetic patient approximately 10 times in the year. The district nurse visits the average diabetic patient 16.6 times. The doctor, however, is consulted by (or sees) 91% of his diabetic patients at some time during the year. The district nurse sees no more than 15.1% of them, and this last is predominantly a special group in which, by reason of age, visual failure, nervousness, or lack of competent family or neighbour help, insulin administration necessitates her services.

The figure of 91% of all diabetics who have consultations either at home or at the doctor's surgery compares with that of 67.6% given for the year ending March, 1954, in the general-practitioner survey of the General Register Office (1956). The high figure for Cornish diabetics is partly explained by the age and sex incidence. The highest consultation rate given in the general-practitioner survey is that for women over 65. A comparative study of the age structure of the Cornish diabetic population and that of the survey shows that 38% of the former but only 9% of the latter are in this age group.

The same consideration arises in comparing consultation rates. The average for all patients in the general-practitioner survey is 3.7 per annum; for all Cornish diabetics it is approximately 10. But for women over 65 it is 5.7 per annum in the general-practitioner survey, and in the year most closely approximating in time to that of our Cornish survey it is 6 per female patient over 65. We have no comparable Cornish figures for a normal general-practice population, but if the Registrar-General's general-practitioner figures are representative of the country as a whole it seems clear that Cornish diabetics require about twice as many consultations in a year as the normal population of similar age.

Another significant figure here is that of patients who required neither home visits nor surgery attendance in the year. There were 43 (9.3%) of these in a total of 462 about whom information was available. Twenty-eight of these are recorded as on an insulin regime and 15 as having no insulin. These figures are of some importance in view of attempts to calculate the incidence of diabetes from surgery attendances and home visits over a given period. Although it must be assumed that those using insulin would at some time or times in the year make contact with the doctor's surgery, there seems no good reason why those on diet alone should do so.

In association with the figure of 49% of patients requiring some domiciliary visiting during the year by the general practitioner should be considered the figure from Table VII of 70 to 82% of patients having no days in bed. From this it seems clear that a large number of the doctors' visits are made to patients who are not confined to bed. The patient, for example, with an attack of bronchitis or a sore foot will stay indoors and send for the doctor, but he will often weather the illness in his armchair by the fire. Further

confirmation of this view is obtained by considering the figure of 3.7 domiciliary visits by the general practitioner in association with the figure of 4.9 bed days. It would seem probable, indeed, that the amount of genuine illness in the life of a diabetic patient might be indicated with fair accuracy by doubling the number of days spent in bed—that is, 10 days' illness per annum.

The last section of Table XI compares the attention given to male and female patients at home and at the doctor's surgery. It will be seen that while the male diabetic gets most of his attention at the doctor's surgery the female patient gets it mainly at home. This is no doubt related to the fact that men can more readily arrange to fit in surgery attendances on the way to and from work. It is, however, in contrast to experience at the clinics, where the tendency is for a better attendance of women than men in proportion to their overall numerical distribution.

Hospital Requirements

What are the hospital requirements of diabetic patients? Why do they go there and how long do they stay? The answers to these questions were sought by inviting each of the 22 hospitals in the area to supply information regarding diabetic admissions during the year 1954-5 and by taking a census of diabetics in hospitals in the area on a day in November, 1955. Table XII shows the result of the first of these investigations. About 42% of those seeking admission to hospital did so for reasons not directly associated with the diabetes. This is not surprising in view of the findings for illness in the home. The diabetic picture is, however, more prominent in Table XII than in that of home illness (Table VIII). This is largely accounted for by the large number who have to come in for initial stabilization. The comparatively low incidence of respiratory ailments contrasts sharply with the high figure for illness in the home. This is in keeping with the general trend of medical care to-day, for the general practitioner can usually cope with such ailments even in the diabetic patient unless severe complications arise.

TABLE XII.—Reasons for Admissions—All Hospitals

Diabetic Causes		Non-Diabetic Causes	
Stabilization	51	Congestive failure	6
Restabilization	10	Other heart disease	3
Cataract and other eye com-		Cerebrovascular disease ..	8
plications	17	Gall-bladder	6
Skin sepsis	12	Thyroidectomy	2
Hypoglycaemia	12	Cancer	6
Diabetic coma	9	Fracture (5) other injuries (2) ..	7
Pre-coma	4	Respiratory infection	5
	115	Pulmonary tuberculosis	6
		Other infections	8
		Intestinal haemorrhage	4
		Prostatic enlargement	3
		Acute abdomen	2
		Social reasons	6
		Pregnancy	3
		Other	10
			85

Altogether 227 patients were admitted to hospital in the year surveyed. Full particulars regarding the reasons for admission are available for 200 of these. The total number of bed days for these 200 was 3,122, and the average duration of stay 16 days. If the figures for the geriatric service and for the tuberculosis patients are excluded the average duration of stay was 13.5 days. Diabetes and its recognized complications accounted for 1,623 days, and the average duration of stay of this group was 14 days (115 cases).

A census taken in the hospitals of the area at the end of November, 1955, showed 44 diabetic patients in hospital. Of these, 15 were in the wards of the general hospitals in Truro, Penzance, Redruth, and Falmouth; 13 in St. Lawrence's Mental Hospital, Bodmin; 13 in the hospitals of the geriatric service; and 3 (all cases of pulmonary tuberculosis) in Tehidy Sanatorium.

From a study of all the hospital figures it is possible to hazard a guess at the bed requirements for diabetics which may serve as a guide to those planning diabetic units. On

the basis of an average stay of 14 days a 15-bed unit in this area should be able to cope with approximately 300 admissions per annum, which is 73 in excess of the total admission figure for the area for the year 1954. Approximately three-fourths of the admissions were to medical wards, so that, given an even admission rate, 12 beds could deal with all medical admissions. There are, however, seasonal fluctuations, so that the actual range of bed occupancy for medical diabetics is from 5 to 18. There are also generally some patients waiting for admission. Taking all these factors into consideration, it seems reasonable to suggest a figure of one medical bed for every 100 of the diabetic population. The provision of a similar number of long-stay beds to cope with social and special clinical problems seems necessary.

Table XIII shows the cause of death, so far as can be ascertained, of 76 diabetics who have died in this area in the past three years. The figures were obtained partly from a follow-up of all diabetics who cease to attend the

TABLE XIII.—Recorded Causes of Death in 76 Diabetic Cases

	No. of Cases	%	Registrar-General (Total Pop.)
Cerebrovascular accidents	17	51.3	.59
Coronary thrombosis	11		
Congestive failure	9		
Kidney disease	2		
Diabetic coma	7	9.2	7
Respiratory infection	6	7.9	
Cancer	6	7.9	
Post-operative deaths	4	5.3	
Senility	4	5.3	17
Bleeding peptic ulcer	2	2.6	
Pulmonary tuberculosis, suicide, appendix abscess, subcutaneous abscess, pulmonary embolism, acute pancreatitis, urinary infection, gangrene	8	10.5	

clinics at Truro and Redruth and partly from a search of hospital records. It is believed that these are more accurate than those obtained from registration returns. Joslin *et al.* (1952) estimate that in the United States only about two-thirds of diabetics who die have their diabetes mentioned on the death certificate. It seems probable that the practice here is similar.

The figures in the third column provide a comparison with those derived from the Registrar-General's (1955) returns for Cornwall for the year 1954. It seems surprising that there is not a greater incidence of death from cardiovascular and renal disease in diabetics than in the general population. The series, however, is a small one, and it is possible that the figures might look substantially different if the investigation were extended for 10 years. The number of deaths from diabetic coma (seven in three years) is small, but could probably be further reduced with better liaison between hospital and general practitioner. In a concentrated urban area, with good hospital facilities for dealing with this emergency, deaths from diabetic coma might well be eliminated.

Table XIV shows the age at death of 129 cases (50 men and 79 women) for which this information was available. It is worth noting that the peak decade for deaths is the

TABLE XIV.—Age at Death of 129 Patients

Age:	0-39	40-49	50-59	60-69	70-79	Over 80
Men	2	2	8	11	18	9
Women	2	1	16	32	26	2
Total	4	3	24	43	44	11

Average age at death: men, 68.9 years; women, 66.1 years.

eighth for men and seventh for women. The average age at death of the 50 men was 68.9 years and of the 79 women 66.1 years; for both sexes it was 67.1 years. The figure compares with that of 61.9 years given by Joslin *et al.* (1952) for 656 of their own patients. We are, however, dealing with an older population in West Cornwall, which may partly explain the high average age at death. It should be remembered also that the figures given are derived from a follow-up of clinic patients personally supervised: the

figures for the whole diabetic population would almost certainly be less favourable.

A study of registration figures for the administrative County of Cornwall for the year 1954 discloses the fact that 54% of male deaths and 67% of female deaths occurred at 70 or over. The corresponding figures in this diabetic series are 54% (male) and 35% (female). It appears, therefore, that diabetes as seen in Cornwall has a definite effect on the expectation of life in the female patient, but little or no effect on that of the male.

Summary

In a survey of diabetes in West Cornwall the incidence of the disease has been found to be 5.6 per 1,000 of the population. This figure is high in comparison with previously published figures for the country as a whole.

The sex distribution (3 female : 2 male) follows that of the country as a whole.

An attempt has been made to measure the load on the community by assessing the work done by general practitioners, district nurses, hospitals, and the patients' relatives and families.

A study of the age of onset shows a tendency for males to predominate up to age 40, with a striking excess of females in the sixth and seventh decades. There is a tendency for male predominance to reappear in old age.

Less than 50% of the patients present with the classical symptoms of thirst, polyuria, and weight loss.

Seventy-seven per cent. of the patients attend a general medical or diabetic clinic.

The problems of the home are mainly those of insulin administration and intercurrent illness: 80% of male and 62% of female diabetics requiring insulin administer it themselves.

For those unable to give their own insulin the wife usually provides the necessary assistance for the helpless husband; to the dependent female patient help is most often given by the daughter, district nurse, or husband.

More than three-fourths of the patients attending the clinics had no illness requiring rest in bed during the course of a year. The average number of days spent in bed with sickness was 4.9 in the year 1954 and 4.1 in 1955. The female patient tended to have rather more illness than the male. Respiratory illnesses predominated in the home. Diabetes and its complications were responsible for only 15.5% and 13.8% of all days in bed in the two years investigated.

The district nurse visits 15.1% of all diabetic patients in a year and is responsible for daily insulin administration in about 6.4% of all patients.

The general practitioner sees the average diabetic patient 10 times in the year. He is consulted by 91% at some time during the year. Home consultations predominate for women and surgery consultations for men.

The commonest reason for admission to hospital was initial stabilization: 42.5% of all admissions were for reasons not connected with the diabetes.

It is suggested that one medical bed for every 100 of the diabetic population is required in this area.

Cardiovascular disease was responsible for death in more than half of 79 patients followed up. The disease in Cornwall appears to affect the expectation of life in the female but not in the male. The age at death for both sexes was 67.1 years.

I am indebted to the 48 general practitioners who co-operated in this survey; to Miss A. White, who provided the information on the work of the district nurses; and to the hospital secretaries and matrons for figures of diabetic admissions.

APPENDIX : QUESTIONNAIRE SENT TO G.P.s

Patient's initials.....
 Present age if known .....
 Sex .....
 Whether attending a clinic:
 (a) Diabetic clinic .....
 (b) General medical .....
 (c) Other .....
 Whether at present a hospital in-patient:
 (a) General medical ward .....
 (b) Chest hospital .....
 (c) Geriatric service .....
 (d) Mental hospital .....
 (e) Other .....
 If no hospital contacts, diagnosis based on
 (a) Previous hospital investigations .....
 (b) Symptoms and urinary findings .....
 (c) Blood-sugar examination .....
 During 1954 how many visits did this patient require:
 (a) Home visits .....
 (b) Surgery attendances .....
 Is this patient at present having insulin ? .....
 Number of list patients surveyed .....
 Number of diabetics found .....
 Private diabetic patients under your care .....

Remarks:

If information not available enter O. Signature: _____

REFERENCES

- General Register Office (1956). *Studies on Medical and Population Subjects*, No. 9. H.M.S.O., London.
 Joslin, E. P., Root, H. F., White, P., and Marble, A. (1952). *The Treatment of Diabetes Mellitus*, 9th ed. London.
 Registrar-General (1954). *Quarterly Returns for England and Wales*. H.M.S.O., London.
 — (1955). *Statistical Review of England and Wales for 1954*. H.M.S.O., London.

OUTBREAK OF STAPHYLOCOCCAL INFECTION IN A SURGICAL WARD

BY

R. A. SHOOTER, M.D.

J. D. GRIFFITHS, F.R.C.S.

JOSEPHINE COOK, M.B., D.C.H.

AND

R. E. O. WILLIAMS, M.D.

From the Departments of Bacteriology and Surgery and the Children's Department, St. Bartholomew's Hospital, and the Central Public Health Laboratory, Colindale

Outbreaks of staphylococcal infection in surgical wards are not very uncommon, but they are rarely recorded in print. This is partly because opportunities are not often available for extensive investigation, and in consequence the source of infection can rarely be traced. Nevertheless, the paucity of published records makes more difficult the study of new outbreaks as they occur.

It so happened that the outbreak described in this paper could be investigated in some detail, and, perhaps for this reason, we have been able to indicate a possible way in which infection may have been spread.

Clinical Features

The surgical unit concerned consisted of an operating theatre and a male and a female ward, each containing

25 beds. The wards were large and airy and the beds were spaced approximately 6 feet (1.8 m.) apart. Before May, 1956, sporadic cases of staphylococcal infection had occurred in both wards. On May 2 a wound-book was begun in order that a more reliable measure of the incidence of sepsis could be obtained. Against each patient's name was entered the name of the surgeon, details of the operation, and the condition of the patient's wound during convalescence. Wounds in which pus could be seen were regarded as septic; patients whose operations were through infected areas were excluded in assessing the sepsis rate.

Between May 2 and June 25, 86 clean operations were performed: three of the patients died post-operatively from causes unrelated to sepsis; of the remaining 83 patients, 24 developed sepsis in their wounds (Table I). Most of this

TABLE I.—Sepsis Developing in Clean Wounds, May 2 to June 25, 1956

	No. of Operations	No. Infected	Other Staphylococcal Infections
May 2-6 ..	7	4	
" 7-11 ..	7	4	
" 12-16 ..	8	2	1 chest infection
" 17-21 ..	3	0	
" 22-26 ..	10	0	
" 27-31 ..	12	3	
June 1-5 ..	7	2	1 urinary "
" 6-10 ..	4	3	
" 11-15 ..	9	3	1 chest "
" 16-20 ..	9	2	2 chest, 1 urinary infection, 2 enteritis
" 21-25 ..	7	1	1 chest, 1 urinary infection, 1 enteritis

sepsis was in the male ward, where 18 out of 43 patients became infected as compared with 6 out of 34 women. Six children brought from the children's unit in another part of the hospital to the operating theatre and returned after operation had no septic complications. *Staphylococcus pyogenes* was recovered in culture from 17 of the infected wounds. From two coliform bacilli were grown, and from five no swabs were taken.

Seven of the male patients developed staphylococcal infection of sites other than the operation wound, including five chest infections and three urinary infections. Finally, by June 25 three patients were found to have staphylococcal enteritis, and, because of this and the number of other infections, it was decided to stop operating, and the male ward was closed.

In addition, the male ward contained six patients operated on before May who, before or during May, became infected with staphylococci. Of these six, one had a pelvic abscess, two had infected wounds, two had urinary infections, and three had chest infections. One of these last patients developed an abscess involving most of the upper lobe of the left lung.

Staphylococci Responsible for Infection

The staphylococci isolated from infected wounds were all of a strain of *Staph. pyogenes* resistant to penicillin, streptomycin, and tetracycline, but at first sensitive to chloramphenicol and erythromycin. Later, as a result of treatment, variants resistant to chloramphenicol or erythromycin appeared. Twelve of the 17 staphylococci isolated from lesions were typed by the bacteriophage method and all were found to belong to one type, with the phage-sensitivity pattern 47/53/75/77. Some strains were lysed only by phages 53, 75, and 77, and some by 75 and 77 alone, but all were considered to belong to the same type. There was no correlation between these minor variations in phage-sensitivity pattern and the change in antibiotic sensitivity. Staphylococci that were not tested, but had the same sensitivity to antibiotics, were considered to belong to the same type in view of our failure at that time to find any strain with the same characteristics but belonging to a different phage type. With one exception, all the staphylococci isolated from patients with infective complications belonged to the epidemic type.