BRITISH MEDICAL JOURNAL

LONDON SATURDAY MARCH 9 1957

THE ROLE OF VALVULAR INCOMPETENCE IN HEART FAILURE*

BY

J. McMICHAEL, M.D., F.R.C.P.

Professor of Medicine, Postgraduate Medical School of London, Ducane Road, London, W.12

AND

J. P. SHILLINGFORD, M.D., M.R.C.P.

Lecturer in Medicine, Postgraduate Medical School of London, Ducane Road, London, W.12; Member of External Staff, Medical Research Council

Within fifty years of Laennec's introduction of the stethoscope a practical understanding of the meaning of cardiac sounds and murmurs had been reached and auscultation held a pre-eminent place in cardiac diagnosis. Then early in this century Mackenzie, having made by his polygraph the most spectacular advances in clinical analysis of cardiac irregularities, began to preach his doctrine of the all-important myocardium to the detriment of interest in the meaning of murmurs. His pupil, Lewis, perpetuated this policy, and the authoritative position of these two men discouraged further auscultatory studies. Since 1945 phonocardiography has become accepted as a diagnostic technique, and, by focusing interest on the auscultatory signs of valvular disease, has led to renewed interest in an old and important subject.

As the left side of the heart is more sheltered from our modern instrumental methods it is better to begin this discussion on the place of valvular incompetence in heart failure on the right side, where observation can be made with greater ease and frequency.

Historical Note on Tricuspid Incompetence

In 1836 Benson noted accurately some characters of the systolic venous pulsation in the neck in a case of tricuspid incompetence proved at necropsy. King (1837), of Guy's Hospital, expressed the view that "on occasions of most copious influx [the right ventricle] becomes distended; upon which the curtains of the tricuspid valve are drawn aside, an aperture of reflux is produced, and the force of the ventricle is diverted from the pulmonary circulation." He termed this the safety-valve action of the tricuspid valve. Rosenbach (1878) recorded that avulsion of the tricuspid valve in the dog was followed by a loud systolic murmur. Byrom Bramwell (1884) was able to give a nearly perfect clinical account and haemodynamic interpretation of tricuspid incompetence, and pointed out that "the tricuspid sphincter is thinner and weaker than the muscular wall of the left ventricle, and relative and muscular incompetence are therefore more

readily produced at the tricuspid than at the mitral orifice. Relative incompetence secondary to an obstruction either in the lungs or at the mitral orifice is extremely common . . . it may be a temporary and curable condition."

James Mackenzie (1908), though recognizing the rarity of organic affections of the tricuspid valve, thought that functional tricuspid incompetence was extremely common, "so common indeed that I am inclined to look upon the valves as being unable to close the orifice properly." He showed in jugular phlebograms the stages of development of the systolic venous wave of tricuspid incompetence, though in later years neither he nor Lewis paid much attention to the condition. Because of the confusion of tricuspid with other systolic murmurs more recent French writers, such as Laubry and Routier (1928) and Soulié *et al.* (1952), have been hesitant in recognizing the importance and frequency of tricuspid incompetence. Fishberg (1937) took a somewhat similar view of the difficulties of diagnosis and regarded systolic pulsation of the liver as the only unequivocal sign of tricuspid incompetence.

Methods of Study

Right Atrial Pulse

Gross incompetence of the tricuspid valve has long been associated with positive systolic pulsation in the jugular veins; detection of lesser degrees of tricuspid incompetence, however, has only recently been made possible by the analysis of right atrial pressure pulses recorded through a cardiac catheter (Korner and Shillingford, 1954). The normal right atrial pulse consists first of the "a" wave, which occurs at the time of right atrial systole. Then towards the end of atrial diastole ventricular contraction pulls the closed valvular diaphragm towards the left, and, as the atrium is anchored by the superior and inferior venae cavae, it becomes elongated and a further drop in pressure in the right atrium (the X descent of Mackenzie) occurs. With continuing inflow into the atria from the great veins the atrial pressure again rises (V ascent) until, with relaxation of the ventricles, the tricuspid valve opens and blood passes from the atrium to the ventricle. This again

*The substance of this paper formed the Sydney Watson Smith Lecture given by one of us (J. McM.) to the Royal College of Physicians of Edinburgh on November 28, 1956.

produces a fall in atrial pressure (the Y descent). Normally the X descent is much greater than the Y. In atrial fibrillation without tricuspid regurgitation the same pattern is found but without the "a" wave (Fig. 1).

FIG. 1.—Right atrial pulse in patients with sinus rhythm and atrial fibrillation. The dotted line shows the point at which the X descent begins.

As tricuspid incompetence develops the normal fall in pressure in the right atrium during ventricular systole is counteracted by the blood regurgitating through the incompetent valve. A higher pressure than normal is then built up in the right atrium, which in turn falls more rapidly when the tricuspid valve opens. With the development of tricuspid regurgitation, therefore, the X descent becomes less and the Y descent greater, and as the regurgitation becomes gross the X descent is obliterated and is replaced by the familiar positive systolic pulsation (Fig. 2).

FIG. 2.—Right atrial pressure curves in sinus rhythm and in atrial fibrillation. Top line: normal X descent. Middle line: impaired X descent. Bottom line: X descent replaced by a positive systolic wave.

Some modern writers are still impressed by the difficulty of differentiating the jugular pulse of atrial fibrillation from that of tricuspid incompetence (Scherf and Boyd, 1948; Wood, 1956). Daley *et al.* (1955) claimed that the experimental production of atrial fibrillation in dogs in fact produced mitral incompetence. Using a somewhat similar technique, Friedman *et al.* (1956), however, failed to confirm their findings. The tracings in

Fig. 1 show quite clearly that a true X descent can occur in the presence of atrial fibrillation. In pathological conditions of the heart it is apparent that atrial fibrillation and tricuspid incompetence may occur together not infrequently; no doubt this association has added to the confusion. Any doubts should be readily resolved by observing the ease with which the poor descent of the base pattern is converted to a positive systolic pulse by mild exercise or even by deep inspiration.

Blood Flow in Right Atrium

The direction of blood flow in the right atrium and great vessels from moment to moment during the cardiac cycle can be demonstrated by using a double-lumen cardiac catheter as a Pitot flow-meter (Müller and Shillingford, 1955). The catheter is placed with the end hole in the right atrium and the side hole in the superior vena cava. The other end of the catheter is then connected to a differential manometer and the recording apparatus, so arranged that if the pressures in the side hole and end hole are the same there is no deflection; if the pressure in the end hole is greater than in the side hole the instrument will record downwards, while if it is higher in the side than in the end the deflection will be upwards. In this way direction of flow can be indicated. Fig. 3 shows a flow curve between

FIG. 3.—Blood flow between the superior vena cava and right atrium in patients with and without tricuspid incompetence.

the superior vena cava and right atrium in a normal patient and in one with tricuspid incompetence. When tricuspid incompetence occurs the normal forward flow during ventricular systole and descent of the base of the heart is replaced by a back-flow. Although this method shows the presence of a regurgitant flow it does not accurately measure the amount of blood passing back.

Estimation of Valvular Incompetence by Dye Dilution Curves

The quantitative estimation of the part played by valvular incompetence in cardiac failure has been made possible by the recent observation that indicator dilution curves are altered in a specific way by the presence of valvular insufficiency (Korner and Shillingford, 1955, 1956). Stewart (1894) and later Hamilton *et al.* (1932) showed that if an indicator substance—for example, Evans blue—was injected into the venous side of the circulation and collected on the arterial side the dye particles would spread out during their passage through the heart and vessels and produce a time-concentration curve at the point of sampling. They pointed out that from this curve it was possible to measure not only the cardiac output but also the volume of blood through which the dye had passed. Korner and Shillingford (1955) have demonstrated that for a given calculated cardiac output and volume the spread of the dye particles is constant, but that the addition of valvular incompetence causes an increase in the spread of the particles quite apart from that produced by variations in the cardiac output and cardio-

pulmonary volume. This effect can be used to compute the amount of regurgitant flow, and, tested in a model circulation, has been found to be accurate, with a maximum error of 15%.

This new approach to the problem of valvular incompetence in cardiac failure has enabled us not only to recognize valvular incompetence of lesser degree than was formerly possible, but also accurately to assess the part it plays in the haemodynamics of cardiac failure.

Functional Tricuspid Incompetence in Congestive Cardiac Failure

Functional tricuspid incompetence has long been recognized as a frequent late development in congestive cardiac failure. A recent analysis (Korner and Shillingford, 1955) of the right atrial pressure pulse in a series of 50 patients with various forms of heart disease with and without congestive cardiac failure has revealed a remarkably close relationship between the level of the mean venous pressure and the development of functional tricuspid incompetence. As the venous pressure rises so the normal X descent becomes impaired and takes on the form of a positive systolic pulsation. The critical level of venous pressure at which tricuspid incompetence becomes recognizable is approximately 8 mm. of mercury above the normal mean zero pressure. Development of functional tricuspid incompetence takes place about this level of venous pressure in many aetiological varieties of heart failure, and when the venous pressure becomes more grossly raised functional tricuspid incompetence is even more obvious and beyond argument (Fig. 4).

FIG. 4.—Relationship between mean venous pressure and the form of the atrial pulse in 50 cardiac patients.

FIG. 5.—Forward output and phasic regurgitant flow estimated by the indicator dilution method in 12 patients with varying degrees of congestive cardiac failure.

Thus a considerable number of cardiac patients with grossly raised venous pressure also have functional tricuspid incompetence. The only exceptions to this are found in constrictive pericarditis and conditions with a very rapid venous inflow (anaemia, early thyrotoxicosis). The importance of these clinical phenomena has been overlooked, probably because of the difficulty of timing the pulse waves in the neck at the bedside, especially when the heart is beating rapidly. Only by graphic recording methods is it possible accurately to assess the form of the venous pulse.

The magnitude of the back-flow through the functionally incompetent tricuspid valve in cardiac failure can be con-

siderable. As the right ventricle and the tricuspid valve dilate, the aperture of reflux may become as great as, or greater than, the pulmonary valve orifice when fully open (Lottenbach and Shillingford, in press). As the mean right atrial pressure is considerably lower than the mean pulmonary arterial resistance to back-flow the oscillating back-flow may easily exceed the forward flow into the pulmonary vessels. Measurement of this back-flow by dye-dilution methods shows it in some cases to reach 6 or 7 litres a minute, while the forward cardiac output is only 2 or 3 litres a minute (Fig. 5). In other words, the stroke regurgitant flow oscillating backwards and forwards through the valve would be of the order of 60–70 ml. back and 80–100 ml. forward at a pulse rate of 100 beats a minute. This observation of a total high right ventricular output in the presence of heart failure and a low forward cardiac output can be appreciated by palpating the excessively large right ventricular pulsations even in patients in advanced congestive cardiac failure.

The physical signs correlate with the presence of this backward flow in congestive failure. The systolic wave in the venous pulse is apparent on careful examination; a systolic murmur heard best at the lower end and just to the left of the sternum is present in most cases; and in the same area, when the phasic regurgitant flow is great, there may be a low-pitched early diastolic murmur; this murmur is probably associated with the rapid inflow of blood from the right atrium to ventricle during this phase of the cardiac cycle, and does not necessarily denote an organic change in the valve itself. It must be remembered that diastolic inflow to the right ventricle under these circumstances equals the sum of backward and forward flow and may therefore be very large. Liver pulsation becomes apparent to nearly all clinical observers if the regurgitant stroke output exceeds 40 ml.

The regurgitant flow through an incompetent valve is not a fixed reflux but is varying from minute to minute under the influence of rest and exercise, emotion, respiration. The effect of these factors may be studied clinically or by means of the newer techniques for quantitative measurement of regurgitant flow.

Influence of Changing Inflow: Respiration: Rest: Exercise

Functional tricuspid incompetence varies with the degree of distension of the right ventricle. It may be seen to vary from beat to beat; for example, under the influence of respiration a deep inspiration may produce or intensify the tricuspid systolic murmur and bring out a strong systolic venous pulse in the jugular vein (Fig. 6). In atrial fibrillation a long diastolic pause may permit an increase in right

FIG. 6.—Effect of deep inspiration on jugular pulse and tricuspid systolic murmur in a patient with congestive heart failure.

FIG. 7.—Effect of exercise and rest on jugular venous pulse in a patient with chronic rheumatic heart disease and early congestive cardiac failure. A, before exercise. B, after three minutes' mild exercise, showing development of positive systolic pulsation and increase of functional tricuspid systolic murmur (phonocardiographic record taken at lower end of sternum). C, after three minutes' rest. D, after six minutes' rest, showing return to X descent pattern and diminution in intensity of tricuspid murmur.

ventricular distension which makes the valve more incompetent; and as a result tricuspid systolic murmurs may diminish in intensity after a short diastole and increase again when the pause is long between ventricular beats.

Rest and exercise have a profound effect on the amount of regurgitant flow through the tricuspid valve (Figs. 7 and 8). Fig. 9 shows two examples of this in patients with

chronic rheumatic heart disease and mild or moderate venous congestion. It will be seen in the second patient (M. B.) that the regurgitant flow on the first dye injection was 2.2 litres a minute, while the forward output was 3.7 l./min. The mean jugular venous pressure was 10 mm. Hg. On resting quietly for 20 minutes the venous pressure had fallen to 7 mm. Hg and the regurgitant flow to 0.5 l./min., but the forward cardiac output had risen to 4.6 l./min. Gentle exercise by moving the legs for two minutes reversed this process and diminished the forward cardiac output to 4.3 l./min., with a rise in the regurgitant flow to 3.6 l./min. and the jugular venous pressure (Fig. 6). This increase in regurgitant flow is confirmed by measurements by the flowmeter (Fig. 8).

It will thus be seen that exercise in congestive heart

failure may increase the total right ventricular output but decrease the forward output of the heart, while rest has the opposite effect.

Organic Tricuspid Incompetence

Although functional tricuspid incompetence is common, organic lesions of the valve are comparatively rare. Rheumatic disease affects the tricuspid valve in 10 to 15% of all our cases of chronic rheumatic heart disease coming to necropsy. Nearly all these have more incompetence than stenosis, valve-cusp fusion being limited to the margins. In the earlier stages it is possible to have all the signs of tricuspid incompetence without much dilatation of the right ventricle. In fact, the right ventricle may be able to carry the additional load of mild regurgitant flow for many years without signs of failing; the patient during this time has a raised venous pressure and may even have cardiac cirrhosis, ascites, and oedema.

Influence on General Haemodynamics of Cardiac Failure

The study of cardiac output in heart disease has revealed that the resting output often remains in the normal range until the last bedridden stage, when it falls towards half the normal; at this last stage the venous pressure is grossly raised. Venesection in these patients may be followed by a rise in cardiac output, while exercise, producing elevation of the venous filling pressure, often engenders a fall in cardiac output.

More than a decade ago McMichael and Sharpey-Schafer (1944) compared these changes with the measurements made by Starling and his school on the isolated dog heart. An elevation of venous filling pressure which normally increases the cardiac output may, if pushed to extremes, have the reverse effect of bringing about a fall in output resulted from some breakdown of the cardiac contractile mechanism, because it was at this stage of human heart failure that favourable digitalis and venesection responses were often encountered.

Sarnoff and Berglund (1954) have made extensive studies of the conditions under which Starling's law becomes applicable. They have demonstrated its general validity when applied to a single ventricle—for example, the left—and have reconstructed curves which show the rising cardiac output curve flattening off in a plateau with continuing elevation of venous pressure, but a falling output response was seen only when atrio-ventricular valve incompetence was induced by overstretching.

If we accept this finding and link it to our own more recent observations the following tentative suggestions seem permissible:

1. The falling cardiac output with gross venous congestion may almost entirely be accounted for by reduction of forward cardiac output consequent on the development of tricuspid incompetence.
2. The improvement in cardiac output after venesection is simply the result of a decreased filling pressure which diminishes right ventricular dilatation; the tricuspid valve becomes more competent and back-flow is reduced.
3. Falling output and elevation of venous pressure on exercise—for example, like that noted by Hickam and Cargill (1948)—may result from increased distension of the right ventricle increasing the tricuspid leak, while the reverse occurs with rest.

4. There is no necessity to postulate a metabolic breakdown of the myocardium at this stage. Such a breakdown invoked by our earlier hypothesis was subject to the criticism that it would have entailed "vicious circle" consequences, increasing venous pressure leading to further and further degrees of myocardial breakdown.

FIG. 8.—Effect of exercise on blood flow between superior vena cava and right atrium in a patient with chronic rheumatic heart disease and early congestive cardiac failure. The development of a regurgitant flow through the tricuspid valve on exercise is shown by the deep downward deflection in systole (measured by differential manometer).

FIG. 9.—Effect of rest and exercise on right ventricular output in congestive cardiac failure with functional tricuspid incompetence (see text).

A further comment may be made in relation to the flow-work of the right ventricle. In the presence of tricuspid incompetence the bidirectional output of this chamber at each systole may be nearly double the normal. The falling forward output may thus reduce only the work of the left ventricle, while the right ventricle is maintaining a greatly increased output. No doubt this local overload imposed by an incompetent tricuspid valve helps to produce the final right ventricular breakdown. It is amazing sometimes to see how long it may be tolerated—six or seven years—before final dissolution. We must think again about low cardiac output states. It may be only the forward output which is reduced, as an incompetent valve may actually increase the total output of the affected chamber. Older subdivisions of heart failure into "high" and "low" output types will give way to the concept that nearly all failing hearts have been overworking for a very long time against the mechanical embarrassments of diseased valves or increased pressure loads.

Valvular Incompetence and Failure of the Left Ventricle

Owing to the technical difficulties encountered in studying left atrial and ventricular pressures the information on left heart failure is not so complete as that on the right. It is likely, too, that there is a considerable basic difference in the haemodynamics of the two sides in cardiac failure. Whereas the tricuspid valve probably only just closes in normal subjects and is readily made incompetent by dilatation of the right ventricle, the mitral valve attached by its chordae to the thicker-walled left ventricle does not seem to become functionally incompetent so easily. At the same time the venous reservoir of the left atrium and lung bed is considerably less than that of the right atrium, caval veins, and their large valveless tributaries. This results in a difference in pressure-volume relationships in the veno-atrial systems on the right and left sides which tends to limit the amount of blood regurgitating through the mitral valve compared with that which may flow back through the tricuspid valve.

Recent work on organic mitral incompetence has shown that the to-and-fro regurgitant flow in established cases may be 2 or 3 litres a minute, which, added to the forward flow, makes a high total output of the left ventricle. The regurgitant flow is not constant, but varies from moment to moment. Any factor tending to lower the peripheral resistance and thus lessen the obstruction to the aortic outflow will at the same time reduce the back flow through the mitral valve relative to the forward flow through the aortic valve: this occurs in exercise. The administration of amyl nitrite causes the forward output to increase considerably, while the regurgitant flow becomes less, a finding which may also be suspected clinically by noting the diminution in the intensity of the murmur of mitral incompetence. The administration of a vasoconstrictor substance, on the other hand, produces an increase in the regurgitant flow. Rest and exercise may thus have opposite effects on the volume of regurgitant flow through the tricuspid and mitral valves.

Aortic Incompetence

The haemodynamics of aortic incompetence present several important differences from those of the atrio-ventricular valves. The regurgitant flow in the latter takes place during systole and is dependent on the full force of ventricular contraction during the part of the cardiac cycle. In aortic incompetence the regurgitant flow depends on the potential energy built up by distension of the aorta and its branches during systole; the amount of this energy developed depends on the stroke volume of the left ventricle, and on resistance and elasticity of the arterial system. In the first part of diastole this energy is rapidly expended forwards through the peripheral arteries and backwards through the incompetent aortic valve. As in mitral incompetence, therefore, lowering peripheral resistance re-

duces the regurgitant flow through the valve, and so does loss of elasticity in the aorta. The peripheral resistance in aortic incompetence is in fact reduced by physiological reactions producing vasodilatation and thus reducing the back-flow. The mechanism of this is not clear, but may well be a reflex initiated in the pressor-receptor areas in the arch of the aorta and carotid sinus. This vasodilatation also has the effect of accommodating a high stroke output of the left ventricle over and above that due to the additional load of regurgitant flow.

Without detracting from the importance of the myocardium it is clear that the volumes of back-flow involved in valvular incompetence are often very large, and constitute a factor of major importance in the development of the syndromes of heart failure. While measurements of valvular regurgitation on the right side of the heart may be made with relative ease and safety by a variety of complementary methods, the techniques applicable to the problems of the left side are more restricted and difficult of application. Observations made so far suggest that the anatomical and haemodynamic factors differ on the left and right sides and indicate somewhat different mechanisms in the process of final cardiac breakdown.

Summary

The part played by valvular incompetence in cardiac failure is discussed.

Functional tricuspid incompetence often occurs in congestive cardiac failure when the venous pressure is raised above 8 mm. of mercury. This regurgitation can be demonstrated by the form of the venous pulse, the tricuspid systolic murmur, and special techniques to demonstrate direction and volume of flow between the right ventricle and atrium during the cardiac cycle.

The development of tricuspid incompetence causes an increase in the total output of the right ventricle at the expense of the forward cardiac output, and even in advanced clinical congestive failure with low cardiac output the right ventricular output remains high.

Exercise increases the functional tricuspid incompetence, with a corresponding fall in forward cardiac output, and probably explains both the phenomena of fixed or falling cardiac outputs on exercise in congestive cardiac failure and the breakdown of Starling's law with a rising venous pressure.

The haemodynamics associated with mitral and aortic insufficiency are discussed and are shown to differ considerably from those involving the tricuspid valve.

REFERENCES

- Benson, C. (1836). *Dublin J. med. chem. Sci.*, 8, 324.
- Bramwell, B. (1884). *Diseases of the Heart and Thoracic Aorta*. Edinburgh.
- Daley, R., McMillan, I. K. R., and Gorlin, R. (1955). *Lancet*, 2, 18.
- Fishberg, A. M. (1937). *Heart Failure*. Kimpton, London.
- Friedman, B., Daily, W. M., and Wilson, R. H. (1956). *Circulat. Res.*, 4, 33.
- Hamilton, W. F., Moore, J. W., Kinsman, J. M., and Spurling, R. G. (1932). *Amer. J. Physiol.*, 99, 534.
- Hickam, J. B., and Cargill, W. H. (1948). *J. clin. Invest.*, 27, 10.
- King, T. W. (1837). *Guy's Hosp. Rep.*, 2, 104.
- Korner, P., and Shillingford, J. (1954). *Brit. Heart J.*, 16, 447.
- (1955). *Clin. Sci.*, 14, 553.
- (1956). *Ibid.*, 15, 417.
- Laubry, Ch., and Routier, D. (1928). *Bull. méd. (Paris)*, 42, 91.
- Mackenzie, J. (1908). *Diseases of the Heart*. Oxford Univ. Press, London.
- McMichael, J., and Sharpey-Schafer, E. P. (1944). *Quart. J. Med.*, 37, 123.
- Müller, O., and Shillingford, J. (1955). *Brit. Heart J.*, 17, 163.
- Rosenbach, O. (1878). *Arch. exp. Path. Pharmacol.*, 9, 1.
- Sarnoff, S. J., and Berglund, E. (1954). *Circulation*, 9, 706.
- Scherf, D., and Boyd, L. J. (1948). *Cardiovascular Diseases*. Heinemann, London.
- Soulé, P., Rouvrain, Y., and Di Matteo, J. (1952). *Bull. Soc. méd. Hôp. Paris*, 68, 332.
- Stewart, G. N. (1894). *J. Physiol. (Lond.)*, 15, 31.
- Wood, P. (1956). *Diseases of the Heart and Circulation*. Eyre and Spottiswoode, London.