EMERGENCIES IN GENERAL PRACTICE

PAROXYSMAL TACHYCARDIA

BY

GRAHAM HAYWARD, M.D., F.R.C.P.

Physician, National Heart Hospital

Paroxysmal tachycardia is a condition characterized by attacks of rapid beating of the heart; the onset is sudden, the duration is usually short, but occasionally lasts days or weeks, and the rate of the heart is between 140 and 240 a minute. There are four different types of paroxysm: (1) supraventricular or auricular tachycardia, due to a rapid regular succession of ectopic beats arising from a focus in the auricle or auriculoventricular node; (2) ventricular tachycardia, with an irritable focus in the ventricle and a rapid succession of ventricular ectopic beats; (3) auricular fibrillation, with rapid irregular response of the ventricle; and (4) auricular flutter, usually with 2:1 block so that the ventricle responds to every second stimulus reaching it from the auricle. The recognition of the different types of paroxysm is important, as their prognosis and treatment vary. The nature of the attack may often be decided by ordinary clinical methods of examination, but electrocardiographic confirmation is advisable when possible, particularly in long-lasting attacks, in attacks occurring in patients with evidence of organic heart disease, or if the attacks began after the age of 50, when they may be a manifestation of coronary artery disease. The commonest are supraventricular (auricular) tachycardia (60% of cases) and auricular fibrillation (30%), while auricular flutter (6%) and ventricular tachycardia (4%) are infrequent.*

Aetiology

Paroxysmal tachycardia occurs most commonly between the ages of 20 and 40, in patients with normal hearts or in association with rheumatic heart disease or hyperthyroidism. The onset of attacks in older patients is usually due to the presence of ischaemic or hypertensive heart disease. Attacks are rare in infancy or childhood, are usually supraventricular in type, but they are serious when they do occur, as there may be an unusually rapid ventricular rate of 250–300 a minute.

Nearly two-thirds of the patients with paroxysms of supraventricular tachycardia have no organic heart disease; the remainder have rheumatic or ischaemic heart disease. In contrast, over 80% of patients with ventricular tachycardia have serious heart disease, either recent cardiac infarction or severe congestive failure. Paroxysmal auricular fibrillation is commonly caused by hyperthyroidism or rheumatic or ischaemic heart disease; less than one-sixth of the cases occur in patients without evidence of heart disease. In this last group the attacks are sometimes precipitated by severe acute infections. Only one-tenth of the patients with paroxysmal flutter have normal hearts; in most cases there are signs of rheumatic, hypertensive, and ischaemic heart disease or occasionally of hyperthyroidism. Paroxysmal tachycardia, usually auricular flutter or fibrillation, may occur as an early manifestation of a bronchial carcinoma; it is then caused by irritation or infiltration of the auricular muscle.

Symptoms

The chief symptom of an attack is palpitation, the patient being conscious of the exact moment at which the heart

rate increased. The end of the attack may be equally abrupt, although not infrequently the attack may seem to pass away gradually or is found to have stopped when the patient awakens from sleep. Other symptoms at the beginning of the attacks are variable and depend to a larger extent on the rate of the heart and the mental make-up of the patient. Dizziness, a feeling of tightness in the chest, faintness, or actual syncope may occur, and the symptoms of anxiety may be added. Later symptoms in the attacks depend largely on the health of the heart muscle and the duration of the paroxysm. Even in a normal heart a long attack may cause alarming symptoms and signs of congestive failure, although these disappear rapidly when the attack stops. Anginal pain during an attack does not mean that cardiac infarction has occurred, and it should be remembered that the tachycardia itself may cause depression of the ST segment or flattening or inversion of the T wave in the electrocardiogram which may persist for several days after the attack has stopped. The rate of the heart during a paroxysm is between 140 and 240 a minute, in most cases between 160 and 200, and remains constant. The rhythm, except in attacks of auricular fibrillation, is regular.

It is often impossible to assess the state of the heart by examination during an attack: the tachycardia itself may make the heart appear to be enlarged on clinical examination; heart murmurs cannot be heard or timed owing to the rapid rate; and the blood pressure commonly falls because the shortened diastolic pause allows insufficient time for proper ventricular filling and the cardiac output falls. It is only by examination between attacks that it is possible to decide whether or not heart disease is present.

Diagnosis

The essential feature in making a diagnosis of paroxysmal tachycardia is the abrupt onset of the attacks. The change from normal rhythm occurs within one heart beat and the patient is conscious of the exact moment at which an attack starts. If the change in heart rate occurs more slowly, over the course of 15–30 seconds or longer, a diagnosis of paroxysmal tachycardia should not be made. It is helpful also to get the patient to tap out with his finger the rate and rhythm of the heart during an attack, as slow but forceful beating of the heart, described as palpitation, may sometimes be confused with true paroxysmal tachycardia.

The next stage in diagnosis is to establish the nature of the paroxysm, for treatment may be more effectively planned if this is known. Paroxysmal auricular fibrillation may be distinguished by the rapid irregular beating of the heart, whereas in the other varieties the rhythm is regular, except in some paroxysms of auricular flutter, when there may be a rapid variation in the degree of block. The clinical features which may enable differentiation of the regular tachycardias are the age at the onset of the attack, the rate, the presence or absence of underlying heart disease, the response to carotid sinus pressure, and inspection of the neck veins.

Age.—Attacks starting under the age of 40 are most commonly due to supraventricular tachycardia, while auricular flutter or ventricular tachycardia are commoner over the age of 50.

Rate.—In supraventricular tachycardia the rate is between 160 and 220, commonly 180 to 200, a minute and does not

*Campbell, M., *Lancet*, 1947, 2, 641.

vary from minute to minute during the attacks. In ventricular tachycardia the rate is within the same limits but is less constant, the heart sometimes quickening or slowing within this range. In auricular flutter there is usually 2:1 block and the ventricular rate is usually 140–150 a minute, slower than in either of the other varieties of regular tachycardia.

Underlying Heart Disease.—Supraventricular tachycardia occurs most commonly in patients without any organic heart disease, and a paroxysm in such a patient is most likely to be due to this cause. Ventricular tachycardia is rare except in patients with obvious and serious heart disease such as congestive failure or recent cardiac infarction, and an attack in such a patient should be presumed to be of this nature unless cardiographic evidence to the contrary is available. In auricular flutter heart disease is usually present, but this is generally less serious than that associated with ventricular tachycardia, and the slower rate should enable the differentiation to be made.

Response to Carotid Sinus Pressure.—In supraventricular tachycardia carotid sinus pressure will either stop the attack abruptly or have no effect on the rate. In auricular flutter it may cause temporary irregularity of the heart or sudden slowing of the heart, but on release of pressure the heart returns to its original rate of beating. It has no effect at all on the rate in ventricular tachycardia. The carotid sinus is situated at the bifurcation of the common carotid artery at the level of the upper border of the thyroid cartilage, and pressure should be applied over it for about 15–20 seconds, first on the right side and then on the left (but *not* on both sides simultaneously). The patient should be sitting, with the head supported and the neck muscles relaxed, the head being slightly rotated to the opposite side to that on which pressure is being applied. The manoeuvre is best avoided in the elderly arteriosclerotic patient. In some cases gentle stroking of the carotid sinus is more effective than sustained pressure.

Venous Pulsation in the Neck.—The neck veins should be inspected with the patient lying as flat as is comfortable, so that the veins are distended. In some cases of auricular flutter the rapid flutter waves can be seen at the root of the neck, with a rate of 280–300 a minute.

Complications and Prognosis

The significance of anginal pain and congestive failure has been discussed above. Failure coming on late in a long attack is far less significant than failure which appears soon after the onset. Arterial emboli may be liberated in patients with mitral stenosis, and the fall in cardiac output with an excessively fast rate, coupled with other factors such as anaemia or infection, may be responsible for the occasional occurrence of venous thrombosis and pulmonary embolism.

The immediate prognosis is excellent in nearly all cases, whether or not heart disease is present, and this provides good reason for the confident reassurance which should be given to patients during an attack. The exceptions to this good prognosis are those attacks of paroxysmal ventricular tachycardia which occur after cardiac infarction in patients already severely ill with congestive failure, and the rare attacks in infants when the unusually rapid ventricular rate may cause syncope or heart failure. The long-term prognosis depends on the state of the heart between the attacks. There is no evidence that even frequently recurring paroxysms occurring with a normal heart cause any permanent damage to the heart or shorten life.

Supraventricular Tachycardia

Supraventricular tachycardia is in nearly all cases a benign condition, even though in children or in attacks with a very rapid ventricular rate syncope or congestive failure may be alarming. It is therefore rarely necessary to use powerful

drugs which may produce unpleasant side-effects. In many cases the patient will have himself discovered some way of stopping the attack, such as a change in posture, putting the head between the knees, taking a full inspiration and then attempting to expire forcefully with the glottis closed, drinking iced water, or by self-induced vomiting. If these manoeuvres have been unsuccessful carotid sinus pressure as described previously or eyeball pressure enough to cause pain should be tried.

If the attack persists and the patient is not unduly distressed a sedative such as butobarbitone, 3–4½ gr. (0.2–0.3 g.), should be given; it will be found in many cases that the attack has stopped when the patient wakes. In those cases where the attack persists for longer than 24 hours, or if distressing symptoms or signs of congestive failure have occurred, digitalis is the drug of choice. It should be given as digoxin, 1 mg. by mouth followed by 0.5 mg. six hours later. If a more rapid action is needed the initial dose may be given by slow intravenous injection, provided the patient has not previously been receiving digitalis; but this should rarely be necessary. Quinidine is less effective than digitalis in the treatment of paroxysmal supraventricular tachycardia, but it may be tried, giving 5 gr. (0.32 g.) two-hourly by mouth to a total dose of 30 gr. (2 g.). Drugs which act by causing powerful vagal stimulation will often stop attacks, but they rarely need to be used and the side-effects are unpleasant. The best is methacholine chloride, 20 mg. subcutaneously. A syringe containing atropine sulphate, 1/50 gr. (1.3 mg.), should be prepared so that if the methacholine causes colic, diarrhoea, vomiting, or collapse, the atropine can be injected intravenously immediately. If methacholine fails to stop the attacks carotid-sinus pressure should again be tried, as the sinus may now be more sensitive.

Digitalis is the most useful drug for the treatment of paroxysms in infants, quinidine, carotid-sinus stimulation, and the vagal stimulants such as methacholine being usually ineffective. For babies under 15 lb. (6.8 kg.) in weight tincture of digitalis, 2 min. (0.1 ml.) six-hourly for up to four days, should be given. For older children 0.125 to 0.25 mg. of digoxin initially and repeated in six hours is usually effective.

The prevention of these attacks is often difficult. In those patients in whom emotional disturbances precipitate attacks the regular use of phenobarbitone may abolish or lengthen the interval between them. Quinidine is the best drug for prophylaxis (quinidine sulphate, 5 gr. (0.32 g.) twice or three times a day), and its administration should be continued for several months. Rarely, attacks will recur with such frequency or severity as to incapacitate the patient, and in these cases the use of thiouracil drugs should be considered. If a state of mild hypothyroidism is induced the attacks often stop, and if treatment is carried out for three to six months and then discontinued the attacks may not recur. Methylthiouracil, 400 mg. daily, should be given at first, the dose being gradually reduced to a maintenance dose of 75–150 mg. daily, depending on the response.

Paroxysmal Auricular Fibrillation

Digitalis and quinidine are the two drugs which are useful in paroxysmal auricular fibrillation. If the ventricular rate is rapid (over 200 a minute) or if signs of congestive failure are present, digoxin should be given as described in the previous section. The drug often stops the attacks, and the risk of it turning paroxysmal fibrillation into established fibrillation is small. If no serious symptoms are present, quinidine sulphate should be used (5 gr. (0.32 g.) two-hourly for six doses), and it is the most effective drug for the prevention of paroxysms (5 gr. (0.32 g.) twice or three times a day). In those cases with recurring paroxysms of auricular fibrillation particular attention should be paid to the thyroid gland, as paroxysmal fibrillation may be the first sign of an otherwise masked hyperthyroidism from a small nodular goitre.

Paroxysmal Auricular Flutter

The drug of choice in paroxysmal auricular flutter is digitalis, which if given in full doses will either stop the attacks or convert the flutter to fibrillation, when the ventricular rate is more easily controlled. The ventricular rate is relatively slow in auricular flutter (140–150 a minute), so that there is usually no urgency in treatment. Digitalis leaf, 2 gr. (0.13 g.) three times a day, should be given for two days, then 1 gr. (65 mg.) four times a day until the attack stops or the rhythm changes to auricular fibrillation; the drug should then be stopped and the heart will usually revert to normal rhythm. These large doses may need to be continued until the patient shows signs of digitalis poisoning, such as nausea or vomiting. If digitalis fails to stop the attack it should be continued in a maintenance dose of $\frac{1}{2}$ gr. (32 mg.) three times a day and quinidine sulphate given as described previously. Quinidine slows the auricular rate, and the digitalis is given simultaneously to prevent the ventricle suddenly responding to every auricular stimulus with a rapid rate often of over 250 a minute.

Paroxysmal Ventricular Tachycardia

As over 80% of attacks of paroxysmal ventricular tachycardia occur in patients with serious heart disease, the rapid ventricular rate often causes heart failure or ischaemic pain, and it is particularly important to try to stop the attack as soon as possible. In cases of extreme urgency, procaine amide hydrochloride should be given intravenously, initially 200 mg. slowly, followed by 100 mg. a minute until the attack stops or until a total dose of 1 g. has been given. A careful watch should be kept on the blood pressure during the administration of the drug, as it sometimes causes marked hypotension; whenever possible it is wise to have electrocardiographic control in case the drug causes conduction defects. If there is less urgency it is better to give the procaine amide by mouth, 1 g. (four tablets) initially, followed by 0.5 g. four-hourly for 24 hours. Quinidine is also sometimes effective in the treatment of the attack—5 gr. (0.32 g.) two-hourly to a total of 30 gr. (2 g.)—and it is the best drug for preventing attacks (5 gr. (0.32 g.) three times a day). Digitalis is contraindicated. If there is any doubt about the nature of an attack of paroxysmal tachycardia it is wise not to use digitalis until the exact diagnosis has been established by electrocardiography.

Conclusion

It should again be emphasized that the great majority of cases of paroxysmal tachycardia seen in practice are benign, particularly if the patient is young and there is no previous history of severe heart disease. Simple sedatives alone will stop many attacks, and the temptation to use in rapid succession a number of powerful drugs, which often have unpleasant side-effects, should be resisted.

Next article on Emergencies in General Practice.—
“Sprains of the Limb Joints,” by Mr. J. I. P. James.

Refresher Course Book.—Copies of the second volume of collected articles from the Refresher Course for General Practitioners are still available at 25s. (postage—inland 1s. 6d., overseas 1s.) each. The first volume is now sold out.

Clinical Pathology Book.—“Clinical Pathology in General Practice,” a collection of 39 articles on clinical pathology that appeared in the *Journal* as part of the Refresher Course for General Practitioners, is now available, price 21s. (postage—inland 1s. 3d., overseas 9d.).

Both these volumes are obtainable from the Publishing Manager, B.M.A. House, Tavistock Square, London, W.C.1, or through any bookseller.

Nova et Vetera**DANIEL McNAUGHTON'S SIGNATURE**

On January 20, 1843, Edward Drummond, private secretary to Sir Robert Peel, the Prime Minister, was walking along Whitehall towards Downing Street. Between the Admiralty and the Horse Guards headquarters he was shot in the back by a man who mistook him for Peel. He died on the morning of January 25. His assailant, who had an imaginary grudge against Peel, was arrested and taken to Gardiner's Lane police station, where, according to *The Times* of January 21, 1843, he gave his name as Daniel M'Naughten. The inquest on Drummond was reported in *The Times* of January 27, which used the same spelling.

The depositions of witnesses and others, taken immediately after the crime, and the coroner's inquest report are preserved at the Public Record Office, London (Crim. 1/4). These documents without exception use the form McNaughton. Among them is a statement made before Mr. T. J. Hall, the Bow Street magistrate, by the prisoner on January 21 and signed by him:

This is the only document bearing McNaughton's signature that can be traced, and unfortunately it fails to make clear just how McNaughton spelled his name. An authority at the British Museum is of the opinion that this signature is “Daniel McNaughtun”; but experts at the Public Record Office, while agreeing that the penultimate letter resembles “u” rather than “o,” point out that it does not appear to be formed in the same way as the “u” earlier in the signature. The form McNaughtun does not occur in the following authoritative works, all of which include McNaughton or McNaghtan, sometimes with as many as 16 variants: H. Barber, *British Family Names*, London, 1903; G. F. Black, *Surnames of Scotland*, New York, 1946; M. A. Lower, *Patronymica Britannica*, London, 1860; P. Woulfe, *Irish Names and Surnames*, 1923. It does not occur in the Glasgow registers of births and baptisms for the years 1811 to 1815. Consequently the most likely spelling seems to be McNaughton.

McNaughton's trial at the Old Bailey began on March 3, 1843. The indictment, now in the Old Bailey records, refers to Daniel MacNaughton. Among evidence submitted was a letter from the prisoner to his bank concerning a deposit of £750 made by him. In reprinting this letter, and in fact throughout the trial, *The Times* (March 4 *et seq.*) consistently used the spelling M'Naughten. Evidence given by the prisoner's father, a Glasgow wood turner, showed that Daniel was illegitimate. He had been apprenticed to his father some years previously. After evidence of his insanity had been submitted, the jury found him not guilty of murder, and he was ordered to be detained during Her Majesty's pleasure. The official report of the trial is published in *Central Criminal Court Minutes of Evidence Taken in Shorthand* by Henry Buckler, London, 1843, 17, 721; it uses the spelling M'Naughten.

There was a debate in the House of Lords on March 13, 1843, when the Lord Chancellor (Lord Lyndhurst) promised to obtain the views of the judges on the need for changing the law on crime and insanity. The questions put to the judges and their answers thereto are recorded in the *Journals of the House of Lords*, 1843, 75, 400–402 (June 19), which uses the form M'Naughten.

Registered at Bethlem

After trial Daniel was committed to Newgate and on March 13 he was transferred to Bethlem, where, in the