speculation, for the pulmonary plasmocytosis might just be part of a general tissue infiltration; but this is rather unlikely, since, of all the tissues examined, only marrow, spleen, and lungs were involved by plasma cells. A further complication is the propensity of the lungs for retaining and extracting cells from the blood stream (Osogoe, 1950; Vaughn, 1953), which may have influenced the final histological composition.

JOHN VAUGHN, M.D.,
STANLEY S. RAPHAEL, M.B., B.S.,
Department of Pathology,
Southend General Hospital, Southend-on-Sea, Essex.

REFERENCES

- Bayrd, E. D. (1948). *Blood*, 3, 987.
Custer, R. P. (1935). *Amer. J. med. Sci.*, 189, 507.
Darling, R. C., Parker, F., and Jackson, H. (1936). *Amer. J. Path.*, 12, 1.
Fadum, R. S., and McBirnie, J. E. (1950). *Blood*, 5, 191.
Falconer, E. H., and Leonard, M. E. (1948). *Ann. intern. Med.*, 29, 1115.
Hayhoe, F. G. H., and Smith, D. R. (1951). *J. clin. Path.*, 4, 47.
Macintyre, W. (1850). *Med.-Chir. Soc. Trans.*, 33, 211.
Osogoe, B. (1950). *Anat. Rec.*, 107, 193.
Snapper, I., Turner, L. B., and Moscovitz, H. L. (1953). *Multiple Myeloma*. Grune and Stratton, New York.
Vaughn, J. (1953). *Blood*, 8, 1.
Whitby, L. E. H., and Britton, C. J. C. (1953). *Disorders of the Blood*, p. 103. Churchill, London.
Wintrobe, M. M. (1946). *Clinical Hematology*. Kimpton, London.

Use of "Anirrit" in the Treatment of Epilepsy

Vossen (1949) reported the first clinical trial of "anirrit" on 36 cases of epilepsy, treated over a period of two and a half years. The substance, which is a mixture of an organic bromine-hydantoin compound and phenylethyl barbituric acid, was found to have remarkably slight toxic side-effects in relation to its therapeutic effectiveness. The dose varied usually between 0.8 and 1.2 g. daily, but was increased in some cases to 1.6 g. a day. The absence of side-effects was stressed by Gordonoff and Kramer (1949), although they comment on the undesirable side-effects of the hydantoins as a group. Janz (1953) found anirrit most effective in the treatment of purely major epilepsy and inadequate in the control of minor and minimal epilepsy. No serious side-effects were observed in this group, although four cases showed a slight leucopenia. The most frequent side-effects were anorexia and loss of weight. Rett (1953) stated that where the hydantoins are used in large doses to control epilepsy depression of the bone marrow may be observed, but that this complication can be avoided by the use of anirrit, in which the hydantoin and bromide act synergistically and only small doses are needed. Almost complete freedom from major fits was obtained in 14 of the 21 cases, although the period of follow-up is not stated.

Recently a pilot observation was carried out for six months on the use of this drug in a small group of epileptic patients, and the results are given in the following report.

Sixteen patients suffering from epilepsy were studied. In 15 of them the attacks were idiopathic in nature, and in one they were due to brain damage from middle-ear and mastoid infection (brain abscess) in childhood. The patients had not responded to full doses of phenobarbitone, phenytoin sodium, primidone, or combinations of these drugs. They had been followed up long enough to judge that they were reliable witnesses, and it was reasonable to assume that a reduction in the frequency of the seizures would be observed over a short period if the drug was to be regarded as effective.

Nine of the patients were suffering from major epilepsy and four were having major and minor attacks; one had psychomotor attacks and one had minor akinetic seizures; only three of the patients averaged less than one attack a week. The ages of the patients varied from 3 to 39 years, three of them being under 10.

The dose of anirrit was slowly increased under observation in order to determine a suitable maintenance dose. Children between the ages of 3 and 9 received maximum daily doses of 0.4 to 0.8 g. The dose for adults varied between 0.6

and 2 g. daily (one case), although the majority of patients had 1.6 g. a day. The drug was used alone except in four cases; in one it was added to amylobarbitone and in the other three to primidone, these two drugs having been taken for some time before anirrit was given. Treatment has been continued for periods varying from one week to six months.

RESULTS

Of the 16 patients, seven (44%) were improved, and two of these were without attacks for three and four months respectively; in four others there was considerable reduction in the frequency, and in one the attacks were much less severe, though no less frequent. Nine patients (56%) were no better, and two of these were worse. In one the attacks continued to progress in frequency, and the patient went into status epilepticus; in the other case aplastic anaemia arose, but recovery occurred gradually under treatment, when the drug was withdrawn. The other seven patients had anirrit for from one week to five and a half months. In some there was a fluctuation in the frequency of the attacks, but over a period of some months no improvement could be observed beyond that seen with other treatment.

The results of treatment in this group are less satisfactory than in the group reported by Rett (1953), in which 66% became almost completely free from fits. In the present series less than half were improved, and only two patients were free from attacks over a period of three to four months. The numbers suffering from different patterns of epilepsy were too small to judge the response of various kinds of seizure to anirrit, but the patients suffering from major and minor attacks responded as well as those with grand mal alone.

Seven patients had an electroencephalogram taken before and during treatment. The records were changed in two—one improved and one worsened—both changing in the same way as the clinical condition of the patient. The record was not significantly changed in five patients, although clinically one was worse, one was unchanged, and three were better.

Toxic Manifestations.—No charted weight loss was seen in the patients who weighed themselves regularly, although one patient stated that there was an initial loss of weight, which did not continue in spite of treatment being carried on. Anorexia was noted in only one of the group. Gingivitis, present before treatment with anirrit, persisted, but it did not occur during treatment with anirrit alone. In only one patient, a child of 6 years, was the white blood count reduced. The severe fall in granulocytes necessitated repeated transfusions, but eventually the patient made an excellent recovery. In none of the other 11 patients whose blood counts were checked did the white-cell count fall below normal levels.

CONCLUSIONS

Patients were submitted for trial with anirrit when they had failed to respond to treatment with phenobarbitone, phenytoin sodium, and primidone; they were considered to belong to the worst group of epileptics. Seven of the 16 cases showed definite improvement and two were relieved from attacks for several months. This preliminary observation indicates that anirrit has a place in the treatment of epileptics, but toxic side-effects (leucopenia) indicate that its use must be accompanied by routine blood counts and careful control.

I thank the medical committee for permission to carry out this investigation and Dr. Stuart Phillips, Ph.D., for the supply of anirrit from Multipax Chemicals Ltd.

P. K. ROBINSON, M.B., M.R.C.P.,
Registrar, National Hospital, Queen Square, London.

REFERENCES

- Gordonoff, T., and Kramer, H. (1949). *Ther. Umsch.*, 10, 163.
Janz, D. (1953). *Nervenarzt*, 24, 208.
Rett, A. (1953). *Paracelsus (Wien)*, 6, Fasc. 11.
Vossen, R. (1949). *Ther. Umsch.*, 10, 165.