BRITISH MEDICAL JOURNAL

LONDON SATURDAY MARCH 31 1956

PRINCIPLES OF SURGICAL TECHNIQUE*

BY

Sir JAMES PATERSON ROSS, K.C.V.O., M.S., F.R.C.S., F.A.C.S.(Hon.)

Professor of Surgery, University of London; Director of Surgical Professorial Unit, St. Bartholomew's Hospital

I wish first of all to express to the Council of the University of Liverpool my deep gratitude for their invitation to deliver this lecture in memory of Sir William Mitchell Banks, who won for himself not only high honour for his great services to the University and to surgery but also warm affection for his sterling character and gentle kindness. The attempt to pay a worthy tribute to the memory of so great a figure in the history of the Liverpool School of Medicine is a formidable task, which is rendered still more arduous by the high standard set by my predecessors.

The first difficulty was to find a subject which could do justice to the occasion. The original intention was that the Memorial Lecture should deal with some subject connected with the study or practice of surgery in which Mitchell Banks had been interested. His teaching, his practice, his lectures, and his writings all indicate that he devoted particular attention to cancer of the breast, to the surgery of hernia, and to medical education. Since my own interests also lie in these directions the difficulty in finding the right subject for the lecture may not be apparent until one comes to realize how many others seem to share these interests, and how much has been written and said on these matters during the past few years.

I was somewhat relieved to find that Mitchell Banks himself had experienced similar difficulty, and I am glad to follow his advice as best I can. As the introduction to an address on medical education in 1891 he said:

"To one whose duty it becomes to give an address, the most difficult part of the undertaking is to choose a subject. Personally I have found under such circumstances that I have done best by speaking about something in which I was at the moment deeply interested, and with the details of which I was thoroughly conversant. This mode of procedure is apt to cripple all flights of genius, and is an effectual bar against attempts to soar into the realms of the oratorical or the ideal. On the other hand, it permits the speaker the use of plain language and the exercise of common sense, so that, whether his audience thinks much of his performance or not, at all events they feel that he knows what he is talking about."

Surgical Craftsmanship

The craft of surgery was a major interest to Mitchell Banks, as it must have been to many of his contemporaries who were seeing the more extensive surgical procedures that could be undertaken with the help of

*The William Mitchell Banks Memorial Lecture delivered in the University of Liverpool on January 19, 1956.

ether and chloroform. The modern age of anaesthesia had dawned while he was a child in Edinburgh. He was dresser and house-surgeon to Syme shortly after Lister had left Edinburgh for the Royal Infirmary at Glasgow, but there must have been many opportunities for Mitchell Banks to meet Lister either in Edinburgh or during the subsequent period when he was a demonstrator of anatomy in Glasgow. In his writings there are many references to Lister, of whose work he always speaks with deep admiration and respect, though he was frankly critical of some of the leading surgeons of his time, even of Syme himself.

A surgeon may on occasion have various qualities attributed to him. He may be regarded as somewhat of a scientist, something perhaps of an artist, but certainly, if he is any good, he is a craftsman. The craft of surgery is to the art much as tactics is to strategy; and, while the art may develop with the addition of increasing knowledge into an applied science, surgery without craftsmanship is as dead as faith without works.

To meet with some novel application of science to our practice is to any of us a stimulating and even an exciting event; but to the surgeon, however experienced and however gifted as an experimentalist, there can be nothing which gives such deep satisfaction as to watch a good brother craftsman at work. This idea was so excellently expressed by H.R.H. the Duke of Edinburgh on the occasion of his formal admission to the Honorary Fellowship of the Royal College of Surgeons of Edinburgh that I venture, with due respect, to quote his words:

"In these days when everything is either raised or reduced to a science, which really means that the human element is removed as much as possible, it is refreshing to find the word 'craft' applied to something as august as surgery. But it is certainly the right word, for the surgeon is the craftsman who draws together the laboratory work of the chemist and the physicist, the nutritional expert and the bacteriologist, the biochemist and the psychologist, and, through the skill of his hands, is the person ultimately responsible for the multiplying of human enjoyments and the mitigation of human suffering."

When thinking over the methods whereby this skill may be acquired—the training of a surgeon—there can be no better guide than Wilfred Trotter (1935), whose ideas about Medicine as a practical art are so illuminating: "An art is taught first in precepts and rules, secondly in experience of its material, and thirdly—and most important—by example. The method of

apprenticeship is thus the keynote of education in the practical arts, because it brings the pupil into familiar contact with his material and gives him the constant example of his teacher in the actual things he himself will ultimately have to do."

For my own part, I have been fortunate in having come under the influence of several master surgeons. To every one of them I owe something, and to some of them I owe almost everything. I would like to acknowledge, with affectionate regard, my deepest debt to Sir Thomas Dunhill for his example not only in technical skill but also in wisdom and sound judgment. I owe much to the privilege I enjoyed of assisting Lord Moynihan; and still more to Harvey Cushing, who was the Mitchell Banks Memorial Lecturer in 1910. I went to Boston to work for him partly in order to study neurological surgery, but chiefly because he was an outstanding master in the craft of surgery. It was he who taught me that the fundamental principles of surgical technique are asepsis, gentleness, and the control of haemorrhage.

Asepsis

Mitchell Banks came to Liverpool in 1868, the year after Lister's publication of the use of carbolic acid as an antiseptic, and when reading his papers one gains a vivid impression of the controversy which followed this innovation in surgical technique. Writing in 1882 on the results of operation for cancer of the breast, he gave an account of the principles upon which he based his treatment, the first being Listerian antiseptics. He refers, however, to his fear of the risk of increasing the shock of a long operation by exposing the patient to the carbolic spray, and for this reason he does not lay down a hard-and-fast rule. "As a rule I employ them [antiseptics]; in hospital always. . . . For this reason [risk of shock], in private, where there is but little fear of septic influences, if the patient be weakly I content myself with simply washing out the wound well with carbolic solution and maintaining a modified antiseptic dressing."

In spite of all his pains, however, he lost 5 of the recorded 46 cases of cancer of the breast from sepsis, and, deploring this tragic figure, he adds: ". . . but we know that had these same patients been all in perfect hygienic conditions—had they for instance been in the atmosphere of Zermatt—they would have recovered. Fortunately every year is improving our knowledge of how to ward off the poisonous conditions, and when we shall have banished them the range of operative surgery will hardly know any bounds."

Considering that this was written over 70 years ago it is sad to think that the goal he regarded as being almost in sight is still unreachd; and it is very disturbing to discover how little difference there may be between the conditions Mitchell Banks was describing and those which exist in many operation theatres to-day. The situation is rendered particularly dangerous by the natural but false assumption that a modern theatre which has been fitted with what appears to be adequate ventilation must be beyond suspicion, and that if wounds are going wrong the cause of the trouble must be sought elsewhere.

Our experience of cross-infection in hospitals during the second world war had opened our eyes to the importance and complexity of the problem (Miles *et al.*, 1940), and when, later on, we were confronted with repeated outbreaks of wound infection we proceeded to investigate them by the methods which had proved so valuable during the war. We noted at once that there was an important difference between the earlier infections and the more recent ones—the former were almost entirely streptococcal, whereas the infections now giving us trouble were with penicillin-resistant staphylococci. This suggested the possibility that the whole problem—channels of infection and the methods to be adopted for blocking them—might be quite different from that of streptococcal cross-infection. Realizing our diffi-

culties, we appealed to Professor L. P. Garrod and Dr. R. A. Shooter, whose interest, guidance, and advice in the bacteriological investigations have been invaluable, and are most gratefully acknowledged.

Although we realized that the problem might be in some respects a different one, we went first to the wards, where we had previously found the source of the trouble. On this occasion, however, we could not find fault with the technique of ward dressings. But it was soon discovered that samples of dust collected from various parts of our wards were laden with penicillin-resistant staphylococci, and since this seemed the most likely source of the infection there remained the question of how the organisms were reaching the wounds.

We had noticed that most of the patients who suffered these wound infections had pyrexia which appeared almost immediately after the operation; furthermore, most of the infections extended into the deep layers of the wounds, and for these two reasons we assumed that infection must be getting in at the time of the operation, and therefore we shifted the scene of our inquiry to the theatre. From the beginning of the investigation we had kept a "wound book" in which are recorded the surgeon's name, the nature of the operation, and its duration, the solution used to clean the skin, and any antiseptic introduced into the wound before closure. When the stitches were removed further notes were added to indicate the state of the wound, and, if infection had occurred, its severity and the infecting organisms were recorded. These records have been kept in full since July, 1954, and in the course of the following seven months the monthly percentage of "clean" operation wounds which suppurred varied between 5 and 11%, the average for that period being 9%, in spite of our initial efforts to discover the channels of infection.

It soon became evident that these wound infections were not confined to the practice of one surgeon but were equally distributed throughout the staff of the unit. It was therefore not surprising that the search for throat and nasal carriers or skin carriers of staphylococci among the surgical and nursing staff was unprofitable. Various solutions were employed to cleanse the skin before incision, but none of these well-approved agents seemed to have any influence upon the incidence of infection. The duration of the operation did not appear to be of prime importance.

Ventilation of Theatres

Past experience had taught us the danger of ventilating an operation theatre with an exhaust fan; but as our own theatre was supplied with filtered air from a supply fan we had assumed that

the fault could not be there. However, the results of Dr. Shooter's examination of the theatre air, using a slit sampler, forced us at last to direct our attention to the ventilation of the theatre. He found that samples of 5 cubic feet (0.14 cubic metre) of air grew varying numbers of colonies at intervals in the course of an operating session (Fig. 1). In order to find out whether this variation might be due to introducing organisms with the patient's clothes or by some other means whereby infection might be carried from ward to theatre in the course of an operation "list,"

FIG. 1.—Colony counts from samples of air in operation theatre during working hours before change in ventilation. Each upright represents colonies from 5 cubic feet (0.14 cubic metre) of air. Samples of 200=200+.

FIG. 2.—Colony counts from samples of air in empty operation theatre and before ventilation change.

samples were taken one day when the theatre was not in use, but the air was found to be just as heavily infected as ever (Fig. 2).

It had been assumed that the theatre was being supplied with air from outside, and, in order to determine the content of this outer air, samples were taken outside our theatre window and also outside a window on the second floor of the Pathological Laboratory Block, but all these samples showed a very low bacterial content (Fig. 3).

FIG. 3.—Colony counts from samples of air taken outside window of operation theatre and outside window on second floor of Pathology Block.

FIG. 4.—Colony counts from samples of air in operation theatre during working hours after change in ventilation (except February 22, when supply fan was stopped).

FIG. 5.—Graph to show incidence of infection of clean wounds before and after change in ventilation of theatre; date of change indicated by arrow.

removed the filters which had been impeding the action of the supply fan and blocked up all the exit vents in the walls of the theatre. To our great relief the smoke test then showed that the air current was passing from the theatre out into the corridor. Subsequent sampling of the theatre air showed that the bacterial colony counts were much lower so long as the supply fans were kept running. The air supplied by this fan is warmed by radiators, but on a particularly cold day, February 22, 1955, I became concerned about the cooling effect of the fan and had it stopped. The bacterial content of the theatre air rose immediately (Fig. 4), and unfortunately the wound of an amputation which was carried out with particular care to avoid any "neighbourhood" source of infection became heavily infected with penicillin-resistant staphylococci. Since this experience we have never operated without the supply fan in action, and there has been a striking decrease in the incidence of wound infection. Of the next 500 "clean" cases done after altering the ventilation only 5 became infected, including the one which occurred on the day the supply fan was stopped (Fig. 5).

By good fortune we had only one important source of infection to deal with, and by still better fortune our rather rough-and-ready method of dealing with it had been success-

ful. Whenever a forced draught is introduced into a room there is a tendency for turbulent eddy formation to occur. We traced the air currents in our theatre by means of smoke and found that the supply fan was providing a constant current of pure air at the level of the operation table, and, although there was some turbulence in the air above it, the stream was moving rapidly enough to prevent much admixture with the turbulent air above. The scheme of ventilation which is generally regarded as the best at present available is one in which the supply of filtered air enters near the roof of the theatre and descends to exit vents near the floor. Even with this arrangement turbulence may lead to contamination of the air at the level of the top of the operation table by air currents sweeping upwards from the floor unless the inlet and outlet systems are carefully regulated. It would appear, therefore, that the arrangements for ventilating a theatre, however good they may look on paper, should always be tested with smoke or otherwise to demonstrate what is actually happening.

It had not previously been realized that certain exit vents in the walls of the

theatre which were connected through ducts to a ventilating shaft which extended from top to bottom of our stack of theatres were acting as a powerful exhaust "fan," whether in fact the fan on the roof was working or not, merely because of the upward draught of warm air in the vertical shaft. This suction action of the exhaust shaft was removing air from the theatre much more rapidly than the input fan could supply it, and therefore air was being drawn in through the doorways.

In order to reverse this air current we

removed the filters which had been impeding the action of the supply fan and blocked up all the exit vents in the walls of the theatre. To our great relief the smoke test then showed that the air current was passing from the theatre out into the corridor. Subsequent sampling of the theatre air showed that the bacterial colony counts were much lower so long as the supply fans were kept running. The air supplied by this fan is warmed by radiators, but on a particularly cold day, February 22, 1955, I became concerned about the cooling effect of the fan and had it stopped. The bacterial content of the theatre air rose immediately (Fig. 4), and unfortunately the wound of an amputation which was carried out with particular care to avoid any "neighbourhood" source of infection became heavily infected with penicillin-resistant staphylococci. Since this experience we have never operated without the supply fan in action, and there has been a striking decrease in the incidence of wound infection. Of the next 500 "clean" cases done after altering the ventilation only 5 became infected, including the one which occurred on the day the supply fan was stopped (Fig. 5).

Asepsis or Antisepsis

The contemplation of the numerous channels whereby staphylococci can reach an operation wound may quite understandably lead to the conclusion that contamination is inevitable, and that the only hope must lie in measures to destroy the organisms in the wound rather than to prevent their access to it. Our experience recounted here indicates how dust will infect the air of the theatre unless care is taken to ensure that the rate at which clean air is supplied is greater than that at which air is extracted from the theatre. But airborne infection may also be conveyed by the patient's clothes or blankets unless special precautions are taken to prevent it. Furthermore, it must be understood that the infected dust need not settle directly on the wound in order to infect it; particles which settle on instruments, towels, or swabs, or in bowls containing water or saline, may ultimately find their way into the wound.

Those who regard this kind of air contamination as unavoidable rely upon ultra-violet light to destroy the organisms, and there is no doubt that the numbers of living bacteria in air can be very greatly reduced by this method. In my opinion it is not correct to introduce such an antiseptic procedure unless every possible means of preventing the access of organisms to the air of the theatre has been tried and has failed.

Furthermore, it is important to remember the work of Devenish and Miles (1939), which showed how staphylococci from skin carriers could get into wounds through punctures in gloves and through the sleeves of gowns, and it may be that nasal carriers can infect wounds if efficient masks are not worn. For all these reasons surgeons have sought refuge in antiseptics—either introducing an antiseptic substance into the wound before it is closed, or else operating under an umbrella of penicillin or some other antibiotic. We have tried both these methods, and agree with Blowers and his colleagues (1955), who found that they did not work. The indiscriminate "prophylactic" use of penicillin is probably dangerous, as well as useless, for there is good evidence that the prevalence of resistant organisms depends upon the amount of antibiotic used; this may account for the widespread increase in the incidence of wound infection by penicillin-resistant staphylococci which has been reported recently from the United States (Howe, 1954) as well as from surgical centres in Britain.

If I may quote Mitchell Banks again, there are no "flights of genius" in what I have had to say about the importance of asepsis in surgical technique. The matters I have brought to your notice are common knowledge; yet in operative technique, as in diagnosis, it is the neglect of something which is well known and even obvious, rather than ignorance of something which is unusual or obscure, that lands us in difficulty.

Gentleness

An essential element in Cushing's principle of gentleness has been described by Moynihan in these words: "Infinite

gentleness, scrupulous care, light handling and purposeful, effective, quiet movements which are no more than a caress, are all necessary if an operation is to be the work of an artist, and not merely of a hewer of flesh." Since the healing of a wound is a vital process it can be carried through with perfection only in tissues which were initially healthy and which have suffered only the minimal amount of injury during the operation. The term "gentleness" can therefore be amplified, as Moynihan taught that it should be, to include everything conducive to the proper care of the tissues. This involves the whole process of pre-operative preparation, the correction of fluid and electrolyte imbalance, the adequate treatment of anaemia, and the supply of proper nourishment to restore depleted stores of protein, sugar, and vitamins.

In the course of the operation itself consideration must be given to the protection of the patient from any harmful effects arising directly or indirectly from the anaesthetic agent employed; and the tissues exposed in the wound must be shielded so far as possible from all kinds of injury, whether mechanical, physical, or chemical. Clumsy manipulation or powerful retraction are obvious faults in technique; but the damage done by drying or cooling of tissues or excessive coagulation by diathermy, or by the introduction of chemical substances or solutions which are excessive either in amount or in concentration, must never be forgotten.

Control of Haemorrhage

One can judge the surgical craftsman by the way he deals with bleeding. The master knows the appropriate method to employ under differing circumstances, and saves time by recognizing the vessels which will seal themselves off spontaneously; he knows how to keep the wound dry while he is working, and how to avoid a haematoma after he has finished. Even in the absence of organisms a mass of blood clot seriously delays the healing of a wound; but the few bacteria which could be destroyed in a dry wound will inevitably cause suppuration in a haematoma.

I remember hearing Sir D'Arcy Power say that a surgeon's attitude to haemorrhage changes in the course of his career. As a young man he tends to fear it too much; he then becomes overbold; and when he is mature his experience tells him when he may be bold but also when he must be particularly cautious. Since blood has become readily available for transfusion one may perhaps add the observation that the inexperienced surgeon tends to give blood when it is not really necessary. Provided the patient's blood condition is satisfactory before operation it should not be necessary to give a transfusion for a radical mastectomy, a gastrectomy, or a block dissection of the neck unless some unusual complication is encountered. Since blood transfusion is not free from risk, a "small" transfusion should never be given as a precaution against post-operative collapse—a patient needs either several pints of blood or none.

The replacement of blood lost in injury or in disease processes such as peptic ulceration permits the surgeon to operate with deliberate precision rather than in the hurry and excitement of a hazardous enterprise which requires supreme skill and good fortune if it is to be successful. Without blood transfusion many of the prodigious extensions in the scope of surgery which have taken place in the last 30 years would not have been possible; yet, oddly enough, in the same period transfusion has tended to eliminate the dramatic from the operation theatre.

The more recently introduced methods of controlling the circulation during operation by hypotension and hypothermia have already opened up fresh fields for surgical enterprise which would have been unattainable without their aid. It may sometimes be tempting to employ these methods for big operations which nevertheless can be performed without them, but it is probably wrong and may even be adding to the surgical risk to do so. Every unnecessary stitch is an error of judgment; so also is the use of an elaborate method when a simpler one will give the same result, for simplicity is still the hall-mark of good surgery.

The Gentle Doctor

The worker in wood or metal or stone gets his reward from the good craftsman's joy in work well done, and he may even endow his craftsmanship with something of his own personality so that his work survives as a memorial to his memory. The surgeon who pursues his craft in a medium which is already living may also derive somewhat the same satisfaction from his handiwork, though he will usually be prepared to acknowledge his debt to Nature for rounding off the rough corners he may have left her to finish for him. A few may have the satisfaction of adding something substantial to the science or to the practice of surgery; but every surgeon may reap his reward in the confidence and the gratitude and even the affection of his patients, which he gains not merely by his skill but also by the sympathy, the friendly help, and the encouragement he has given when these were sorely needed. Mitchell Banks gave expression to this idea in "The Gentle Doctor," which was an address to students of the Yorkshire College at Leeds, when he described his ideal as "an honest man and a gentle man doing his best to relieve suffering."

That he had his reward in his lifetime is shown by references made in the many tributes which were paid to his memory by professional colleagues and by laymen alike at the time of his death, and I will conclude this memorial lecture with a few sentences from the eulogy delivered by the Rev. Dr. Watson, better known perhaps as "Ian Maclaren," the Minister of the Rodney Street Scottish Presbyterian Church where Mitchell Banks had worshipped. "He was a safe and sound man, who created and merited confidence. . . . Above all there dwelt in him that soul of kindness without which no man can reach the height of his calling, in medicine or any other profession. . . . True science honours life in every form, but chiefly in its highest, in man; it exists to study, to deliver, to reinforce, to glorify it; it is the eager, patient, tender, pitiful servant of every human being. Unto this divine service Mitchell Banks gave himself, and we may say now he is gone that he fulfilled his high commission. . . . Because he was so brotherly to every fellow creature and so true to his friends we loved him living, and now when he is gone we shall keep his memory green."

I wish to express my gratitude to Dr. R. A. Shooter for Figs. 1-4.

REFERENCES

- Blowers, R., Mason, G. A., Wallace, K. R., and Walton, M. (1955). *Lancet*, 2, 786.
Devenish, E. A., and Miles, A. A. (1939). *Ibid.*, 1, 1088.
Howe, C. W. (1954). *New Engl. J. Med.*, 251, 411.
Miles, A. A., Schwabacher, H., Cunliffe, A. C., Ross, J. P., Spooner, E. T. C., Pilcher, R. S., and Wright, J. (1940). *British Medical Journal*, 2, 855 and 895.
Trotter, W. (1935). *Ibid.*, 2, 609.

One of the services which the Patent Office renders to scientists and industry is the publication of abridgments of accepted patent specifications. These give essential particulars of such specifications in a very concise form with line drawings where necessary, and show the trend of invention in the United Kingdom. The abridgments are published in series of 20,000 specifications, divided into 44 groups, each group covering a field of invention; medicine and surgery fall in group VI, together with beverages, food production, and tobacco. Initially they are sold in sets of sheets, each part consisting of 16 pages and containing about 50 abridgments. Individual volumes are then assembled and indexed by subject and name of applicant. During the last war the publication of abridgments had to be suspended, but the Board of Trade announces that group volumes of specifications accepted during the war period are now on sale. A complete range of volumes, which includes specifications published before November, 1951, is now available. Abridgments in sheet form include specifications published before August, 1955. The volumes in bound and sheet form can be obtained from the Sale Branch, Patent Office, 25, Southampton Buildings, Chancery Lane, London, W.C.2.