Our incidence of toxæmia was only 6.1%, compared with their 14.8%. We are unable to give a satisfactory explanation of this difference. Our stricter criterion with regard to weight gain has not altered the incidence to any appreciable extent, but our wider selection of women booked for delivery may have contributed significantly to the difference.

With regard to the close association between toxæmia of pregnancy and essential hypertension, it is interesting to note that Maxwell and Maxwell (1955) found no significant alteration in the distribution of the blood groups in the latter condition.

Summary

The distribution of blood groups has been examined in a series of 11,086 pregnant women, 675 of whom developed toxæmia of pregnancy.

There was no significant difference in the distribution of ABO blood groups in the toxæmic as compared with the non-toxæmic group of women.

We are indebted to Dr. Fraser Roberts for help in the statistical analysis, and grateful to Dr. J. V. Dacie for access to the records of the Blood Transfusion Laboratory at Hammersmith Hospital; we also thank Professor J. C. McClure Browne for his advice and encouragement.

REFERENCES

- Discombe, G., and Meyer, H. (1952). *Amer. J. clin. Path.*, 22, 543.
Hooper, I. (1947). *Irish J. med. Sci.*, p. 471.
Maxwell, R. D. H., and Maxwell, K. N. (1955). *British Medical Journal*, 2, 179.
Pike, L. A., and Dickinson, A. M. (1954). *Ibid.*, 2, 321.
Richardson, J. R. E. (1954). Quoted by Pike and Dickinson (1954).
Sachs, H. (1940). *Irish J. med. Sci.*, p. 145.

CORTISONE TREATMENT OF THE LOW-SALT SYNDROME

BY

I. G. GRABER, M.D., F.R.C.S.Ed.

PETER BEACONSFIELD, M.D., Ph.D.

AND

OWEN DANIEL, M.B., F.R.C.S.

From the Department of Surgery, Postgraduate Medical
School of London, W.12

After any major surgical procedure there is usually a decrease in the concentration of plasma sodium (Wilkinson *et al.*, 1951). This hyponatraemia can be aggravated by vomiting, diarrhoea, paralytic ileus, or indiscriminate infusion, and can generally be corrected by suitable electrolyte replacement. We have encountered recently five patients who developed a low-sodium syndrome following major surgical procedures and who failed to improve despite what was regarded as adequate replacement therapy. Acting on the supposition that this intractable hyponatraemia was due to transient exhaustion of adrenocortical function, such as is known to occur sometimes in the presence of severe stress (Bland, 1952), we administered cortisone. The results suggest that cortisone therapy is of considerable value in these forms of hyponatraemia, and we report in this paper details of three typical patients. These patients were not the subjects of a planned investigation, and the biochemical data are those obtained during the course of routine management.

Case 1

A man aged 48 was transferred to Hammersmith Hospital because of severe and persistent electrolyte imbalance which had developed following right hemicolectomy, performed six days previously for ileo-caecal intussusception. On each post-operative day he had been given approximately 270 mEq of

sodium and chloride in the form of 3,000 ml. of intravenous fluid consisting of 0.9% (normal) saline alternating with 0.18% saline with dextrose. On admission the patient complained of severe diarrhoea, which had started 24 hours previously.

On examination he was cold, clammy, greatly dehydrated, vasoconstricted, and very restless. To exclude the possibility of a jejuno-colic anastomosis as the cause of the diarrhoea, he was given a capsule containing 200 mg. of indigo carmine, and the dye appeared in his stool six hours later. Blood analysis showed haemoglobin 14.5 g., protein 6.3 mg., and urea 45 mg. per 100 ml. Plasma electrolytes were sodium 114, chloride 83, potassium 3.5, and bicarbonate 37 mEq/l.

He was immediately treated by oral administration of 2 g. of sodium chloride and 1 g. of potassium chloride every four hours, together with half-normal saline by mouth to the amount of 2,500 ml. a day. This regime provided a minimum daily intake of 382.5 mEq of sodium, 463.5 mEq of chloride, and 81 mEq of potassium. His clinical appearance and diarrhoea improved, but there was little change in his plasma chemistry. After four days' treatment plasma sodium was 114, chloride 88, potassium 4.2, and bicarbonate 22 mEq/l.; blood urea was 23 mg. per 100 ml. The 24-hour urine output was 720 ml., containing sodium 14.4, chloride 71, and potassium 24 mEq/l. The volume of the stools at this time was not excessive, and it was believed that the salt balance was positive.

In view of the persistent hyponatraemia and hypochloræmia a course of cortisone acetate, 25 mg. every six hours, was started on the fourth day after admission and the administration of salt by mouth stopped. Twelve hours after starting the cortisone the plasma sodium had increased to 145, chloride to 100, and potassium to 4.5 mEq/l. Urine output during this 12-hour period amounted to 840 ml., containing sodium 50, chloride 71, and potassium 24 mEq/l. Cortisone administration was continued for four days, during which time clinical improvement and normal plasma chemistry were maintained. When the cortisone was stopped the plasma sodium decreased slightly for two days, then returned again to normal levels and the patient made an uneventful recovery. It should be noted that, before cortisone was given, low-plasma sodium and chloride persisted despite a positive salt-and-water balance. Within 12 hours of starting cortisone treatment plasma sodium and chloride returned to normal, although the administration of extra salt had been stopped and the urinary excretion of sodium had nearly quadrupled.

Case 2

A married woman aged 54 was admitted to hospital suffering from jaundice and intermittent hepatic fever. A year previously she had undergone a successful mitral valvulotomy. A week after admission an impacted stone was removed from the ampulla of Vater, cholecystectomy was performed, and a T-tube was inserted in the common duct. The patient received 500 ml. of blood during the operation. During the first week after operation bile drainage varied from 500 to 1,500 ml. a day and contained sodium 140, chloride 118, and potassium 4.4 mEq/l. On the sixth post-operative day plasma sodium was 116, chloride 88, potassium 4, and bicarbonate 24 mEq/l. The 24-hour urine output was 1,200 ml., containing sodium 38, chloride 40, and potassium 115 mEq/l. During this period the patient took about 2,000 ml. of water by mouth every day and a slowly increasing mixed diet, but additional sodium chloride replacement was withheld because of feared cardiac decompensation, except during the first two days, when 1,000 ml. of 0.45% saline with dextrose was given intravenously each 24 hours.

On the seventh, eighth, and ninth days after operation she was given 12 g. of sodium chloride and 6 g. of potassium chloride by mouth in addition to her normal diet. This supplement provided 200 mEq of sodium, 281 mEq of chloride, and 81 mEq of potassium. During the same period 800 ml. of the 1,200 ml. of bile drainage was returned by rectal drip. Clinical improvement was made during these

three days and the bile discharge decreased, but the hyponatraemia and hypochloraemia remained virtually unchanged. Cortisone was then given orally in 25-mg. doses every six hours and the administration of extra salt stopped. Twelve hours after starting cortisone therapy plasma sodium increased to 128, chloride to 94, and potassium 5.2 mEq/l. Cortisone administration was continued for six days, by which time plasma sodium was 130, chloride 99, and potassium 4.5 mEq/l. During this period bile drainage averaged 500 ml. a day; thereafter it slowly decreased and the patient made a satisfactory recovery (see Chart).

Case 3

A married woman aged 38 was admitted to hospital for refashioning of a terminal ileostomy. A year previously colectomy had been performed for ulcerative colitis, after which she had made a good recovery until the ileostomy became partially stenosed. After refashioning the ileostomy 2,000 to 3,000 ml. of ileal content was discharged every day. The total electrolyte loss was measured and an adequate replacement of sodium and potassium was given intravenously, in addition to a normal diet and an oral intake of fluid of approximately 2,000 ml. a day. The excessive ileal discharge continued, and, although a positive sodium chloride, potassium, and fluid balance was maintained, it was obvious that she was going slowly downhill. Fourteen days after operation the plasma sodium was 93, chloride 61, and potassium 3.2 mEq/l. She was then given 25 mg. of corticotrophin every six hours for four days. Twenty-four hours after starting corticotrophin the plasma sodium was 133, chloride 90, and potassium 4.0 mEq/l. After a further 24 hours, plasma sodium was 134, chloride 102, and potassium 3.8 mEq/l. After the administration of corticotrophin the excessive ileostomy excretion ceased, intravenous therapy was discontinued, and the patient made an uneventful recovery. The dosage of corticotrophin was decreased gradually.

Comment

Post-operative patients with a low plasma sodium and chloride usually respond well to the administration of salt. Occasionally, abnormally low plasma sodium and chloride levels persist despite the establishment of positive salt-and-water balance. We have found that cortisone can quickly remedy such a condition, and this observation throws some light on the effect of adrenocortical hormones on cell and capillary permeability. It has sometimes been assumed that the mineral activity of the posterior pituitary and adrenocortical hormones affects solely renal tubular function and that changes produced by these hormones on intracellular and extracellular fluid are a consequence of renal activity rather than an expression of hormone action on cell permeability (Gaunt *et al.*, 1949). Investigations by Eliel and Pearson (1951) on patients with hypopituitarism and by Graber and Beaconsfield (1955) on adrenalectomized patients indicate that large shifts of sodium and potassium occur across the cell membrane under hormonal control.

Thorn *et al.* (1955) have postulated that 11- and 17-oxygenated steroids have an important effect on the intracellular and extracellular concentration of sodium and potassium in many chronic disease states, and that following the

administration of large doses of cortisone or hydrocortisone a shift of water and electrolytes occurs, with the overall result that the sodium content of the extracellular fluid compartment is increased, thus increasing the load of sodium presented to the kidneys. It has also been shown in adrenalectomized dogs (Grollman, 1954) and rats (Ledingham, 1954) that there is a significant increase in the sodium and decrease in the potassium content of the cells, especially in skeletal muscle and liver.

The rapid changes in plasma electrolytes following the administration of cortisone in the cases reported in this paper can best be explained on the basis of cortisone altering cell permeability, allowing sodium and chloride to pass quickly from the cell into the plasma, thus restoring a normal plasma electrolyte picture and relieving intracellular

Case 2. Post-operative plasma sodium and chloride concentration expressed as milliequivalents per litre.

oedema. This explanation is supported by the fact that cortisone administration was followed almost immediately by a greatly increased urinary output of sodium and a relatively diminished urinary output of potassium.

Our experience leads us to believe that cortisone has a place in the treatment of otherwise intractable post-operative low-salt syndrome. But the value of this treatment must be weighed carefully against the known dangers of cortisone administration.

Summary

The administration of cortisone can quickly correct severe post-operative hyponatraemia and hypochloraemia persisting despite adequate intake of sodium, potassium, and chloride. Three illustrative case histories are given, and their bearing on current theories on the mode of action of adrenocortical hormones is briefly discussed.

REFERENCES

- Bland, J. H. (1952). *The Clinical Use of Fluid and Electrolyte*. Saunders, Philadelphia and London.
- Eliel, L. F., and Pearson, O. H. (1951). *J. clin. Endocr.*, **11**, 913.
- Gaunt, R., Birnie, J. H., and Eversole, W. J. (1949). *Physiol. Rev.*, **29**, 281.
- Graber, I. G., and Beaconsfield, P. (1955). *British Medical Journal*, **2**, 704.
- Grollman, A. (1954). *Amer. J. Physiol.*, **179**, 36.
- Ledingham, J. M. (1954). *Clin. Sci.*, **13**, 535.
- Thorn, G. W., Laidlaw, J. C., and Goldfien, A. (1955). *Ciba Foundation Colloquia on Endocrinology*, **8**, 352.
- Wilkinson, A. W., Billings, B. H., Nagy, G., and Stewart, C. P. (1951). *Lancet*, **1**, 315.