Northern Hospital, under the direction of Dr. C. A. Clarke, and with the help of Miss S. M. Manning, working under a grant from the Research Committee of the United Liverpool Hospitals. We are also grateful to Dr. A. M. Cooke, physician-in-charge, Diabetic Clinic, Radcliffe Infirmary, Dr. M. M. Pickles, Director of the Serology Laboratory, Radcliffe Infirmary, and Mr. D. Hewitt, of the Department of Social Medicine, Oxford, for their help.

REFERENCES

- Aird, I., Bentall, H. H., Mehigan, J. A., and Roberts, J. A. F. (1954). *British Medical Journal*, 2, 315.
 ——— and Roberts, J. A. F. (1953). *Ibid.*, 1, 799.
 ——— (1955). *Ibid.*, 2, 321.
 Craig, J., and Wang, I. (1955). *Glasg. med. J.*, 36, 261.
 Dobson, A. M., and Ikin, E. W. (1946). *J. Path. Bact.*, 58, 221.
 Harris, H. (1949). *Ann. Eugen.*, 14, 293.
 ——— (1950). *Ibid.*, 15, 95.
 Holt, S. B. (1948). *Ibid.*, 14, 144.
 Joslin, E. P. (1947). *The Treatment of Diabetes Mellitus*, 8th ed. Kimpton, London.
 McConnell, R. B. (1955). *Proc. roy. Soc. Med.*, 48, 291.
 Maxwell, R. D. H., and Maxwell, K. N. (1955). *British Medical Journal*, 2, 179.
 Snedecor, G. W. (1946). *Statistical Methods*. Iowa State College Press, Ames, Iowa.

FURTHER OBSERVATIONS ON ABO BLOOD-GROUP FREQUENCIES AND TOXAEMIA OF PREGNANCY

BY

A. M. DICKINS, M.D., M.R.C.O.G.

J. R. E. RICHARDSON, M.R.C.S., L.R.C.P.

L. A. PIKE, M.B., B.S.

AND

J. A. FRASER ROBERTS, M.D., D.Sc., F.R.C.P.

From Perivale Maternity Hospital; the Department of Pathology, King Edward Memorial Hospital, Ealing; and the London School of Hygiene and Tropical Medicine

In a previous paper (Pike and Dickins, 1954) a comparison was made of the frequencies of the ABO blood groups of toxæmic and non-toxæmic women in 3,651 consecutive deliveries. The toxæmic mothers showed a higher frequency of group O than the remainder, the difference being fairly highly significant. It was realized, however, that larger numbers were needed, and the hope was expressed that other series would be studied at other hospitals.

As a first step the original material was re-scrutinized, with stricter criteria for the diagnosis of toxæmia. Ninety-three women were transferred from the toxæmic series to the non-toxæmic on the following grounds: (1) that the arterial pressure had reached 140/90 in labour only; (2) that the prescribed level of arterial pressure was attained by the systolic or diastolic reading only and not by both; and (3) that the history of toxæmia in a previous pregnancy, though probable, was not beyond all doubt. The modified figures, with the addition of cases over a short further period, were described briefly by one of us (Dickins, 1955). The significance of the difference was reduced, but the evidence remained fairly strong.

The original series comprised deliveries from March, 1951, to April, 1954. A second series, comprising deliveries at the same hospital from May, 1954, to December, 1955, has now been analysed. The second series does not confirm the original result, and it must now be concluded that we have no acceptable evidence that toxæmic and non-toxæmic mothers differ in their ABO frequencies. Some mystery remains, however—a mystery we cannot explain—as to why the original series gave the result it did. The Table shows the results for the original period, with the amended figures on re-scrutiny, and the later series.

ABO Frequencies in Toxæmic and Non-Toxæmic Women: Numbers and Percentages

	Toxæmic				Non-Toxæmic			
	O	A	B	AB	O	A	B	AB
<i>Basic Data</i>								
1. March, 1951–April, 1954	290	203	34	14	1,416	1,342	263	89
2. No. 1 on re-scrutiny	237	170	30	11	1,469	1,375	267	92
3. May, 1954–Dec., 1955	106	116	29	11	695	670	115	43
Total 2+3	343	286	59	22	2,164	2,045	382	135
<i>Corresponding Percentage Frequencies</i>								
1.	53.6	37.5	6.3	2.6	45.5	43.2	8.5	2.9
2.	52.9	37.9	6.7	2.5	45.9	42.9	8.3	2.9
3.	40.5	44.3	11.1	4.2	45.6	44.0	7.6	2.8
2+3 ..	48.3	40.3	8.3	3.1	45.8	43.3	8.1	2.9

If groups O and A only are compared the series as originally published gave a χ^2 for the difference between toxæmic and non-toxæmic mothers of 9.09 for one degree of freedom. If group O is compared with the total of the other three the χ^2 was 11.75 with one degree of freedom. The effect of the transfer of the 93 mothers on re-scrutiny is to lower these figures to 6.18 and 7.82 respectively, the second figure corresponding to a probability of less than 1 in 100. The new series (3 in the Table) shows an excess of A in the toxæmic mothers, but the difference from the non-toxæmic mothers is far below the level of significance ($\chi^2=0.77$ for the OA comparison).

When the whole material is combined the excess of group O in the toxæmic mothers is not significant. χ^2 is 2.17 for the OA comparison, and 1.72 when group O is compared with the total of the other groups. The two series are, however, significantly heterogeneous. With due regard to correct weighting, χ^2 for heterogeneity, with one degree of freedom, is 4.75 for the OA comparison and 8.48 for the comparison of group O against the other groups. Thus the material taken as a whole shows no significant excess of O in toxæmic women. The results for the two periods, however, do not agree, the discrepancy being fairly highly significant.

We have tried to find reasons for the discrepancy, but without success. Exactly the same criteria were applied in the diagnosis of toxæmia of pregnancy. The frequencies amongst all women grouped in the area and those delivered in hospital do not differ and there has been no secular change in either. For example, dividing women grouped from mid-1948 to the end of 1955 into 15 half-yearly periods, the total in groups O and A is 10,844. The relative proportions of O and A are perfectly homogeneous, χ^2 being only 8.26 for 14 degrees of freedom ($P=0.87$). The two series of toxæmic women are homogeneous within themselves, the first for high O and the second for low O. If each is divided into four periods the χ^2 's for heterogeneity within the two series, with 3 degrees of freedom, are as follows: for the OA comparison 6.73 for the first series, 3.68 for the second; for the O versus (A+B+AB) comparison, 5.44 and 1.73. All these values are non-significant at the 5% level. We have also subdivided the toxæmic patients into those with pre-eclamptic toxæmia of pregnancy and those with essential hypertension in pregnancy according to criteria described previously (Dickins, 1955). Both types show high O during the first period and low O during the second.

When it was originally thought that there was an excess of group O in toxæmia it was decided to reinvestigate the question of ABO heterospecific pregnancy as a possible explanation (Richardson, 1955). During both the original period of high O and the later period the proportion of heterospecific pregnancies corresponded closely to what would be expected on the basis of the gene frequencies. The total number of pairs of mothers and babies tested was 236. In all pregnancies which were ABO compatible a

Coombs test was carried out on infants' cells sensitized in mothers' serum. No other incompatibility was detected.

So far as we are able to analyse the figures, the first period, as shown above, consistently yields a fairly highly significant excess of O in toxæmia, the second period equally consistently yielded no such excess. It may be that there is a phenomenon to explain, but on the present evidence it seems likely that the original finding was due to chance. This experience serves to emphasize the need for large numbers in work in this field, the desirability of demanding a high level of significance before results are taken as more than an indication for further inquiry, and also the desirability of comparing results from different hospitals and different areas.

Summary

A further series does not confirm the finding (Pike and Dickins, 1954) that there is an excess of women of group O among those suffering from toxæmia of pregnancy.

REFERENCES

- Dickins, A. M. (1955). *Proc. roy. Soc. Med.*, **48**, 140.
Pike, L. A., and Dickins, A. M. (1954). *British Medical Journal*, **2**, 321.
Richardson, J. R. E. (1955). *Proc. roy. Soc. Med.*, **48**, 142.

ABO BLOOD GROUPS AND TOXAEMIA OF PREGNANCY

BY

M. G. PEARSON, M.B., F.R.C.S.Ed., M.R.C.O.G.
Senior Registrar

AND

G. D. PINKER, M.B., M.R.C.O.G.
Late Registrar

Institute of Obstetrics and Gynaecology, University of London

In a study of 541 patients with toxæmia of pregnancy, Pike and Dickins (1954) found a significant excess of women of blood group O as compared with non-toxaemic pregnant women. It was suggested that some of the features of toxæmia might be the result of an antigen-antibody reaction. We thought it would be of interest to repeat their investigation on a similar series of patients admitted to another hospital.

Material

The investigation was carried out on 11,086 consecutive admissions to the maternity department of Hammersmith Hospital between 1949 and 1954.

The majority of patients, both primigravidae and multiparae, are booked for delivery at this hospital in order of their application for admission at the antenatal clinic. Patients attending for the first time in late pregnancy are admitted on obstetrical, medical, or social grounds only.

For the purposes of this investigation and in order to conform with the criteria selected by Pike and Dickins, patients were regarded as having toxæmia of pregnancy if they showed two of the following signs: (1) a blood pressure of 140/90 or over after the 28th week of pregnancy; (2) clinical oedema or an average weight gain of more than 2 lb. (0.9 kg.) a week from the 28th week; (3) albuminuria in a catheter specimen, not due to infection. The level of weight gain which we regarded as significant in the absence of clinical oedema was greater than that selected by Pike and Dickins, but otherwise our criteria were identical with theirs.

Results

The distribution of blood groups in our series of 11,086 cases differed significantly from those of Pike and Dickins (1954), Richardson (1954), and Discombe and Meyer (1952), although the latter three series were compiled in an area

within a few miles of this hospital (Table I). If our series is compared with that of Discombe and Meyer the χ^2 for three degrees of freedom is 18.68.

It is suggested that this difference may be due to the fact that a high proportion of our maternity cases are drawn from the Hammersmith and Shepherd's Bush areas, with the result that approximately 25% of our bookings are Irish women. In Ireland the distribution of the blood groups is such that the ratio of group O to group A and that of group B to group A are higher than in England. Two series of figures illustrating the distribution of blood groups in Ireland—those of Hooper (1947) and Sachs (1940)—are shown in Table II. The expected distribution in a population containing 25% Irish has been calculated, using the figures of Sachs for Ireland and those of Discombe and Meyer for England.

The distribution of groups in toxæmic patients has been compared with that in non-toxaemic patients and the results have been contrasted with those of Pike and Dickins (Table III).

A comparison of our toxæmic and non-toxaemic patients shows there is no significant difference in the distribution of the blood groups (χ^2 for three degrees of freedom is 5.67); comparing the relative proportions of group O and the other groups taken together, χ^2 is 1.30; comparing the proportions of group O and group A, χ^2 is 3.16. A comparison of group B and group AB taken together against group O and group A taken together gives a χ^2 of 1.86. None of these differences is significant.

TABLE I.—Group Frequencies in Totals of Pregnant Women

Group	Present Series		Pike and Dickins		Richardson		Discombe and Meyer	
	No.	%	No.	%	No.	%	No.	%
O	5,105	46.04	1,706	46.73	4,579	45.79	4,578	45.78
A	4,448	40.12	1,545	42.32	4,238	42.38	4,219	42.19
B	1,138	10.26	297	8.13	866	8.66	890	8.90
AB	395	3.56	103	2.82	317	3.17	313	3.13
Total	11,086		3,651		10,000		10,000	

TABLE II.—Percentage Group Frequencies by Nationality in Pregnant Women

Group	Hooper (Ireland)	Sachs (Ireland)	Discombe and Meyer (England)	Distribution Expected with 25% of Irish Women	Present Series
O	54	53.6	45.8	47.7	46.0
A	32	32.4	42.2	39.8	40.0
B	11	11.5	8.9	9.6	10.2
AB	3	2.5	3.1	3.0	3.6

TABLE III.—Group Frequencies in Toxaemic and Non-toxaemic Patients

Group	Present Series				Pike and Dickins			
	Non-toxaemic Patients		Toxaemic Patients		Non-toxaemic Patients		Toxaemic Patients	
	No.	%	No.	%	No.	%	No.	%
O	4,809	46.2	296	43.9	1,416	45.5	290	53.6
A	4,150	39.8	298	44.1	1,342	43.2	203	37.5
B	1,080	10.4	58	8.6	263	8.5	34	6.3
AB	372	3.6	23	3.4	89	2.9	14	2.6
Total	10,411		675		3,110		541	

Discussion

Pike and Dickins, in their series of 541 toxæmic patients, found a significant increase in the proportion of women of blood group O as compared with the non-toxaemic group. They suggested that in some cases certain of the features of toxæmia might be due to an antigen-antibody reaction.

In a similar series of 675 toxæmic patients we have been unable to confirm their findings. We have not found any significant difference in the distribution of blood groups in these cases as compared with non-toxaemic women.