Medical Memoranda

Phimosis

The usual operation of circumcision is generally admitted to be crude and a mutilation. Many hold that its effects are disfiguring; it certainly deprives a tender part of its natural protective covering.

The following simple plastic procedure has, I believe, many advantages as an alternative for it, and I commend it with confidence to all interested in minor surgery. The figures are partly diagrammatic.

A director and then a narrow knife are inserted at the orifice in such a direction radially as to avoid blood vessels (Figs. 1 and 2). The incision is better done by transfixion,

and should be of such a length that, when opened out into a circle, it should make a comfortable fit round the organ (Figs. 3 and 4). When sure that the glans will just go comfortably through, a single absorbable suture is placed at the angle of the V, to hold the two layers together (Fig. 4). Both layers of the prepuce are then drawn right down towards the base, the inner one coming into a position of eversion (Fig. 5).

The edges will now encircle the corpus penis completely except for a tiny bridge of skin on the ventral aspect. This is what constituted the original orifice. The edges will

probably remain in contact all the way round—not as indicated in Figs. 5 and 6—and without any puckering, so that no further stitching may be needed.

However, one more suture is advisable or adhesive plaster on each side at the base to keep the whole skin drawn well down to prevent the cut edges from turning in on themselves (Figs. 5 and 6). These should remain until healing is complete, when the whole foreskin can be drawn again into its natural position. Little, if any, dressing is required, the cut being so far removed from the meatus urinarius. The frenum is not cut and there is no crushing.

W. REGINALD WILSON, M.B., B.Ch.

Tardive Cutaneous Leishmaniasis

The incubation period of cutaneous leishmaniasis is variable and sometimes unexpectedly prolonged. Manson-Bahr (1954) states that it may be as long as 15 months or even more, and Smith (1955) has drawn attention to the possibility of an incubation period of several years by recording a case in which a typical Oriental sore first appeared three years after the patient's return to England from an endemic area (North Africa).

As long ago as 1936 Napier and Halder reported a similarly prolonged incubation period. Their patient had moved from an endemic area in North-West India (Rajputana) to Calcutta, and had lived there continuously for over three years before developing typical facial cutaneous leishmaniasis. Two comparable cases occurring in Canada have been described by Forsey (1950) and Fidler (1952). In each of these the lesion first appeared between three and four years after the patient's final departure from endemic areas in Europe and the Middle East. The Russian workers Latyshev *et al.* (1953) state that the incubation period of cutaneous leishmaniasis may be prolonged to four or even five years. They mention the case of a woman who lived continuously in Leningrad after leaving an endemic area (Kokand) in Soviet Middle Asia in August, 1943, and developed a leishmanial skin nodule in September, 1948—an incubation period of over five years.

In the following case the incubation period, judged by the same criteria as in the cases quoted above, was not less than 10 years.

CASE REPORT

The patient, a labourer aged 50, attended the dermatological out-patient department at St. Mary's Hospital, Portsmouth, on July 25, 1953, with a tender lump on the ulnar side of his right forearm which had appeared suddenly six months previously. There were no constitutional symptoms. The lesion was a brownish crusted intradermal nodule $\frac{1}{4}$ in. (0.6 cm.) in diameter. A tentative clinical diagnosis of lupus vulgaris was made and excision biopsy was performed.

Histology.—Sections showed a superficially ulcerated intradermal granuloma characterized by the presence of large numbers of parasites, nearly all within the cytoplasm of large vacuolated histiocytes. The parasites were morphologically identical with Leishman-Donovan bodies and the appearances were typical of cutaneous leishmaniasis.

Past History.—The patient served in Egypt, Libya, Palestine, and Syria from 1940 to 1942. At the end of 1942 he returned to this country and was demobilized in April, 1943; he had not been abroad since, and had been in good health during the whole period.

COMMENT

From the above case it is evident that cutaneous leishmaniasis must be considered in the differential diagnosis of nodular cutaneous lesions, even when the patient has not been exposed to infection for as long as 10 years. In all the cases mentioned above the evidence for the prolonged incubation period is circumstantial, but Berberian (1944) has shown in human volunteers that the appearance of lesions at the site of intradermal inoculation with suspensions of leishmaniae may be delayed for as long as four years seven