cells in the anterior chamber. There were no posterior synechiae. The patient had generalized glandular enlargement and numerous skin lesions. A skin biopsy confirmed the diagnosis of sarcoidosis.

Local cortisone treatment was given in the form of drops to both eyes, with in addition three subconjunctival injections of 6.25 mg. each to the right eye. There was rapid improvement in the vision with the disappearance of the precipitates. Three weeks after the start of treatment vision was 6/6 in both eyes. At the time of writing the patient's vision was still 6/6, but it had not yet been possible to stop applying drops, for cessation of treatment was at once followed by recurrence of inflammatory signs. Until three months previously, she was using cortisone drops, 0.5%, which she had to apply five times a day to keep the condition under control. For the last three months she had been using hydrocortisone drops, 1%, and twice-daily application of this had kept the iridocyclitis controlled. She has also had two courses of systemic cortisone therapy to control her generalized sarcoidosis during periods of exacerbation.

Treatment of Other Ocular Signs

With regard to the conditions other than iridocyclitis, no ocular treatment was given to the patient with Eales's disease or to the patient with macular oedema. The former's condition has remained stationary for two years, while that of the latter subsided spontaneously over a period of three months. The case of phlyctenular conjunctivitis was treated with atropine and sulphacetamide drops, clearing up after three weeks. Of the cases of keratoconjunctivitis sicca two were mild, though there was definite reduction in the quantity of tear secretion, as indicated by Shimer's test, and early filiform keratitis was present in each case. Symptomatic treatment, using Ringer-Locke solution, sulphacetamide, and methyl cellulose drops, brought considerable relief. One of these patients had moderately enlarged lacrimal glands, which returned to normal size over a period of eight weeks.

The third case had much-enlarged lacrimal glands, severe filiform keratitis, and much-reduced lacrimation. The enlarged glands subsided to normal size spontaneously after three months. The only beneficial treatment in this case was hydrocortisone drops applied to the conjunctival sac. These brought great symptomatic relief.

No treatment was effective in reducing the swelling in the case with infiltration of the eyelids, though systemic cortisone, local radiotherapy, and excision of the sub-cutaneous tissue were tried.

Summary

Of 100 patients with histological evidence of sarcoidosis 28 had ocular abnormalities. Iridocyclitis, the most serious manifestation, was present in 20 cases.

If untreated, sarcoid iridocyclitis may lead to blindness.

Early treatment of the iridocyclitis with cortisone or hydrocortisone offers the best chance of controlling the disease.

Treatment is unlikely to be beneficial when started at a late stage. Early diagnosis is therefore essential.

Methods which assist in arriving at the diagnosis of sarcoidosis are discussed and a brief description of the Kweim antigen test is given.

[Figs. 1-3 are reproduced by permission of the *Archives of the Middlesex Hospital*, in which they first appeared (Ainslie and James, 1953).]

REFERENCES

- Ainslie, D., and James, D. G. (1953). *Arch. Middx Hosp.*, 3, 71.
Crick, R., Hoyle, C., and Mather, G. (1955). *British Medical Journal*, 2, 1180.
James, D. G., and Thomson, A. D. (1955). *Quart. J. Med.*, 24, 49.

PHAECHROMOCYTOMA WITH SUSTAINED HYPERTENSION

BY

K. N. V. PALMER, M.D., M.R.C.P.
Senior Lecturer

AND

G. A. McDONALD, M.B., Ch.B.
Senior Research Fellow

From the Department of Medicine, University of Aberdeen

Phaeochromocytomas are functioning tumours of chromaffin tissue, the clinical syndromes which may occur being due to the presence of excess amounts of noradrenaline and adrenaline in the circulation (Goldenberg *et al.*, 1949). Patients may show either paroxysmal hypertension or persistent hypertension mimicking benign or malignant hypertension, with or without increased metabolism or glycosuria (Goldenberg, 1954). Only a quarter to a third of the cases have the classic picture of paroxysmal hypertension (Green, 1946), and a sustained hypertension is the commoner presentation. Usually, however, other manifestations that are uncommon in essential hypertension are associated with sustained hypertension due to a phaeochromocytoma: the most frequent of these are generalized sweating, pallor and coldness of the skin, goose-flesh, unexplained fever, postural hypotension and tachycardia, a normal cold pressor test, a raised basal metabolism, and glycosuria (Smithwick *et al.*, 1950).

Diagnostic Procedures

Adrenolytic agents cause a prompt fall in blood pressure, usually to normotensive levels, in cases of sustained hypertension due to phaeochromocytoma, but only during a hypertensive phase in the paroxysmal type of case. Phentolamine ("rogitine") has rapidly established itself as the adrenolytic agent of choice, for it is almost free of side-reactions, is easy to administer, and, unlike piperoxan, it does not produce a rise in blood pressure in cases of essential hypertension (Helps *et al.*, 1955). False-positive tests may occur if the drug is given intramuscularly (Gifford *et al.*, 1952) and when it is given during barbiturate sedation and in uraemia.

Provocative tests are mainly of value in the diagnosis of patients with the paroxysmal type of hypertension during a normotensive phase. In these cases intravenous injection of histamine causes an abrupt rise in blood pressure (Roth and Kvale, 1945). If a positive result is obtained with the phentolamine or histamine tests, the urinary excretion of the catechol amines (adrenaline and noradrenaline) should be estimated, and this is the most reliable diagnostic test for the presence of a phaeochromocytoma.

Treatment is surgical and consists in the removal of all tumours secreting adrenaline and noradrenaline. During operation two dangers have to be dealt with—namely, severe exacerbations of hypertension during the induction of anaesthesia, positioning of the patient, and handling the tumour; and a severe hypotensive phase when the adrenal veins are clamped before removal of the tumour. The first of these contingencies is met by the use of phentolamine, the second by the use of noradrenaline.

Case Report

A housewife aged 26 was admitted to hospital on April 24, 1955. She complained of frontal headache associated with attacks of sweating for one year. The headache was described as a dull ache, which usually came on in the morning and was made worse by stooping. The attacks

lasted about half a day and were not associated with sickness or gooseflesh. She had always looked pale, and suffered from lifelong constipation.

On examination she was thin, pale, and nervous. The thyroid gland was not enlarged. The skin was warm and moist. There were no signs of cardiac enlargement or cardiac failure. The blood pressure was 210/130 mm. Hg, and remained at this level during the period before operation. The fundi showed numerous small flame-shaped haemor-

FIG. 1.—Phentolamine test.

rhages, white fluffy exudate in both maculae, but no papilloedema. Neither kidney was palpable. She had a low-grade fever up to 100° F. (37.8° C.).

Investigations.—The urine contained albumin, 1 g./litre. Concentration and dilution tests, blood urea, and urea-clearance tests were within normal limits. There was polyuria, the average daily urine output being 3,500 ml. Intravenous pyelography revealed no abnormality. There was an area of calcification in the right loin. The haemoglobin was 85%, the white-cell count 11,400 per c.mm. (73% polymorphs). The sedimentation rate was 38 mm.

in the first hour. X-ray examination of the chest was normal. The B.M.R. was +22%. A dextrose-tolerance test (50 g. dextrose) showed impaired dextrose tolerance (resting level 140 mg. per 100 ml. rising to 210 mg. at one hour and 205 mg. at two hours). A glucose-insulin tolerance test did not show any evidence of insulin resistance. An intravenous phentolamine test (5 mg.) produced a fall in blood pressure from 215/140 to 115/65 mm. Hg (Fig. 1). Repeated loin-compression tests gave inconclusive results. The cold pressor test was positive (a rise in blood pressure from 200/120 to 240/150 mm. Hg being obtained). Bio-assay of the urine showed that it contained 2,730 µg. of catechol amines per 24 hours, of which 98% was noradrenaline (normal excretion of catechol amines is less than 150 µg. per 24 hours).

Operation.—On July 4, 1955, the tumour, weighing 65 g., was successfully removed by Mr. J. F. Philip through a right loin incision. Premedication: 1/150 gr. (0.43 mg.) of scopolamine and 5 mg. of phentolamine intramuscularly. Anaesthesia (Dr. R. Milne): thiopentone, suxamethonium, nitrous oxide, oxygen, and pethidine. The blood pressure during the operation was controlled by phentolamine and noradrenaline (Fig. 2). During induction of anaesthesia there was a sharp rise of pressure, which was readily controlled by phentolamine, 5 mg. intravenously. During mobilization of the tumour the blood pressure rose to 220 mm. Hg systolic. Phentolamine, 5 mg. intravenously, was given and the pressure dropped to normotensive levels. The total amount of phentolamine used during the operation was only 10 mg. As the blood supply from the tumour was interrupted there was a profound fall in blood pressure to 60 mm. Hg systolic, and a noradrenaline drip (4 mg. per 1,000 ml. in 5% dextrose) was started. The drip rate was regulated to maintain a blood pressure of approximately 120/70 mm. Hg. The total amount of noradrenaline given during operation was 5 mg. Noradrenaline was given for 11 hours after the operation at an approximate rate of 5 µg. per minute, and 50 mg. of cortisone was given intramuscularly six-hourly for 24 hours.

The post-operative course was uneventful; the patient was free of symptoms and the blood pressure remained normal. Histological examination verified that the tumour was a phaeochromocytoma. Biopsy of the right kidney taken at the time of the operation showed no abnormality. Analysis of the tumour showed that it contained noradrenaline, 3.08 mg. per g., and adrenaline, 0.16 mg. per g. The noradrenaline content was 94%, which accounted for the high pre-operative level of urinary noradrenaline. The urine excretion of catechol amines on the seventh post-operative day was normal (76 µg. per 24 hours).

The patient was readmitted eight weeks after operation. She felt very well, had gained a stone (6.4 kg.) in weight, and had lost her pallor. She had had no recurrence of the headaches or attacks of sweating. The blood pressure was normal (116/80 mm. Hg), and remained so during her 16 days in the ward.

She was afebrile. The urine contained a very faint trace of albumin. The urine output was now normal (1,500 ml. per day). The blood urea was normal. The area of calcification in the right loin was still present. The haemoglobin was 88% and the white-cell count 7,200 per c.mm. (normal differential). The sedimentation rate was 5 mm. in the first hour. The B.M.R. was +10%. A

FIG. 2.—Blood-pressure record during operation.

dextrose-tolerance test was normal. A provocative histamine test was negative and a cold pressor test was normal. The urinary excretion of catechol amines was normal (44 μ g. of noradrenaline per 24 hours). The fundi showed no exudates or haemorrhages.

Discussion

Persistent hypertension associated with attacks of sweating and pallor suggested the diagnosis of phaeochromocytoma in this case. It is desirable, as Rosenheim (1954) suggests, that all severe hypertensives should be screened for phaeochromocytoma. The phentolamine test enables this to be done readily, for it is simple to perform and does not cause side-effects in patients with essential hypertension. If this test is positive the diagnosis should be confirmed by an estimation of the urinary catechol amine excretion.

Phentolamine is of great value in the prevention of hypertension during the operation. It should be remembered, however, that in large doses it has a sympathicolytic effect (Meier *et al.*, 1949). Knox and Slessor (1955) suggest giving 20–30 mg. orally or intramuscularly 20 minutes before induction of anaesthesia, and 10–20 mg. intravenously during induction. They used this dosage in their patient who had paroxysmal hypertension, but it did not prevent a major rise in blood pressure following the injection of tubocurarine. They remark that doses of phentolamine in excess of requirements seem to be harmless, because rapid clearance of the drug soon allows normotensive levels to be reached. Our case showed a sustained hypertension, and because of this we concluded that it would be safe and wise to give 40 mg. of phentolamine intramuscularly 20 minutes before the operation. This was followed by a profound fall in blood pressure which lasted two hours and necessitated the postponement of the operation. We think it is safer to try the effect of increasing doses of phentolamine intravenously and intramuscularly before the operation in order to arrive at the right dose for each individual case.

Perirenal air insufflation was not done in this case. The right side was explored first for the following reasons: (1) the surgeon thought that he could feel a mass on the right side; (2) single tumours are commoner on the right side (Symington and Goodall, 1953); and (3) there was an area of calcification in the right loin which was thought to be in the tumour (although as the calcification was still present after the operation it was probably a calcified lymph node).

Reserpine. — Reserpine ("serpasil") when given parenterally has been shown to produce a hypotensive effect in most cases of essential hypertension; this comes on about half an hour after injection, is maximal at 2½ hours, and persists for from 4 to 5 hours (Freis and Ari, 1954; Hafkenschiel and Sellers, 1954; Finnerty and Sites, 1955). It is brought about by vasodilatation and a reduction in cardiac output, both actions being secondary to a central inhibition of reflex sympathetic activity (Trapold *et al.*, 1954a, 1954b). Bein (1953) was unable to find a peripheral sympathicolytic or ganglion-blocking action in animals, although McQueen *et al.* (1954, 1955) found in rabbits that, in addition to

diminishing reflex vasomotor responses, it did have a direct effect on the peripheral vessels. In our case the hypertension may be presumed to be due solely to circulatory pressor amines, since it fell to normotensive levels when a small dose of phentolamine was given. There was no significant fall in blood pressure when 2 mg. of reserpine was given intra-

FIG. 3.—Reserpine test.

muscularly (Fig. 3), which suggests that this drug has no adrenolytic effect in the human subject in seven hours in the dose given.

Chlorpromazine.—Chlorpromazine has been shown to possess adrenolytic and noradrenolytic properties in the dog, the hypertensive effect of adrenaline being abolished and the pressor effect of noradrenaline given in the same dose being reduced. It has little or no effect on the blood pressure of dogs (Courvoisier *et al.*, 1953), although in large doses in human beings it may cause postural hypotension. In phaeochromocytoma, chlorpromazine may be expected to

FIG. 4.—Chlorpromazine test.

FIG. 5.—Post-operative chlorpromazine test.

have a hypotensive effect by virtue of its adrenolytic properties. Knox and Slessor (1955) found that 25 mg. of chlorpromazine intramuscularly did not affect the height of a histamine-induced rise in blood pressure in their case, but abolished the symptoms, such as headache, pallor, and sweating, which were associated with the rise. In our case 25 mg. of chlorpromazine was given intramuscularly to the patient in the supine position. It produced a fall in blood pressure to normotensive levels which lasted for 4½ hours (Fig. 4). There was no significant fall in the blood pressure when chlorpromazine was given in the same dosage under the same conditions after the operation (Fig. 5). This suggests that chlorpromazine has a powerful adrenolytic and noradrenolytic action in the human subject, and that it may be used as an alternative to phentolamine in the premedication of patients before operative removal of a phaeochromocytoma. Dundee (1954) used chlorpromazine intramuscularly for this purpose, and found that, although it did not completely prevent a rise in blood pressure during positioning the patient on the operating table, it was very effective in preventing a hypertensive attack during handling of the tumour.

Summary

A case of phaeochromocytoma is described which showed sustained hypertension, increased metabolism, impaired dextrose tolerance, polyuria, and albuminuria. These findings became normal after removal of the tumour.

The use of phentolamine in the diagnosis and operative management and the effect of reserpine and chlorpromazine on the blood pressure are described.

Our thanks are due to Dr. G. B. West, of the Department of Pharmacology in the University College, Dundee, for assays of catechols in urine and tumour tissue.

REFERENCES

- Bein, H. J. (1953). *Experientia (Basel)*, 9, 107.
 Courvoisier, S., Fournel, J., Ducrot, R., Kolsky, M., and Koetschet, P. (1953). *Arch. int. Pharmacodyn.*, 92, 305.
 Dundee, J. W. (1954). *Brit. J. Anaesth.*, 26, 357.
 Finnerty, F. A., and Sites, J. G. (1955). *Amer. J. med. Sci.*, 229, 379.
 Freis, E. D., and Ari, R. (1954). *Ann. N.Y. Acad. Sci.*, 59, 45.
 Gifford, R. W., Roth, G. M., and Kvale, W. F. (1952). *J. Amer. med. Ass.*, 149, 1628.
 Goldenberg, M. (1954). *Ciba Foundation Symposium on Hypertension*, p. 91. Churchill, London.
 — Faber, M., Alston, E. J., and Chargaff, E. C. (1949). *Science*, 109, 534.
 Green, D. M. (1946). *J. Amer. med. Ass.*, 131, 1260.
 Hafkenschiel, J. H., and Sellers, A. M. (1954). *Ann. N.Y. Acad. Sci.*, 59, 54.
 Helps, E. P. W., Robinson, K. C., and Ross, E. J. (1955). *Lancet*, 2, 267.
 Knox, J., and Slessor, A. (1955). *Ibid.*, 1, 790.
 McQueen, E. G., Doyle, A. E., and Smirk, F. H. (1954). *Nature (Lond.)*, 174, 1015.
 — (1955). *Circulation*, 2, 161.
 Meier, R., Yonkman, F. F., Craver, B. N., and Gross, F. (1949). *Proc. Soc. exp. Biol. (N.Y.)*, 71, 70.
 Rosenheim, M. L. (1954). *British Medical Journal*, 2, 1181.
 Roth, G. M., and Kvale, W. F. (1955). *Amer. J. med. Sci.*, 210, 653.
 Smithwick, R. H., Greer, W. E. R., Robertson, C. W., and Wilkins, R. W. (1950). *New Engl. J. Med.*, 242, 252.
 Symington, T., and Goodall, A. L. (1953). *Glasg. med. J.*, 34, 75.
 Trapold, J. H., Osborne, M. W., Plummer, A. J., and Yonkman, F. F. (1954a). *J. Pharmacol.*, 110, 49.
 — Plummer, A. J., and Yonkman, F. F. (1954b). *Ibid.*, 110, 205.

The proceedings of the National Smoke Abatement Society's conference at Bournemouth last year have recently been published (price 12s. 6d. net from the Society, Palace Chambers, Bridge Street, London, S.W.1). They include Sir Ernest Smith's presidential address, papers on recent progress in domestic heating by centralized methods, on local authorities and the Beaver Report and the Clean Air Bill, and by members of the Beaver Committee. The Des Vœux Memorial Lecture was given by Dr. A. Parker on "The Destructive Effects of Air Pollution on Materials."

SERUM IRON IN THE DIAGNOSIS OF HEPATOBIILIARY DISEASE

BY

L. SCHAMROTH, M.B., F.R.F.P.S., M.R.C.P.Ed.

W. EDELSTEIN, M.B., B.Ch.

General Hospital, Johannesburg

W. M. POLITZER, M.D.

AND

N. STEVENS, B.Sc.

South African Institute for Medical Research, Johannesburg

The differentiation between obstructive and non-obstructive jaundice may sometimes be difficult in spite of the laboratory aids at present available. It is therefore interesting to note reports of an association between viral hepatitis and an abnormally high serum iron level. Earliest publications were those by Warburg and Krebs (1927) and Locke *et al.* (1932), who each described a single case of "icterus" in which a raised serum iron level occurred. Hemmeler (1939) first reported that the serum iron was elevated in "catarrhal icterus" and normal in obstructive jaundice. Since then only a few reports have appeared in the literature on this subject.

The present study was undertaken to investigate the value of the serum iron determination as a routine procedure in the diagnosis of hepatobiliary disease.

Methods and Material

Over a period of 15 weeks 104 serum iron studies were conducted on 63 patients with hepatobiliary disease. This group consisted of 21 cases of acute infective hepatitis, 8 cases of homologous serum hepatitis, 6 cases of "subacute viral hepatitis" (defined below), 14 cases of cirrhosis of the liver, 12 cases of extrahepatic obstruction, and 2 cases of chlorpromazine "hepatitis." There was also one case of haemochromatosis.

The diagnosis of viral hepatitis was based on the history of a constitutional disturbance (anorexia, nausea, lassitude, generalized body pains), pyrexia, tenderness in the right hypochondrium, jaundice, and regression of the disease on conservative treatment. If injections had not been given during the preceding six months the cases were termed "infective hepatitis" (endemic viral hepatitis), whilst those that had had injections two to six months before the onset of the illness were designated "homologous serum hepatitis." None of the patients had been given a blood transfusion or had received iron therapy prior to the onset of the disease.

Specimens were collected at 9 a.m.—that is two hours after the morning meal. Before any glassware was used in the laboratory (including the tubes into which the blood was collected) it was cleaned with chromic acid, then with concentrated nitric acid, and rinsed well with glass-distilled water. The serum iron was determined by the method of Kitzes *et al.* (1944), which takes approximately two hours to perform; several estimations may be carried out simultaneously. Our normal range for this method is 70–185 µg. per 100 ml.

Results

The results are recorded in Figure 1 and Table I. At the initial estimation, which was carried out during the first week of admission, 16 out of 21 cases of infective hepatitis had serum iron values above 220 µg. per 100 ml. The mean value of the 21 initial estimations was 292 µg. per 100 ml.