However, I can only say that as a result of our recent experiences I would be sorry indeed to be without this drug again at the present time. And if frenquel, used alone or combined with E.C.T., helps only a small limited group of cases of depressions and early schizophrenics, and we can learn to pick out these groups clinically, it will still be very much worth while.

Conclusion

May I again stress the importance, with all these new chemical tranquillizers flooding the market, of trying to find the limited and definite indications for any new one used. We must not go on prescribing these drugs at random, replacing one by another without any rhyme or reason. We should always be on the lookout for even the smallest clues to help us in our differential search.

Even several new treatment swallows will hardly yet make a psychiatric summer. Some people, especially in America, have been talking about a new golden era in psychiatry, when it will be possible to give up insulin coma, leucotomy, and E.C.T., and have the chemical tranquillizers as humble handmaidens of universally applicable forms of dynamic psychotherapy for all types of neurotic and psychotic patient. I venture to predict that those psychiatrists who are saying this sort of thing will prove to have been more psychotherapeutically brainwashed than their patients will ever be. What would really happen then is just what was happening twenty years ago. All we had in those days were Freud, Adler, and Jung's psychotherapeutic methods of treatment and the bromides, paraldehyde, and longer-acting barbiturates to help us in the treatment of all types of mental disorder—except G.P.I., for which malaria therapy had become available. Few of the modern physical treatments, such as insulin, E.C.T., leucotomy, the synthetic vitamins, endocrines, and the like, which have made the real difference in actual practice, were then in general use in mental hospitals. The result was that private paying patients and a highly privileged small group of hospital patients got all the psychotherapy, while most of the patients in ordinary mental hospitals were simply kept chemically tranquillized—or “drugged,” as we used to call it in those days—on bromides and paraldehyde. We must not revert to this same old state of affairs by simply substituting new tranquillizers for the old.

A better era in psychiatric treatment is really likely to be achieved only when we start to learn that every method of treatment, empirical and otherwise, not only has its uses but also its very real limitations. There will be no treatment found universally applicable to all groups of patients. Progress in psychiatry will occur gradually and piecemeal, in exactly the same way as has happened in general medicine and surgery. It will certainly not be by the finding of any universal panacea, chemical or psychological. We must therefore go on keeping a very open mind about the value of all sorts of different treatment methods, and try to keep as many of them as possible in *constant* use. Then we shall always be able to compare new methods and new tranquillizers with the old. And it will be found, as elsewhere in life, that “what is new is not always true, and what is true is not always new.”

REFERENCES

- Barbour, R. F., Pilkington, F., and Sargant, W. (1936). *British Medical Journal*, 2, 957.
British Medical Journal, 1936, 2, 1050, 1242, 1286.
Fabing, H. D. (1955). *Neurology*, 5, 319.
— and Hawkins, J. R. (1955). *Diseases of the Nervous System*, 16, 3.
Rinaldi, F., Rudy, L. H., and Himwich, H. E. (1955). *Amer. J. Psychiat.*, 112, 343.

ANXIETY STATES*

BY

GERALD GARMANY, M.B., B.Sc.
M.R.C.P., D.P.M.

From the Psychiatric Department of Westminster Hospital

The purpose of this paper is to present some findings from the records of Westminster Hospital rather than to give a textbook account of anxiety states, though inevitably something must be said about the nature and aetiology of anxiety. Some years ago a front page to the out-patient notes was devised, and the primary purpose of this was a pragmatic one. It was intended that the essential points of any particular case should be easily ascertainable when a patient was seen for the second or third time. A formulation of the case would have been incorporated in a letter sent to the doctor who had referred the case to hospital, and this front page in effect broke down and analysed the letter. In fact, of course, the analysis came first and the letter was usually composed some hours later. Although this was the primary purpose of the analysis the possibility of future research was in mind, and was encouraged by the fact that the hospital is fortunate in having a records officer complete with a Powers-Samas machine. It was not, however, envisaged that anxiety states in particular would be investigated, and, indeed, the results it presented could have been furnished quite easily in respect of any other syndrome.

These facts are mentioned in so far as it is relatively unusual to present work which has been done without motive, and which has been subsequently computed without any unconscious distortion by the observer, since it was done by a machine. The investigation has revealed one extremely irritating defect in the transcription sheets—namely, that by some strange oversight it does not differentiate between male and female. This will be apparent in what follows in that sex distribution is not available.

Nature of Anxiety

By analogy with the generally accepted definition of depression, an anxiety state may be defined as a clinical condition dominated by an affect resembling fear which is not transitory or slight, and is not secondary to some other condition. By “clinical condition” is meant an illness in the usual sense—namely, that material discomfort is caused by it, or that ability to work and maintain social relationships is materially interfered with. The affect of anxiety resembles fear but differs from it in the following particulars. Firstly, fear is usually related to appreciated danger, and to that extent has the quality of normality. Anxiety, on the other hand, often appears inappropriately and in relation to no particular threat, and to that extent seems to the patient to be pathological. Secondly, fear is more often bodily whereas anxiety is not, and the patient usually shows his appreciation of this fact by saying that in physical matters he is not in the least nervous. Thirdly, fear is for a term and the end of it can be seen, while anxiety may offer no such hope. Lastly, the individual seems to be much more totally involved in anxiety, and the patient does not seem able to survey himself dispassionately as he can in fear.

The other item in the definition which requires brief scrutiny is the primary nature of the condition. Anxiety states are often diagnosed when other conditions are basic

*A shortened version of a Maudsley Bequest Lecture read to the Royal Medico-Psychological Association on November 7, 1955.

and the anxiety is a secondary phenomenon. This is particularly the case with depressions, and nearly all "acute anxiety states" referred urgently to hospital turn out to be in this category. Schizophrenics may generate severe and overt anxiety as a response to their inner experiences. Free anxiety in a young person combined with marked ideas of reference is always suggestive of this disease. Anxiety may be a secondary result of missed organic disease. Patients with carcinoma of the lung or of the colon, for example, sometimes seem to have a conviction that all is not well even at a stage when investigations are negative, and there is something a little worrying about the man who quietly and without dramatization says that some part of him "just doesn't feel right" and that he feels he has cancer.

The clinical features of anxiety states are well known, but the importance of tension in the voluntary musculature is stressed not only because it is invariably present but because it may outlive the original anxiety state by producing bad postural habits which perpetuate the condition. Inability to relax, restlessness, fatigue, and difficulty in getting off to sleep are usually present. Chronic overactivity of the autonomic nervous system is present, but its forms are many and varied, and these are presumably decided by such factors as familial predisposition, conditioning in childhood, and special valuation of the part as in the case of footballer's knee and the pianist's hand. It is not helpful to think of sympathetic or parasympathetic predominance, for both divisions are overactive and in a state of unsettled and varying equilibrium with one another. Indeed, this very variation is itself a source of anxiety in the man whose colour changes quickly from pink to pallor, and who alternates between a tiresome constipation and a troublesome diarrhoea. Work such as that done on the stomach by Wolf and Wolff (1942) shows very clearly how anxiety may produce somatic dysfunction which in its turn causes morphological change.

It has to be remembered that an anxiety state may not always be a continuous illness of fairly steady intensity. It may be punctuated by episodes of very intense apprehension amounting to panic—what are called anxiety attacks, panic attacks, or phobic attacks. In between these attacks the patient sometimes feels very well. I am sure that these attacks are frequently misunderstood. I do not think they should be included, as they sometimes are, among the manifestations of an obsessional neurosis, for the phobias of an obsessional neurotic are quite different. The latter are fearful attitudes lasting months or years until remission occurs, and quite unlike the acute and short-lived episodes called anxiety attacks, though these are particularly likely to occur in obsessional personalities, as opposed to sufferers from an obsessional neurosis. It is not at all common to see anxiety attacks unless the doctor has his consulting-rooms high up in a house served by a small lift, when he can be assured of a steady supply of sufferers. It is almost as necessary for a psychiatrist to work on the ground floor as it is in the case of a cardiologist.

The search for homeostasis may be assisted by the development of hysterical or obsessional mechanisms as well as by the taking of sedatives or alcohol. When hysterical mechanisms are well developed the evaluation of the underlying tension state is of great importance.

A woman of 40 years was seen in 1950. She had been virtually bedridden for the last twelve months, and the general picture was one of hysterical indifference. There was, however, a history of depression and she had a marked fine tremor of the hands. Her blood pressure was 200/115, and it is interesting to note that it had been observed to be 220/110 in 1945. In view of the signs of tension, prefrontal leucotomy was undertaken with dramatic recovery. She was followed up over a period of three years, during which time the blood pressure was in the neighbourhood of 170/90, and she was symptom-free.

Aetiology

It is in line with responsible English thought to regard an anxiety state as an interreaction between a constitution which is in some way predisposed and operative existing stress

factors. As shown during the war every possible combination may be present. Now as then, some are so predisposed that the ordinary business of living produces breakdown while others show the most remarkable resistance to all that life can do to them. Each individual must maintain some sort of equilibrium between the demands and pressures of the outside world, the requirements of his conscience, and the need for satisfaction of his instinctual drives. The stress factors derived from the outside world are clear enough and include such factors as domestic relationships, work, social contacts, financial difficulties, housing problems, and physical disease. Both conscience and instinctual drive are included in the list of stress factors under the heading of personality when it was thought that that was the right course. This may not commend itself to everyone, but, as the records are nearly all compiled by me, uniformity at least can be claimed for them.

It is necessary to explain how it was decided whether personality should be regarded as part of constitution or as a stress factor in its own right. Two examples may make the issue a little clearer. Homosexuality is a constitutional anomaly, or so I believe, but it is not one which of itself predisposes to anxiety. Given certain circumstances it may be lived with without tension, and indeed this is sometimes the case. On the other hand, if a homosexual marries or joins the armed Forces, then he is likely to find his anomaly a stress factor of considerable magnitude. An obsessional personality is again a constitutional matter, but in the circumstances of modern living it is a stress factor in that it creates difficulties for itself wherever it goes. This particular method of describing cases has been adopted in order to avoid the usual difficulty of begging the question, as most assessments of constitutional predisposition tend to do.

Constitution has been taken in its ordinary meaning of mental and physical make-up, and the existence of "loading" has been inferred from a number of things. Thus a strong family history of the same condition indicates predisposition, and where constitutional loading has been described as "notable" this history has usually been present. Loading has also been inferred from the personal history when anxiety was present at the period of life when it is usually uncommon and where the tendency has dominated the individual, however well things went for him.

Table I shows the incidence of anxiety states in a consecutive series of 644 cases seen at Westminster Hospital. It will be seen that anxiety states account for 25% of the whole.

TABLE I.—Distribution of 644 Consecutive Cases

Anxiety states	158 (25%)
Phobic states	33
Reactive depression	94
Endogenous depression	107
Involuntional depression	12
Obsessional states	14
Hysteria	28
Somatization reactions	23
Psychopaths	29
N.A.D.	36
N.Y.D.	2
Other	108

This figure can be compared with the results obtained by other workers only if some uniformity is observed in the criteria to be applied. Thus a current textbook of great authority states that anxiety states are relatively rare, but the discrepancy is explained when we read that "hysterical elaboration of minor anxiety symptoms and depressive states with anxiety features" are far more common. The same book remarks that it may be a matter of taste whether the diagnostic label is anxiety state with hysterical features or hysteria with anxiety features. The Geneva International List offers a simple way out of this difficulty, and it would be better that everyone should follow it. Under this arrangement all cases in which tension is evident would be included with anxiety states whether hysterical features were present or not. Hysterical reaction has a separate heading which is called "Hysterical Reaction without Mention of Anxiety Symptoms." This heading includes such conditions as anorexia nervosa, hysterical blindness, fugues, and other

conditions where tension may be quite absent. With these rigid criteria for hysteria, the number is relatively small, as will be seen.

The total number of depressions is a third higher than that of anxiety states. It has been thought worth making a distinction between reactive depression and endogenous depression, even though it is not always easy. The differentiation between anxiety states and reactive depression is much more straightforward despite the fact that the reactions tend to partake a little of each other. In depression sadness dominates the picture, while in anxiety tension is to the fore. The somatization reactions are relatively small in number, merely because these were diagnosed by colleagues; I have a great aversion to the term.

Table II shows the incidence of constitutional loading. If one remembers the restrictive criteria applied to constitution, the number falling within the first two groups is impressive. In practice, "possible" has been equated with "likely," and there has always been some lead available.

TABLE II.—Incidence of Constitutional Loading in 158 Cases of Anxiety States

Notable	36	61%
Possible	62	
Unlikely	57	
Not recorded	3	

TABLE III.—Stress Factors

Personality	111	(70%)
Domestic	56	(35%)
Work	32	(20%)
Other social factors	16	(10%)
Financial	14	(8.7%)
Housing	15	(9.4%)
Physical	43	(27%)
None ascertained	14	(8.7%)

The stress factors which were thought to be operative are shown in Table III. They were not recorded merely because they were there, but only if it was thought that they were operative in producing the neurosis. The factors are not mutually exclusive, and in some cases two or more were present, which is why the total exceeds 158. It will be seen that personality factors were considered important in 70% of the cases, and of these factors an obsessional character was by far the most common. This illustrates the dilemma to which reference was made previously. An obsessional personality has been regarded as a stress factor rather than a predisposing constitutional anomaly when it has begun to produce effects at a given time, as when a man changes his job, or when he can no longer meet the perfectionist demands he makes upon himself either because of added responsibility or because of reduced adaptation, as when he is becoming arteriosclerotic or develops some other physical illness. The obsessional personality creates stresses wherever it goes because of the high standards it demands and its difficulty in delegation. An example might be afforded by the obsessional housewife who copes very well in a flat, and then develops anxiety symptoms after moving into a house; or a patient once seen recently who coped very well with his six companies, but developed bad panic attacks when his father died and left him another nine.

Table IV shows the age distribution. It will be seen that the diagnosis is uncommon in old age, and not very frequent in adolescence. A peak is reached in the fourth decade,

TABLE IV.—Age Distribution (158 Cases)

Age Group (Years)						
0-20	21-30	31-40	41-50	51-60	61-70	71-80
8	36	66	28	15	4	1

when the combination of youthful zeal and increasing responsibility is at its most deadly. There is therefore good support for the widespread belief that life begins at 40.

Treatment

Drugs were relied upon as the principal approach in about one-third of the cases (Table V). This was because the patients had too far to travel for other treatment, because

they could not get the necessary time off work, or because they lacked the necessary intelligence or persistence for anything else; and sometimes it might be because the condition responded satisfactorily to sedative and it did not seem

TABLE V.—Disposal of Cases

Consultation	28	74=50%
Counselling	5	
Supportive therapy	24	
Intensive therapy	17 (11%)	
Group therapy	4	
Relaxation	54	
Environmental adjustment	3	
Hospital	3	
Physical treatment	1	
Drugs	52	
Referred elsewhere	4	

necessary to do more. The principal drug used is amylobarbitone or its sodium salt, because it is quick in action and safe to issue. Patients are advised not to mix drink, drugs, and driving as a routine, though the dangers of mixing amylobarbitone sodium with alcohol have probably been exaggerated. The same cannot be said about quinalbarbitone. Patients with episodic anxiety, or who have panic attacks in well-defined situations which can be anticipated, can with advantage carry amylobarbitone about with them, and take $\frac{1}{4}$ or $1\frac{1}{2}$ gr. (50 or 100 mg.) about 30 minutes before the expected ordeal. So far as chlorpromazine is concerned, it is perhaps most useful as a synergist to amylobarbitone in very severe tension. There is no doubt that it produces sedation without clouding more selectively than other drugs, but the incidence of complications is substantial enough to deter unnecessary use.

Reserpine and rauwolfia have been the subject of a controlled experiment at Westminster Hospital, and the results were recently published by Folkson and May (1955). The results do not, on the whole, encourage the use of these drugs in the treatment of anxiety states, except perhaps when hypertension is present as well.

About half of our cases were given psychotherapy, using the term in its broadest sense. At the most simple level ventilation has a place, particularly with older people, who feel better for talking, without particularly desiring advice. Others require what is sometimes facetiously called "reassurance." The role of assurances and explanations is in allaying secondary misinterpretations and apprehensions, which are perhaps rather commoner than they used to be because of the increase in popular articles dealing with tension and other psychiatric disorders. As a therapeutic weapon for dealing with the biological effects of morbid anxiety, reassurance is of little help. The man with tension headaches is not usually helped by being told that he has no tumour if his headaches go on, and they usually do.

Ordinary counselling has a definite part to play, though it may not be a very exciting one. Patients may not always be very bright, or their judgment may be distorted and simple advice may be required about everyday things. The view has been taken that it is right to give advice when it is sought, and when, in addition, ordinary concepts of morals and ethics are not cut across. A dissenting man and wife may be advised to come together, but no psychiatrist should advise them to separate. Parents may be advised to stop beating their children, but not to start. There can be no objection, it is felt, to intruding into religious questions if some misunderstood issue can be clarified without challenging the patient's values directly and without getting oneself identified with the Devil's Advocate. If this is done it is necessary to have some acquaintance with comparative religion and some sympathy with spiritual aims. Given these things there is no more reason for keeping clear of religious issues than would apply to any other problem in the patient's mind.

I have no great faith in the importance of the repressed unconscious conflict or, at least, have not found the concept of great assistance in my work. The opposite applies to the conscious or near-conscious dilemma which is usually present

if it is searched for. A typical dilemma may be illustrated by the nocturnal panic attacks of the woman torn by resentment for some reason and anxious to punish her husband by refusing him intercourse; while at the same time anxious about the integrity of her marriage if she pursues her desires. It would be ingenuous not to accord such a situation the importance it deserves. It may be found, and indeed it frequently is, that she adopts the course of alternately accepting and rejecting one of the two possible courses of action, and a self-perpetuating cycle is thus established. The habit of switching from one solution to the other may be broken once it is realized how anxiety is being caused, and the psychiatrist may be called on to give a clear lead if he legitimately can.

With an intelligent patient it is worth spending a little time showing how morbid anxiety is established, and how in its turn it produces somatic dysfunction. This is necessary if he is to collaborate with any enthusiasm in the future treatment of his case. He must be able to see where he is going and what the purpose of his efforts is to be. It must not be forgotten that, although only a minority of psychiatrists in this country are analytically orientated, the same does not apply to the general public, who are Freudian to a man. A good deal of time may have to be spent explaining why infantile experience is being ignored. Indeed, it is said that there is now a cult which concentrates on intrauterine experience, and a young woman recently asked that we should discover what her parents talked about when she was a foetus.

We accept it as axiomatic that attitudes and sentiments can be taught to a child, and forget that a similar technique may be taught to an adult. This re-educational or habit-forming psychotherapy requires certain conditions. First, the patient should understand in concrete rather than abstract terms what is being aimed at, and this objective should be framed in positive rather than negative terms. Keeping calm instead of getting panicky is too abstract, and not rushing about the place has the disadvantage of being negative. Secondly, he must consider very carefully whether he objects to the habit, for if he does he will never acquire it. Thus it would be pointless to suggest gentleness to a professional boxer, and dangerous to invite more extroverted aggression from a nursemaid. Thirdly, the desirability of the habit must be sold to him so that he will really want it. The conditions that govern all human endeavour have then been set up.

The usefulness of relaxation therapy in the treatment of anxiety states is becoming increasingly known (Garmany, 1952). The technique involves the teaching of a precise motor habit and is not to be confused with the relaxation of the analyst's couch, which is quite a different matter. The teaching is based on the fact that tense muscles in anxious patients show continuous action potentials, and that these may be dispelled concurrently with the symptoms. The ordinary patient is quite incapable of relaxing to order, and must be taught in the same methodical way that would be shown by a golf professional who hoped to implant a swing. At Westminster Hospital it has been found convenient and effective to entrust this work to physiotherapists who have been specially instructed in what is required, and small classes of six are taken twice a week. There is a separate class for each sex, and a separate class for beginners. The patients who benefit most are those who are bright, obsessional enough to have the drive to master the technique, and middle-aged. In theory patients benefit from both relaxation and from re-educative psychotherapy. In practice it has been found that younger patients stand immobilization less well than those over 40. Conversely, they learn new mental habits more easily, as might be expected. Patients with recently developed panic attacks, when these do not herald a depressive illness, do very well with relaxation.

It will be noted that there is no mention of prefrontal leucotomy in the treatment of anxiety states. This quite truly indicates a belief that the indications for this operation in tension states are very few and far between. There are,

however, occasions on which the operation has been recommended and carried out, but such a case did not occur during this series.

Summary

A study has been made of 158 patients with the diagnosis of anxiety state, which represented about 25% of a consecutive series of patients seen in the out-patient department of a teaching hospital. All cases showing tension were included, whether hysterical features were prominent or not, following the Geneva International List.

The incidence of constitutional predisposition was found to be very high even when restrictive criteria were applied in assessment. The operative stress factors were far more commonly to be found in the personality itself than in external factors, such as housing or domestic difficulties. The incidence of tension states is highest in the fourth decade, and relatively low in adolescence and old age.

The use of drugs in treatment is discussed together with a consideration of the forms of psychotherapy found most useful in this condition.

REFERENCES

- Folkson, A., and May, A. R. (1955). *British Medical Journal*, 2, 1121.
Garmany, G. (1952). *Muscle Relaxation*. Actinic Press, London.
Wolf, S., and Wolff, H. G. (1942). *J. Amer. med. Ass.*, 120, 670.

"MERATRAN"

A NEW STIMULANT DRUG*

BY

WILLIAM G. A. BEGG, M.B., Ch.B., D.P.M.

AND

A. ARNAUD REID, M.B., M.R.C.P., D.P.M.

Research Registrars, Department of Psychological Medicine,
St. Thomas's Hospital, London

There have been a number of reports in the medical press of the United States on the properties of α -(2-piperidyl) benzhydrol hydrochloride ("meratran") and its value in the treatment of depression and fatigue states. Brown and Werner (1954) found that in the larger animals, such as dogs, meratran produced a marked increase in purposeful activity which, although it was more intense than that produced by amphetamine, was not accompanied by the irritability and anorexia produced by the latter. Furthermore, there were no changes in the blood pressure, heart rate, or respiratory function which could be attributed to the drug. There was no depressive phase following stimulation, and the margin of safety was wide. Himwich and Rinaldi (1955) thought that the site of action of the substance was probably on the reticular substance of tegmentum, while Heath (1954) found that meratran, unlike any other known drug, produced rapid high-voltage spikes in the region of the septum of monkeys.

In view of its stimulant powers and its ability to awaken rabbits from barbiturate-induced sleep, Fabing (1954) decided to try the effects of meratran upon cases of narcolepsy. He found it to be highly effective for this condition, and subsequently decided to try it on cases of mild depression (Fabing *et al.*, 1955). They

*This work was done with a grant from the Endowment Funds of St. Thomas's Hospital.