EMERGENCIES IN GENERAL PRACTICE**"DRUNK IN CHARGE"**

BY

A. B. DAVIES, M.B., B.Sc.*General Practitioner, Walsall; formerly Police Surgeon*

Whatever the public impression may be as to the state of the "drunk-in-charge," his clinical condition for certification purposes generally falls considerably short of the music-hall picture. The present legal definition refers to him being at the material time "under the influence of drink or drugs to such an extent as to be incapable of having proper control of a motor vehicle."

Answering the Call

In most areas there are duly appointed police surgeons who are well accustomed to conducting medical examinations for this purpose. Nevertheless every medical man and woman is at risk for such a call, or may be asked to give an opinion on fitness to drive even when actually giving medical or surgical treatment in an emergency. For example, a hospital resident should be most careful and cautious if asked by the police to give an opinion on the sobriety or fitness-to-drive of a person brought in as an accident case. He can refuse to give such an opinion, though he may be subpoenaed to give evidence if required. Only doctors officially appointed to the police are obliged to answer calls to examine an alleged drunk. It is not part of the N.H.S. contract to do so, and a fee for such service may be charged to the police authority or to the patient requesting his attendance. Nevertheless many doctors will attend out of a sense of public duty or to see fair play for their own patients. Sometimes the doctor's knowledge of his patient's past medical history and clinical condition will be of great advantage to the defendant.

In these cases the defendant will have been brought to the police station as a result of his alleged unusual conduct when driving or because he has been involved in an accident. The function of the medical examiner is to determine whether there is a medical cause for this behaviour or any explanation other than the effects of alcohol. The earlier he attends the more accurate is his opinion likely to be about the patient's condition at the time of arrest. The time of arrival must be noted and accurate detailed notes recorded from then on. If, after his examination, the doctor has any doubt about the medical condition of the patient, he should not hesitate to arrange for the necessary observation and treatment. Intoxication and intracranial haemorrhage may coexist, and the results of missing the latter can be disastrous.

General Examination

The patient's consent to being examined must be obtained. It is an assault to examine without this. Unless this is given the notes should record only external observations and words spoken.

The examination should be conducted in a large, warm, well-lit room with maximum privacy—not in the charge-office. On the way to such a room (perhaps the magistrates' room) the doctor should follow the accused so that he can observe his gait and manner of going throughout doorways, turning bends, and going up or down steps, if such hazards

are available. Within the room will be the doctor and the accused. The latter should always be told that he has the right to call his own medical practitioner if the examining-doctor is acting in an official capacity. He may also request his solicitor. The presence of one police officer is usual, but he takes no part in the examination and makes no comment. It must be remembered that the defendant is in custody. The officer will be of assistance in finding writing materials, taking the patient to the lavatory, etc.

From the moment the patient is first observed the doctor will have noted any objective signs such as may be indicated by his colouring, demeanour, speech, odour of his breath, stance, gait, or the state of his clothing, including the presence on it of blood or vomit. Nevertheless a full and comprehensive examination must be methodically conducted, with particular attention to the heart, lungs, and nervous system. Negative findings are just as valuable in evidence as positive, and should be recorded in view of the possibility of other explanations for the accused's behaviour being put forward at a later date. There should be general questions about his previous state of health, when a doctor was last seen and for what condition, whether there has been service in the Forces and if so the grade on discharge, and whether he has been accepted for life insurance. The occupation should be noted. He should be asked if he feels ill at present and if he has recently been injured. These questions aim at eliciting information about any pre-existing condition which may have a bearing on the present state of the accused.

A thorough physical examination may show the presence of needle punctures as in diabetes or drug-addiction. The later examination of the urine may disclose glycosuria, albuminuria, or the presence of ketones. Heart disease, hypertension, asthma and bronchitis, anaemia, mental disorders, organic nervous disease, pyrexia, and other significant physical conditions will usually be detectable by efficient examination assisted by pointed questions. Full allowance must be made for any of these conditions if present, even to their acceptance as being fully responsible for the incapacity in driving. The taking of drugs for the treatment of disease is sometimes put forward as an explanation of the patient's condition: opiates, belladonna, anti-asthmatic and antispasmodic drugs, barbiturates, antihistamines, and even alkalis have all been claimed to have been responsible. Special care must be taken in examining for head injuries. Concussion may be pleaded later, and the examiner should be on his guard for the slowly progressive effects of an intracranial haemorrhage, and indeed for any cerebral catastrophe. Major and minor epilepsy are occasionally suggested, but this admission presupposes a wrongful declaration in applying for a licence. The possibility of malaria, and the effects of petrol vapour or exhaust fumes, must also be borne in mind. Certain vague terms such as "shock," whether surgical or emotional, and "black-out" may be used in court. The medical examination should make it possible to answer such pleas. The temperature, pulse, blood pressure, and respiration rate should all be recorded, the pulse rate at both the beginning and end of the examination (see below).

Finally, it is wise to make a note of the state of the weather at the time of examination. The case may not be heard for months and recollections may not then be clear. Conditions such as fog, snow, rain, extreme heat, or excessive cold may be relevant to the case.

Signs of Alcoholic Intoxication

General Demeanour.—Alcohol affects different people in different ways, and sometimes demeanour may change in the same individual during examination. Some are offensive and abusive, others hilarious and facetious, a few maudlin and depressed. Mood may change from full co-operation to hostility and abuse. Some are garrulous and noisy. Almost without exception these subjects think they have done well in their tests. It is this confidence which makes some of them say, "I don't need any other doctor; I'll be satisfied with you."

Aspect.—The skin is usually flushed. The upper eyelids appear "heavy," the whites of the eyes muddy and blood-shot. The pupils are dilated in varying degrees with varying responses (generally sluggish) to light. The pupil may redilate after contraction and there may be occasional lateral nystagmus. There may be evidence of injury, with wounds and blood stains. The clothing should be examined for damage, blood stains, or vomit. The patient may vomit during the examination, and the vomit may smell of drink. Buttons may be torn off clothing. The trouser or coat buttons may be fastened in the wrong holes.

The Breath.—The smell of drink, whether fresh or stale, is positive evidence that the patient has taken drink, but no more. It is no index of the quantity or type consumed. It must be remembered that he may have been given a "nip" by a kind bystander after an accident. The mouth and throat should be examined. After heavy drinking the throat and tongue are red and dry, but there may be furring as well. Broken dentures or a cleft-palate may affect the speech. Attempts may be made to mask the odour with peppermint or chlorophyll.

Pulse and Respiration.—The pulse rate is always increased by alcohol. The rate is substantially maintained during the examination according to the amount consumed. It should be counted at the beginning and end of the examination and recorded. The respiration rate also increased with drinking.

Eyes.—In addition to the preliminary observations already mentioned, there is also the question of impaired vision. Are glasses usually worn? Were they being worn at the time of the offence? Has the patient his glasses with him, either being worn or in his pocket? Rough tests should be made for distant vision, as well as a reading test for near vision, the patient being allowed to use his glasses if he normally does so. Ophthalmoscopic examination may disclose abnormalities of the optic fundus such as retinitis, arteriosclerosis, or a choked disk.

Co-ordination.—In driving, good co-ordination and a quick reaction-time are most important. Some opinion on the patient's co-ordination will already have been formed from the observations previously made. Balance is important, and Romberg's test should be performed. Any tremors should be elicited. The gait should be further tested by asking the patient to walk across the full length of the room, turn round, and return. (It is hardly fair to use a white line.) All kinds of abnormal gait may be seen, ranging from the rolling, lurching variety as if on a ship's deck in heavy seas to the stamping, military exaggeration of the over-confident. Reeling and lurching are common on turning. On going through a doorway the subject often brushes against the door-posts. Picking up a match is an excellent test of co-ordination. Apart from complete failure there may be groping and fumbling, scraping of the fingers along the floor, and often an indirect approach instead of the normal direct and speedy movement. In dressing after examination the co-ordination is well tested in tying shoe-laces, fastening shirt buttons, braces, or belt. Handwriting is discussed below. It is wrong to ask the patient to perform any acrobatic or other test with which the examiner himself might find difficulty.

Speech and Reading.—Ample opportunity will already have been obtained to note abnormalities of speech. If

there is yawning the time of the last sleep should be ascertained. Hiccup should be noted. The reading should be tested in a good light and a daily newspaper used. The patient should be allowed to make his own selection, as it may be said afterwards that a particularly difficult piece had been selected by the examiner. The parts read should be marked with crosses at the four corners for production in court if necessary. Allowance must be made for illiteracy. The commonest faults are lisping and slurring, but there may be frequent repetition of words and syllables. Sometimes words or whole lines are omitted, at others extra words are added. Allowance must be made for peculiarities of speech due to race or dialect.

Writing Test.—It is helpful to obtain the consent to examination in writing. This provides both proof of consent and a very good test of handwriting. A suggested form is as follows: "I am willing to be examined by Doctor So-and-so," signed and dated. The passages for the writing test should first be read over at normal speaking rate and then repeated slowly at dictation speed, words being spelt where necessary. The patient should be allowed to sit down at a table in good light, and be provided with lined paper and pen or pencil as he wishes. The handwriting by itself, of course, is of little value; some subjects will be illiterate, and others have only a poor mastery of penmanship; moreover, most people have intrinsic peculiarities of pattern and style. A fair assessment can be made only by comparing with a sample written on another occasion. However, gross defects are often seen, in spelling especially and in spite of it having been dictated slowly. Syllables, letters, and strokes are often repeated. The letters N, M, and W cause particular difficulty, excess up-and-down strokes often being made. Sometimes the handwriting shows gross incoordination.

Memory Test.—This is one of the best tests of mental alertness and accurate cerebration. Some objection was made to it by the Scottish judges, but fairly conducted it is valuable and a good response is in the patient's favour. It is wrong to ask leading questions such as, "How much drink have you had?" The accused should be asked if he is willing to give an account of himself during the preceding six hours, to test his memory. If he refuses, the test should not be proceeded with any further. Nevertheless, in spite of warning, some individuals make most damning admissions about the amount of drink they have consumed. In general there is an early account of great accuracy; as the time of maximum intoxication approaches the account becomes vague, devoid of detail, and sometimes there is a complete hiatus; often after an accident there is renewed alertness of memory. Other useful tests of mental awareness are to ask the subject to name the date—this often causes difficulty, especially in the early hours of the morning—and to give an estimate of the time. Sometimes the replies show great errors.

Specimens for Estimation of Alcohol Content

The patient should be asked if he will provide a specimen of urine for examination. Sometimes this has been done before the doctor's arrival. A few people object—and they are entitled to do so—but most defendants are only too glad to oblige. If he agrees, the patient is taken to a lavatory and is offered a clean vessel in which to pass water. The specimen is measured and divided into three equal parts, which are then placed separately in three clean containers. One is for the use of the patient and his own doctor, one for the police, who may send the specimen for analysis, and one is retained for the court in case they require it. All are sealed identically in the presence of the medical and other parties concerned. Routine tests are conducted later. Specimens may also be sent to a laboratory for the determination of the alcohol content.* A small specimen of blood is some-

*For further details see *The Recognition of Intoxication*, 1954, B.M.A., London. Price 1s.

times taken for the same purpose, but this procedure is not free from risks, particularly if the patient is not fully co-operative. Naturally, the syringe must contain no trace of alcohol.

Final Opinion

The complete clinical examination usually takes about forty-five minutes. On this examination, and on this examination alone, the doctor must form his opinion, although this may be amended as a result of a second examination later or a report on the laboratory findings. The examiner should never question the police about the circumstances of the arrest, as such information may unconsciously bias him. The opinion is formed on the general conclusion to be drawn from a large number of observations and responses to tests. *There is no clinical test, taken by itself, which is diagnostic of incapacity to control a car as a result of the effects of alcohol*, but a very fair opinion should be formed from the examination described. In borderline cases the benefit of any doubt should be in favour of the defendant. The only tests which are certain proof of consumption of drink are the smell in the breath, which is no guide to the amount, and the finding of alcohol in the blood or urine.

After the examination, the "police" doctor will be asked to make an entry in the occurrence book. It should be in simple form—for example, "I have examined John Jones at the Borough Police Office at 1 a.m. on Monday, April 9, 1956, and I have formed the opinion that he is (is not) under the influence of drink (or drugs) to such an extent as to be incapable of exercising proper control of a motor vehicle," and signed. He should also communicate his opinion to the accused. The patient's own doctor may record the fact that he has attended and examined him but should on no account commit his opinion to writing at this time; nor should he disclose his opinion verbally to the police unless the accused insists on it. He should tell the police that he will advise the accused's solicitor of his opinion at a later date. If the accused asks for his opinion, this should be given him privately.

In court the doctor should describe all his findings and the responses to tests, reading from notes *made at the time of examination*, specimens of handwriting, etc., being preserved as exhibits, together with the actual reading matter from which a test was made. It is for the court alone, when acquainted with all the facts, including the medical evidence, to decide whether the accused person has committed an offence.

Next article on Emergencies in General Practice.—
"Common Dislocations," by Mr. L. W. Plewes.

CAUSES OF ROAD ACCIDENTS

ERGONOMICS RESEARCH SOCIETY'S SYMPOSIUM AT BRISTOL

[FROM A SPECIAL CORRESPONDENT]

From April 16 to 19 the Ergonomics Research Society met at Bristol University to discuss the human element in road transport. They considered the role of such factors as dazzle, car design, alcohol, and fatigue in causing or aggravating traffic accidents, and they were able to indicate some of the ways whereby this mounting toll of life and limb could be reduced. The meeting was international in character, and results of research from the United States and the Continent were reported as well as those from Britain. Mr. P. G. MASEFIELD, managing director of Bristol Aircraft Ltd., opened the symposium in the presence of Sir PHILIP MORRIS, Vice-Chancellor of the University.

In 1954, said Mr. G. GRIME (Road Research Laboratory), the first speaker, the transport of goods and passengers by road in Britain was estimated to have cost nearly £2,000m.,

about a tenth of the country's total expenditure. Road accidents that year cost over £75m., and resulted in about 5,000 deaths and a quarter of a million other casualties. There could be no doubt, he declared, in the face of such figures, that the business of road transport would repay a very large research effort to improve its efficiency and make it safer. Indeed, much research on the prevention of road accidents is already in hand, as was evident from the papers read at the symposium, some of which are summarized here.

Dazzle and Glare

Dazzle and glare were discussed in relation to both street-lighting and headlamps. It is known that the use of "cut-off" lighting—lighting with a restricted beam—very largely eliminates glare in street lighting, but at the expense of reducing visibility between the illuminated areas of road. The nature of the light also matters. Dr. G. J. FORTUIN (Medical Department of Philips' Works, Holland) reported experiments which had shown that the "glare order" of five types of light was: yellow, sodium, incandescent, fluorescent, and mercury, yellow having the highest glare—twice as much as the mercury lamp, the lowest on the scale. With glare, what is important from a traffic point of view is the reduction of visibility when the eye is turned towards a strong source of light. In this context Dr. B. ROSENGREN (University of Gothenburg, Sweden) distinguished two types of glare: the first due to an adaptation of the retina to a higher value of luminance than the surroundings (here there are photochemical changes in the retina which manifest themselves as after-images, for example); and the second due to dispersal of light from a broad beam around the small particles in the refracting media of the eye. A driver instinctively avoided looking into the headlights of an approaching vehicle. The obscuring of detail when he did this was due to the latter form of glare. This was what had to be overcome, since it was from this type of glare that accidents would result, not from "discomfort" glare. The nearest possibility to a solution, Dr. Rosengren suggested, might be the use of a sharply demarcated light cone and, perhaps, a single headlight. In France, said Dr. A. Y. WISNER (Régie Nationale de l'Usine Renault), experiments had been carried out with two complementary sources of light, yellow and blue, mixing on the road. For the same illumination on the road it had been found that dazzle was greater if both headlights were white than with a blue axial headlight and a yellow lateral one. At present, however, coloured headlights could be obtained only by the filtering of white lights, and this involved a considerable energy wastage, which made the method uneconomical.

In general in England the aim has been to get more light on the road while preventing it from reaching the driver's eyes. To this end, experiments with cut-off beams are being carried out. The tendency on the Continent is to minimize discomfort; the Anglo-American aim is to have as large a range of vision as possible. It requires a beam of 120 ft. (36.6 m.) in order to see to swerve out 4 ft. (1.2 m.) when travelling at 40 m.p.h. (64 km.p.h.). Surveys of the dazzle experienced at night, said Mr. R. L. MOORE (Road Research Laboratory), had shown that it was due in the main to badly aimed headlamps, and in some cases to deteriorated lamps. Figures for glare in England were worse than those in America: in Washington, where there was compulsory inspection of the headlamps, 2% had a glare above 2,000 candles; in Maryland, where there was no compulsory inspection, the figure was 6%; while in England, where lights were on the whole badly maintained, it was 20%.

The use of polarized light was also discussed. If windcreens and headlights are polarized there is no glare. However, the difficulties of applying this method universally are too great. The amount of light absorbed by the polarized screens, and the difficulty of dealing with motor-cyclists and pedestrians, render it impracticable at the moment. Sharp demarcation of the headlight beam seems to be the most easily realized safeguard. In general, the feeling was against the use of tinted glasses or windcreens—indeed, of anything