RESPONSIBILITY OF THE GENERAL PRACTITIONER IN MARITAL PREPARATION AND PROBLEMS

BY

G. B. CARRUTHERS, M.D.

Marriage Counsellor and Medical Officer, Family Planning Association

The serious increase in the breakdown of the marriage relationship has led to the development of the concept of marriage guidance. There are two main objectives: (1) through pre-marital advice, to give couples a clear understanding of the various relationships within marriage, and to help to avoid some of the difficulties which may arise; and (2) if difficulties should occur, to help the partners towards achieving a solution.

The general practitioner is often approached for advice, either before marriage or, later, when difficulties have arisen. He may also be in a position to realize that marital discord is the basic cause of the numerous symptoms of preoccupation with minor ailments which many of his neurotic patients show. The time he can devote to assisting them is often restricted, but by his initial attitude and advice, and in appropriate cases by the suitable reference of a case, he can do much to give the couple a chance of solving their difficulties.

Unfortunately, young couples often have difficulty in finding a sympathetic and understanding adviser to whom they can turn for guidance. The Church has lost much of its influence as a counsellor, and parental intervention in these problems is now unfashionable and is looked upon as interference. The young couple could undoubtedly often gain much from the experience of their elders, who would themselves benefit from the realization that they still have an important part to play after their children have married.

As a result of this parental indifference the young couple often have nowhere to turn, and a sympathetic and understanding general practitioner can render great assistance.

Preparation for Marriage

The importance of adequate pre-marital advice cannot be stressed too strongly. The common problems which the couple bring to the doctor at this time concern their physical relationships; the spacing of their families; and, possibly, eugenic problems. Often they have been confused and disconcerted by reading numerous well-meaning but often disturbing books. They may have guilt or anxieties resulting from previous relationships. It is common experience, from breakdowns in marriage seen at Marriage Guidance Council centres, that much subsequent unhappiness could be prevented at this stage.

During the interviews it will sometimes be possible to detect equally important difficulties not appreciated by the couple themselves. Such facts as immaturity or irresponsibility in one or both partners may become apparent. These factors are important, because they affect the general marriage relationship or the physical relationship. If possible, an effort should be made to help the couple to realize their deficiency and, consequently, their unpreparedness for marriage. Individual or group discussions may prove valuable in these circumstances. The scope of the pre-marital consultation cannot be rigidly defined, and must be left to the individual concerned. Similarly, the extent of the need for pre-marital advice is not clearly laid down, and must be limited to the resources available.

The British Medical Association acknowledges that pre-marital advice may reasonably be given if there is any anxiety about general health, sex functions, or heredity. In fact, there are many other cases in which such advice would be of great value.

The order in which the interviews are handled is a matter of choice. I prefer to see the couple together initially for a few minutes to assess their attitude to one another, and then to talk to each individually.

A routine medical examination of each partner would seem advisable. In the woman this should include examination of the external genitalia to discover if there is any condition that might lead to difficulties in intercourse. The reassurance that she is normal and need not anticipate any difficulty is in itself of great value. Pelvic examination is advisable if this is possible. A routine chest x-ray examination, and in some cases a Wassermann reaction or rhesus grouping, may be indicated. If the wife should be rhesus-negative, and the husband rhesus-positive, both should be reassured regarding the remoteness of the chances of trouble arising.

Planning for a Family

Nowadays, most couples will be anxious to discuss the problem of family planning. The advisability of spacing their family must be decided by the couple themselves. Often this is necessary because of their living accommodation and financial position, apart from the realization that a properly spaced family is less likely to impair the wife's health.

An early child which necessitates the couple having to leave their accommodation or the husband to give up his training for a career is hardly a good basis for a subsequent happy marriage. Under present social conditions it must be accepted that many couples will wish to adopt some method of family spacing.

It is as well to point out, especially to women in their middle twenties or older, that the first baby should not be postponed for too long. The financial considerations are often exaggerated, because the couple do not fully appreciate that the pleasure of an infant more than compensates for the sacrifice which may be entailed. The woman is often unwilling to give up her work at first, but it is remarkable how delighted most women are when they actually start raising a family. One of the disadvantages of birth control is that it occasionally tends to be continued too long after the need has passed. It is important to remember that an early child is an important force in uniting the family. (I am not, of course, advocating having a child to cement a marriage that is already crumbling.)

The selection of the method of contraception is probably a difficult one. It is sometimes advisable for the couple to confine themselves initially to the use of a sheath and lubricant, and subsequently for the wife to try out a cap. Some couples will have decided that the wife should be fitted with a cap before marriage, in which case this can be done. It should, however, be remembered that this requires breaking the hymen in the virgin if internal protection has not been previously used, and occasionally frigidity in marriage can be attributed to the distaste aroused from the fitting and use of a cap as an introduction to the physical side of marriage. This is undoubtedly a danger in a timid or sensitive girl.

It is probably unwise to stress any possible difficulties at this early stage, and it may be best to leave this until they have occurred. In fact, difficulties are uncommon, and above all the couple both need reassuring that they are normal and that the physical side of marriage is less difficult than the books would make it appear.

It is important to stress that they should return a month or so after the marriage if they wish to. This is a suitable time to fit the wife with a cap if she so desires, and to check up and make sure that no physical difficulties have arisen. Any premature ejaculation in the husband should be treated at once. Usually, amylobarbitone sodium, 1 gr.

(65 mg.), at night and practical advice on positioning and technique will correct this. The disappointment and frustration which follow if this condition is left untreated can lead to serious consequences. Frequently middle-aged couples are seen who have had their whole marriage happiness marred by this one factor. Neither may in fact be aware of it. Sometimes the wife will not be obtaining any great satisfaction from intercourse, but at this early stage there is no cause for alarm. She should be advised to consult the doctor again if the condition does not correct itself over the next few months. Deep underlying psychological disorders requiring treatment are rare, and in any case treatment is difficult.

It is important to stress how welcome the couple are to return at any time when they may have problems they wish to discuss. It is valuable to know that the Marriage Guidance Council and Family Planning Association run pre-marital advice services, either for individual couples or through group discussions.

Marital Difficulties

The doctor may often be the first to realize that the marriage is not proving successful. The parties may approach him directly on this account, or, more often, their unhappiness will manifest itself through over-anxiety or the persistence of minor physical symptoms. Inquiry may elicit the fact that the underlying cause of their "neurosis" is marital disharmony. Too often this is accepted as the diagnosis, whilst the treatment remains symptomatic and is not directed to the underlying disorder. Intervention at this level is likely to involve considerable expenditure of time, but much help can often be given by discussion with one or both parties.

Sometimes the problem turns out to be a definite one which can be tackled. Such problems include physical difficulties such as premature ejaculation, frigidity, or impotence, which can be helped by reference to the appropriate consultant, or there may be a housing or social problem which can be helped by contact with the welfare or social services available for this purpose. In most cases the difficulty proves to be a less definable one, such as incompatibility of temperament, unfaithfulness, or mistreatment by one or other party. Sometimes the incompatibilities are relatively minor and can be helped by experienced guidance from the friendly practitioner. Often, however, it will be necessary to see one or both parties on many occasions and the practitioner will feel he cannot afford sufficient time. It may also seem inadvisable to intervene because of the professional relationship, and the need to avoid being involved in a situation that may eventually lead to legal proceedings. In all these cases it would seem advisable to refer them to an appropriate body which handles such problems—namely, the Marriage Guidance Council.

Training the Marriage Counsellor

The Council has about 80 centres, situated all over England. These are under the auspices of a National Marriage Guidance Council, which is responsible for the selection and training of all counsellors. The counsellors are carefully chosen after attending residential selection conferences, when they are interviewed and tested for suitability by a team which includes a psychiatrist. They are selected from middle-aged married men and women, largely professional people and housewives whose personal and general background would seem to make them suitable for such work. They must then attend an extensive training programme spread over four or more week-ends at residential centres.

The training is designed to instruct the person in the technique of counselling and the proper handling of the situations that may arise. The marriage counsellor's object is not to dispense advice or presume to tell people what to do. It is to befriend them and help them to work their way through their difficulties to their own most acceptable solutions. The training includes lectures on the common legal

difficulties that may be involved, and lectures by psychiatrists to help the prospective counsellor to pick out any mentally disturbed or unbalanced cases they may meet, so that these may be referred to an appropriate specialist.

Behind the counsellors in each Marriage Guidance Council centre are a team of medical, legal, and spiritual consultants, and social workers, for consultation in their respective fields. Some counsellors are themselves doctors, and primarily sexual or medical problems are naturally seen by them. The medical specialists available cover psychiatric and sexual problems.

Co-operation with the general practitioner is close in many areas, in others could be much closer. The council is only too eager to have cases referred to it by the general practitioner. There is no danger that counsellors will interfere with his treatment. They should be regarded as medical auxiliaries who can give the large amount of time needed to listen to and help sort out the marital problems which are leading to a broken home, to problem children, or to physical symptoms.

The co-ordinated efforts of the general practitioner and the marriage counsellor—one helping to preserve the individual health, the other helping to retain the family relationship and thus the family "health"—can together prove of immense value to the happiness of the home.

I wish to thank Mr. A. J. Brayshaw, General Secretary of the National Marriage Guidance Council, for his valuable advice and comments in connexion with the preparation of this article.

CASE NOTES

THEIR VALUE FOR RESEARCH

Much of the medical literature of the past is based on material extracted from medical records. It was appropriate, therefore, that the Medical Section of the Royal Statistical Society should devote an evening to considering the real value and reliability of the routine case record. This they did on May 29 at a meeting held at the Westminster Hospital Medical School.

Causes of Error

Dr. A. C. DORNHORST (St. Thomas's Hospital) opened the discussion with an account of the type of information that could be found in most routine medical notes. Non-professional details included those of a personal nature, and even some of these apparently simple facts were often recorded in a misleading fashion. One error he had encountered was that where the age was recorded on every page of the patient's notes it often remained the same despite the passage of many years. Professional information included both clinical and technical details. The taking of a careful clinical history was generally accepted as the foundation of accurate diagnosis, but in fact there was often great variation if the history was taken from the same patient by different observers. Among the reasons for this were the asking of questions with an impossible amount of detail, a poor memory on the part of the patient, and the difficulty he often had in grasping the exact question asked. All too often the patient was thinking in completely different terms. Leading questions were a well-recognized pitfall, and yet often they could be avoided only if the examiner had unlimited time and patience, and preferably a forensic training. If leading questions were asked the whole history-taking became a conspiracy, and the conclusions reached depended largely upon the personality of the individuals concerned.

The clinical examination was often recorded partly in terms of an "exact" nature—measurements which were not in fact reliable. Examples of this were the position of the apex beat and the degree of enlargement of the liver. Recent publications had shown that there was considerable variation if the same fact was recorded by different individuals. Progress notes were usually distinguished by their absence, par-