insipidus itself the action of adrenal cortical hormones uninhibited by A.D.H. may be contributory to the observed polyuria. Both Sirek and Best (1952) and Swingle *et al.* (1953) have produced experimental evidence in support of this. They were able to induce diabetes insipidus by injecting dogs with large doses of cortisone. "Pituitrin" was effective in controlling the polyuria and polydipsia only when administered in large amounts. The latter authors concluded that the increased urine volume was brought about by (1) thirst due to sodium retention, (2) inhibition of tubular reabsorption of water, and (3) probably by reduction of the amount of A.D.H. released by the posterior pituitary.

Clinical Studies

Garrod and Burston (1952) observed the effects of cortisone on water diuresis in patients with either Addison's disease or panhypopituitarism. The diuretic response was defective in all patients initially, but was restored to normal after cortisone was given. And, although cortisone raises the glomerular filtration rate and renal plasma flow, this is only a partial factor. The main role of cortisone is to enable the renal tubules to regulate the reabsorption of water. Another important and interesting feature was that cortisone induced a transient diabetes insipidus in one of their patients with impaired anterior pituitary and neurohypophysial function due to a suprasellar cyst. This is almost analogous to the state of affairs that existed in our case, except that it is uncertain in our patient what part the focal necrosis in the anterior pituitary played.

Prunty (1953), in a study of the effects of cortisone in patients with Addison's disease, came to the conclusion that, while sodium and chloride retention occurs, this may not persist as long as the hormone is administered. Under certain conditions diuresis may occur with increased excretion of these ions.

There is thus some evidence from previous studies to support the hypothesis that an excess of adrenal cortical hormones (in our case exogenous cortisone), acting without or with only negligible A.D.H., can convert temporarily a mild diabetes insipidus to a severe form.

We wish to emphasize one further point. Cortisone actually made this leukaemia worse. This is in accord with the recent observations of Fessas *et al.* (1954). They found that cortisone and corticotrophin were of benefit in the acute lymphoblastic leukaemia, whereas they were of no use, or even accelerated the leukaemic process, in cases of acute myeloblastic or monoblastic leukaemia.

Summary

A case of monocytic (paramyeloblastic) leukaemia with diabetes insipidus is reported.

Necropsy changes in the pituitary and other organs are described.

The effect of cortisone in increasing the polyuria is observed and the possible modes of action are discussed.

We are grateful to Dr. P. R. Evans for his helpful advice, to Drs. R. L. Waterfield and J. T. Cecil for their reports on the marrow and blood films, to Dr. A. J. Smith for the necropsy report, and to Dr. S. J. De Navasquez for the histological report. We are indebted to Mr. R. Nichols for Fig. B and Guy's dental photographic department for Fig. A.

REFERENCES

- Castaigne, P., and Hubault, A. (1953). *Rev. prat.*, 3, 1871.
 Cates, J. E. (1953). *J. Endocr.*, 9, Suppl. XII.
 — and Garrod, O. (1951). *Clin. Sci.*, 10, 145.
 Chalmers, T. M., and Lewis, A. A. G. (1951). *Ibid.*, 10, 127.
 Fessas, P., Wintrobe, M. M., Thompson, R. B., and Cartwright, G. E. (1954). *A.M.A. Arch. intern. Med.*, 94, 384.
 Garrod, O., and Burston, R. A. (1952). *Clin. Sci.*, 11, 113.
 Gaunt, R., Birnie, J. H., and Eversole, W. J. (1949). *Physiol. Rev.*, 29, 261.
 Hickey, R. C., and Hare, K. (1944). *J. clin. Invest.*, 23, 768.
 Jones, G. M. (1944). *Arch. intern. Med.*, 74, 81.
 Kugelmeier, L. M. (1937). *Z. klin. Med.*, 132, 521.
 Lenk, R. (1911). Quoted by Warkany and Mitchell (1939).
 Prunty, F. T. G. (1953). *British Medical Journal*, 1, 852.
 Sheldon, J. H. (1927). *Lancet*, 1, 489.
 Sirek, O. V., and Best, C. H. (1952). *Proc. Soc. exp. Biol. (N.Y.)*, 80, 594.
 Swingle, W. W., Fedor, E. J., Ben, M., Maxwell, R., Baker, C., and Barlow, G. (1953). *Ibid.*, 82, 571.
 Warkany, J., and Mitchell, A. G. (1939). *Amer. J. Dis. Child.*, 57, 603.

POST-MENOPAUSAL OESTROGEN PRODUCTION

BY

R. A. STRUTHERS, M.D., M.R.C.O.G.

(From the Department of Obstetrics and Gynaecology,
St. Bartholomew's Hospital, London)

[WITH SPECIAL PLATE]

It was formerly believed that oestrogen activity ceased with the menopause. Later, that opinion was modified and the view was held that, in some women, some oestrogenic activity continued for a time after the menopause. Little has been done to change this broad generalization into something more precise. The impression conveyed by the literature is that post-menopausal oestrogenic activity, while not a rarity, is not common, but that the true incidence of post-menopausal oestrogen production is not known.

It is accepted that the ovaries are the chief source of oestrogen, and, in particular, the granulosa cells of the follicle, but as the Graafian follicles cease to mature at the menopause one has to look further afield for the source of post-menopausal oestrogen.

It was decided, therefore, to carry out an investigation in order to (1) determine the number of women in whom post-menopausal oestrogenic activity was present; (2) attempt to make a quantitative estimation of that oestrogen; and (3) determine its source. It was with these aims that an attempt was made to provide an answer to three problems. Although the direct purpose was to find out whether there was any significant amount of post-menopausal oestrogen activity in women, and, if so, whether it was ovarian in origin or not, it was thought possible that it might provide some information regarding the importance of oestrogen in malignant disease.

Method

It was realized that to carry out this investigation several hundred patients would have to be examined if the results were to be valid. The method of estimating the oestrogen would have to be simple, quick, inexpensive, and of reasonable accuracy. Methods employing colorimetry and fluorimetry were thought to be impractical.

The method of oestrogen determination of Brown and Bradbury (1949), who suggested the use of the human vaginal smear in the assay of oestrogens, has much to commend it. Post-menopausal women with atrophic vaginal smears were given oestrogen, and the smear results were evaluated into four grades. It seemed that this method had advantages, and possible inaccuracy through using animals is eliminated. Expense was negligible, the method was simple, and large numbers of estimations could be made in a day. The main source of error lay in the cytological interpretation, but with experience this could become minimal. The main adverse criticism was that it was only broadly quantitative. The results were divided into four grades, which were relative and not absolute values: the method could not be more precise. For assessing post-menopausal (natural and artificial) oestrogen activity, a procedure similar to this, in which the individuals concerned were tested directly, is therefore appropriate; the lack of precision for which this method can be criticized is

not of importance. Shorr (1940) believed that such a method of bio-assay on the individual patient was better than other methods of bio-assay, and he supported the criticism by Marrian, who said that the usual methods of assay estimated breakdown products rather than functioning oestrogen. Further, he believed that the substitution of "patient subject bio-assay" for animal assay was more reliable.

Aeppli and Rosenmund (1951) suggested that the smear method had definite practical advantages, and tested its accuracy against the bio-assay method of Allen and Doisy; they found that the two methods gave results that were remarkably similar and that the former could therefore be regarded as accurate enough. For these reasons the smear method was chosen in the present investigation of post-menopausal oestrogen production in women.

The smears for cytological examination were obtained by the use of a spatula, which was stroked firmly down the upper third of the anterior wall of the vagina. The smear was then made on a microscope slide and at once placed in a fixing solution of equal parts of ether and absolute alcohol. The stain used was described by Papanicolaou *et al.* (1948), and is a modification of his original stain. The smear was left in Harris's haematoxylin for eight minutes instead of five, and in E.A. 36 for two minutes instead of one and a half. The purpose of this was to get a more heavily stained smear. The material in this series was scanty and showed up better with heavy staining. With thicker smears—those usually obtained in pre-menopausal women—such a method would lead to overstaining.

Method of Grading

Each smear was graded according to the cytological characteristics. Four grades were used: A, B, C, and D. Grade A was fully oestrogenic (the same as a pre-menopausal woman) and grade D completely non-oestrogenic and atrophic. The majority lay between these extremes and were graded B or C according to the proportion of squames, navicular and parabasal cells, their nuclei, and their staining qualities.

Grade A.—The smear is made up of fully developed squames, large thin cells with curled-up or crenated margins, mostly acidophilic, and with pyknotic nuclei (Special Plate, Figs. 1 and 2).

Grade B.—Some fully developed squames are present, but the majority are smaller navicular cells with larger nuclei. There is a greater proportion of basophilic stained cells (Plate, Figs. 3 and 4).

Grade C.—There are no fully developed squames. The majority of cells are navicular and there are some parabasal cells. The latter are the smallest cells of the exfoliated vaginal epithelium; they are basophilic, round, and have relatively large pale nuclei (Plate, Figs. 5 and 6).

Grade D.—The smear is composed almost entirely of parabasal cells with only an occasional navicular cell (Plate, Figs. 7 and 8).

Several considerations have been omitted from this description of the four grades. No mention has been made of the distribution of the cells, whether discrete or in clumps, or the presence of mucus, leucocytes, and erythrocytes. This was done so that the grading should be straightforward and simple. The characteristics described under each grade were quite sufficient to enable each smear to be allotted to its appropriate grade.

It is accepted that oestrogen production fluctuates after the menopause; it therefore seemed quite possible that one test of each patient might be fallacious. Forty unselected patients had smears taken on two occasions between four and eight weeks apart, and their levels on these occasions were compared. Remarkably little variation was found when assessment was made by this method, the error being 2.6%.

Results after the Natural Menopause

Vaginal smears were taken from 389 patients. All were asked if they had been taking oestrogen or androgen in any form and by any route; 36 patients who were known to have taken either within a month of examination were excluded from the investigation.

Although all the remaining 353 patients were post-menopausal, only 110 had had a natural menopause. The remainder had had the menopause induced artificially either by oophorectomy or by irradiation with deep x rays or radium. The 110 patients who had had a natural menopause were unselected, apart from ensuring that the menopause had been natural and that no oestrogen had been taken. They were either in medical and surgical wards or attending gynaecological out-patient clinics. These latter patients were not attending because of symptoms related to oestrogenic activity or the lack of it. They were therefore regarded as a fair representation of normal post-menopausal women. The oestrogen production of this group is shown in Table I.

TABLE I.—Number with Oestrogen Activity After Natural Menopause, Oophorectomy, and Radiotherapy

	Grade A	Grade B	Grade C	Grade D	Total
Natural menopause ..	6 (5.45%)	58 (52.7%)	29 (26.4%)	17 (15.45%)	110
Oophorectomy ..	2 (1.5%)	55 (42%)	40 (30.5%)	34 (26%)	131
Deep x rays ..	2 (2.7%)	43 (58.1%)	17 (23%)	12 (16.2%)	74
Radium ..	0	22 (57.9%)	8 (21.05%)	8 (21.05%)	38

It will be seen that 5.45% showed as much oestrogenic activity as any pre-menopausal smear, 52.7% showed somewhat less than is seen on a pre-menopausal smear, 26.4% showed a little oestrogen, and in 15.45% there was no detectable oestrogen by this method. Therefore, 64 (58%) of the women who have had a natural menopause show a considerable amount of oestrogenic activity. This figure is at variance with the general opinion and significantly higher than the figures quoted by others. Even so, 29 (26.4%) who were thought to show the presence of a little oestrogen were not included. It seems, then, that it is the exception for a woman to secrete no oestrogen at all after the menopause. This in itself is an interesting conclusion. It immediately raises the question of the source of the post-menopausal oestrogen.

Results of Oophorectomy

It has been shown that the ovaries, and the granulosa cells of the ovaries in particular, are the main source of oestrogen. These granulosa cells surround the maturing ovum of the Graafian follicle. At the menopause ovulation ceases, but partial ripening of the follicles continues for a few months and possibly years (Allen, 1941). Ingraham (1941), using a special reticulum stain, has demonstrated theca cells in the post-menopausal ovary. It is probable, therefore, that cells in the ovary, other than the granulosa cells, may produce oestrogen after the menopause.

Hadfield (1954) has pointed out that the cells of the ovarian stroma make a very substantial contribution to the formation of the early Graafian follicle. They are present in the greatest number during the peak period of production of oestrogen. They might well be a source of oestrogen in themselves. He adds that certain ovarian stroma cells become more active after the menopause. There has been little histological investigation into the post-menopausal ovarian stroma. Shaw and Dastur (1949) described some

particular cells (Shaw-Dastur cells), but were unable to determine their function, associating them with cancer and believing them to have a hormonal action.

It appears, therefore, that the ovary may be a source of post-menopausal oestrogen. To find out if this is so, vaginal smears were taken from women who had had their ovaries removed and from others who had had deep x rays or a radium irradiation of their ovaries. By these methods any ovarian source of oestrogen would be destroyed, provided that the irradiation had been adequate. If the ovaries were the source of post-menopausal oestrogen no evidence of oestrogenic activity would be found from the vaginal smears. A total of 131 women were examined who had had their ovaries removed. Table I shows the oestrogen production in this group.

Grouping grade A and B results together, it can be said that 57 out of 131 (43.5%) women showed considerable oestrogenic activity after oophorectomy. This is only 14.5% less than the oestrogenic activity found after a natural menopause (58%).

If the proportion of women showing substantial oestrogenic activity is only 14.5% less in women without ovaries than it is in women with ovaries, the output of oestrogen by the ovary after the menopause can be only very little at the most. Experimental error is unlikely to account for the whole of this difference, but it could mean that the amount of oestrogen produced from the ovaries in a post-menopausal woman is insignificant.

Results after Radiotherapy

It was next intended to show the figures for post-menopausal oestrogen production in women after radium and x-ray irradiation to the ovaries. If the figures were comparable with those from oophorectomy, then the three artificial groups would be considered together as one for purposes of comparison with the natural menopause group.

The women who had had radium or deep x-ray therapy were with only few exceptions of one symptomatic group. They had had menorrhagia, polymenorrhoea, or irregular bleeding at the time of the menopause. After diagnostic curettage to exclude malignant disease the haemorrhage was controlled by the induction of the menopause by either of the two methods. There was no particular reason in individual cases for choosing x rays rather than radium to produce an artificial menopause. All the women were treated at the same hospital and the same standard method of treatment was used. None had been irradiated less than six months before having a vaginal smear taken.

The radium dose was 50 mg. of intrauterine radium for forty-eight hours—that is 2,400 mg.-hours. The effect of the irradiation on the ovaries is directly related to their distance from the uterine cavity. This was investigated by measuring the distance on patients at abdominal operations with the pelvic organs in their normal position. The average distance was 5 cm., but there can be some variation, and therefore also of intensity of irradiation. With 2,400 mg.-hours, the dose to the ovaries at 5 cm. is approximately 720 r. This dose is much greater than that employed by many. A dose of 200 r at the ovary suppresses menstruation in almost all women. It can be said, therefore, that it is most likely that the patients receiving radium treatment to the ovaries in this series had had their own ovarian function destroyed. The patients who had deep x rays to their ovaries had a dose of 900 r. So both the radium and the x-ray groups can be safely assumed to have no ovarian oestrogen. Table I shows the oestrogen production in these two groups.

The results shown in Table I are compared in the graph. Statistical analysis shows that there is no significant difference between the four groups—that is, between the natural menopause and the artificial, or between the three individual artificial menopause groups.

Table II compares the natural group with the three artificial menopause groups combined. The graph and Table

I make it clear that the two main groups are comparable. Therefore, in appraising the post-menopausal oestrogen production, all the 353 women examined can be considered as one group, and this is done in Table III. Of the women

TABLE II.—Comparison of Natural and Artificial Menopause Groups

	Grade A	Grade B	Grade C	Grade D	Total
Natural	6 (5.45%)	58 (52.7%)	29 (26.4%)	17 (15.45%)	110
Artificial	4 (1.7%)	120 (49.3%)	65 (26.8%)	54 (22.2%)	243

TABLE III.—Grading of Whole Series

Grade A	Grade B	Grade C	Grade D	Total
10 (2.8%)	178 (50.4%)	94 (26.6%)	71 (20.1%)	353

in this series, 53.2% showed considerable oestrogenic activity, 26.6% showed the presence of a little oestrogen, and only 20.1% showed no oestrogen to be present. At the maximum, then, nearly 80% of women exhibit a varying amount of post-menopausal oestrogen production. This is a remarkably high figure, and there seems little reason to doubt its accuracy.

Results after Adrenalectomy

It having been established that there is a high incidence of oestrogen production after the menopause and that this oestrogen most probably does not come from the ovary, it is necessary to look for another possible source. There is some evidence that the adrenal is the source of oestrogen production in such patients. Oestrogen was extracted from the adrenal glands of rabbits in 1930 by Engelhart. The substance which he extracted caused uterine hypertrophy, and an extract of the adrenal taken during pregnancy had a much more pronounced effect. In 1934 Frank reported two cases of carcinoma of the adrenal gland with a very high oestrogen content in the urine, and Simpson and Joll (1938) reviewed five cases of feminization with adrenal tumours. In the field of experimental medicine, Fekete *et al.* (1941) removed the ovaries at birth from a group of mice of a cancer-prone strain. They noted nodular hyperplasia of the adrenals following this, the hyperplastic cells being lutein-like. The uterus, vagina, and breasts remained in a rudimentary state for seven months, but then developed as if oestrous cycles were occurring. It is, of course, dangerous to compare what happens in mice of a cancer-prone strain with any changes in women, but the apparent taking over of ovarian function by the adrenals in mice is of some significance when searching for the source of post-menopausal oestrogen in women.

The coelomic mesothelium is the common embryological origin of both the ovaries and the adrenals.

If removal of the ovaries does not suppress oestrogenic production and the adrenal is really the source of this, then removing the adrenals should result in eliminating the oestrogen altogether, for without both ovaries and adrenals there should be no oestrogenic activity. The advent of cortisone enabled Huggins and Bergenstal (1952) to do this, for without cortisone bilateral adrenalectomy always proved fatal. With the use of cortisone five of their first six cases survived. They carried out this work in women thought to have oestrogen-dependent cancer because the growth might be expected to regress if the source of oestrogen was removed. This did occur in two of the six cases mentioned.

To discover whether the adrenals were the source of oestrogen in post-menopausal women it was necessary to examine vaginal smears from women whose ovaries and adrenals had been removed. If all oestrogen disappeared from women who produced considerable quantities of oestrogen after oophorectomy and before adrenalectomy, then it could be concluded that the adrenals were an important source of post-menopausal oestrogen. Theoretically this was likely to be so, but of course little clinical material is available. Eight such cases were available for examination of vaginal smears, but one of these women died within twenty-four hours of adrenalectomy. In four of the remaining seven an estimate of oestrogen production was made before and after adrenalectomy, in three there was no change in the oestrogen level, and in the fourth there was a slight reduction; of the three whose level remained unchanged, two showed high and one low oestrogenic activity. Three women had smears taken after adrenalectomy only; two of these showed moderately high oestrogenic activity and one showed none. Although control smears taken before adrenalectomy are not available for these women, the fact that two show the presence of much oestrogen means that removing the adrenals has not removed the source of oestrogen.

If any conclusion at all can be drawn from this small series it is that the adrenals are not the main source of post-menopausal oestrogen, and that although they probably produce some oestrogen there is probably some other main source of production. This conclusion can be criticized on the grounds that an insufficient number of patients have been investigated, but had the smears shown a large reduction in oestrogenic activity many more patients would have had to be examined before any conclusions could have been reached. Even if one case had shown the same oestrogenic activity before and after adrenalectomy it could be said that the adrenal is not the whole source of post-menopausal oestrogen, and when six examined patients show continuing oestrogen production after adrenalectomy it is fair to assume that there must be another source of this oestrogen. It is inevitable that these patients belong to a highly selected group. They had extensive carcinoma of the breast with widespread metastases, and metastases were found in the adrenals of three. Each one was in the menstrual phase of life before oophorectomy was carried out. The interval in each case between the oophorectomy and adrenalectomy was between six months and two and a half years, and they were, of course, all receiving cortisone after adrenalectomy.

These patients form a small but abnormal group, for almost certainly they suffered from an abnormal hormonal metabolism. It was essential, therefore, to compare the oestrogen level before and after adrenalectomy, for if it remained unchanged, then the issue was reduced to its simplest terms and the abnormal factors were of no consequence. In such a case, any change of oestrogen level could be attributed only to the one altered factor produced by bilateral adrenalectomy. In view of the findings in these six cases it is probably justifiable to conclude that the adrenal gland is not an important source of post-menopausal oestrogen.

It is essential to investigate further the source of oestrogen production in these cases, and the next gland to be implicated is the pituitary. Oestrogen has been found in the hypophysis of cattle and horses (Callow and Parkes, 1936), and its

relationship to ovarian function is very close. It was interesting to note that one woman of 47 years who had had amenorrhoea for twelve years following childbirth due to hypopituitarism had no oestrogen production according to the smear method of estimation. Further investigation of the pituitary as the main source of post-menopausal oestrogen was not possible at this time. Hypophysectomy has since been carried out on a number of patients, but as yet the effects of this procedure have not been fully evaluated. Professor H. Olivecrona and his colleagues in Stockholm maintained life satisfactorily with hormonal replacements which included either oestrogen or androgen.

Oestrogen Production in Female Breast and Genital Tract Cancer

Some of the patients in this series had advanced carcinoma of the breast, and it is possible that cases of breast carcinoma might be dependent on the presence of oestrogen in the blood stream. In 1896, Beatson, an Edinburgh surgeon, first put into practice a theory born from an impelling desire to do something to alleviate the suffering of the many women sent to him with extensive inoperable fungating carcinoma of the breast. He carried out bilateral oophorectomy on three women with extensive carcinoma of the breast, and there was a remarkable regression of the growth in one case. From this beginning many similar cases have been dealt with in the same way, but it cannot be said that a definite conclusion has yet been reached.

Should carcinoma of the breast depend to any extent on an oestrogenic stimulus, it might be expected that women suffering extensively from this disease would exhibit more oestrogenic activity than normal women. Two such groups of cases were therefore compared.

The first series consisted of 57 women with extensive carcinoma of the breast who either were post-menopausal or had had their ovaries removed. These are compared in Table IV with the women who did not have breast cancer

TABLE IV

	Grade A	Grade B	Grade C	Grade D
Women with carcinoma of breast	2 (3.5%)	27 (47.4%)	16 (28.1%)	12 (21.0%)
Normal women	8 (3.3%)	132 (46.9%)	53 (28.6%)	41 (21.2%)

but were directly comparable in other respects because they were post-menopausal or had had their ovaries removed. The percentage of oestrogenic activity in the four grades of these two groups of patients are remarkably similar.

The oestrogen level is affected by age, and so the ages of the groups had to be compared. The average age of the normal women who had had either a natural menopause or had had a bilateral oophorectomy performed was 56.1 years at the time of taking vaginal smears. The average age of the breast-carcinoma patients when smears were taken was 55.8 years. The ages of the two groups can thus be regarded as comparable, the age ranges being similar.

The conclusion, therefore, that in this series there is no greater oestrogen production in patients with breast carcinoma after a natural or artificial menopause than in normal women after a natural or artificial menopause appears to be valid. This suggests that oestrogen may not be an important factor in "hormonal dependent" breast carcinoma.

Although it does not appear that oestrogen is an important factor in breast carcinoma, it may have some significance in the aetiology of cancer of the female genital tract. Amongst the smears that have been analysed for other purposes, 67 were taken from women with carcinoma of the uterus or ovary. The oestrogenic activity of these smears from cases with genital cancer is compared in Table V with that of smears from a similar group of women without known malignant disease. From the figures there appears to be rather less oestrogenic activity in the women with cancer of the genital tract than in other women. Age

TABLE V

	Grade A	Grade B	Grade C	Grade D
Genital cancer	1 (1.5%)	25 (37.3%)	25 (37.3%)	16 (23.9%)
No genital cancer	8 (3.4%)	132 (56.4%)	53 (22.6%)	4 (1.7%)

was not a source of bias, for the average age of the patients with genital cancer was 57.0 years and that of the other women 56.1 years.

Summary

An investigation has been made into the post-menopausal production of oestrogen on a series of 353 women.

The production of oestrogen was found to be high, 53% of women showing evidence of considerable oestrogenic activity and only 20% showing none at all.

The oestrogen production was not affected by the manner in which the menopause was achieved.

Neither the ovary nor the adrenal was found to be a significant source of oestrogen after the menopause.

Oestrogen did not appear to have an important part to play in either breast or female genital tract cancer.

Part of this work was undertaken during the tenure of a grant from the Yapp Research Fund. I wish to thank Mr. John Beattie and Dr. E. F. Scowen for their help and Mr. Bert Cambridge for his technical assistance.

REFERENCES

- Aeppli, H., and Rosenmund, H. (1951). *Gynaecologia (Basel)*, 131, 404.
 Allen, E. (1941). *J. Amer. med. Ass.*, 116, 405.
 Beatson, G. T. (1896). *Lancet*, 2, 104, 162.
 Brown, W. E., and Bradbury, J. T. (1949). *J. clin. Endocr.*, 9, 725.
 Callow, R. K., and Parkes, A. S. (1936). *J. Physiol. (Lond.)*, 87, 28P.
 Engelhart, E. (1930). *Klin. Wschr.*, 9, 2114.
 Fekete, E., Woolley, G., and Little, C. C. (1941). *J. exp. Med.*, 74, 1.
 Frank, R. T. (1934). *Proc. Soc. exp. Biol. (N.Y.)*, 31, 1204.
 Hadfield, G. (1954). *Ann. roy. Col. Surg. Engl.*, 14, 21.
 Huggins, C., and Bergenstal, D. M. (1952). *Cancer Res.*, 12, 134.
 Ingraham, C. (1941). *Amer. J. Obstet. Gynec.*, 42, 577.
 Papanicolaou, G. N. (1925). *Proc. Soc. exp. Biol. (N.Y.)*, 22, 436.
 — Traut, H. G., and Marchetti, A. A. (1948) *The Epithelia of Woman's Reproductive Organs*, p. 5. New York.
 Shaw, W., and Dastur, B. (1949). *British Medical Journal*, 2, 113.
 Shorr, E. (1940). *Bull. N.Y. Acad. Med.*, 16, 453.
 Simpson, S. L., and Joll, C. A. (1938). *Endocrinology*, 22, 595.

CEREBELLAR SOFTENING

A SURGICAL EMERGENCY

BY

BERNARD FAIRBURN, M.B., F.R.C.S., D.A.

AND

LESLIE C. OLIVER, M.B., F.R.C.S., F.I.C.S.

Consultant Neurosurgeons, Neurosurgical Centre, Oldchurch Hospital, Romford, Essex

Although cerebral haemorrhage is so common cerebellar haemorrhage is rare; Courville (1950) found only 117 examples in 1,484 necropsies of patients dying from spontaneous haemorrhage in the brain (less than 8%). Despite the rapidly fatal nature of cerebellar haemorrhage (Mitchell and Angrist, 1942) a few patients have been saved by timely surgery (Guillaume *et al.*, 1949; Dickmann and Zimmann, 1949).

Softening of the cerebellum is even rarer than haemorrhage. Thus Grinker and Bucy (1949) state that "because of the many anastomoses, softenings of the cerebellum are almost pathological curiosities"; and search of the literature has disclosed only three references to the condition (Germain and Morvan, 1938; Fattovich and Lenti, 1940; Fattovich, 1950). Hitherto, diagnosis has been made at necropsy and no well-defined syndrome has been established. No reference to

posterior fossa decompression having been carried out for this lesion has been discovered.

In this article, records are presented of three patients whose symptoms and signs indicated a space-occupying lesion, and evidence is given that softening and swelling of one cerebellar hemisphere—probably due to vascular occlusion—was the cause in each case.

Case 1

The patient was a man aged 40. On April 23, 1953, while at work he felt faint and vomited several times. He did not lose consciousness, but felt so ill that he had to be sent home to bed. He continued to vomit, and was therefore admitted to hospital on April 25, when he was drowsy and had signs of meningeal irritation. Lumbar puncture showed a raised pressure (reading not recorded). The cerebrospinal fluid contained 60 mg. of protein per 100 ml. His condition deteriorated during the next two days. He became comatose and developed Cheyne-Stokes respiration and transient Babinski responses. Another lumbar puncture revealed turbid fluid containing 260 white cells per c.mm. (89% polymorphs) and 80 mg. of protein per 100 ml. The fluid was sterile. He gradually regained consciousness and was transferred to us on May 1, nine days after the onset of the first symptoms. On examination he was found to be fully conscious but could not remember the first three days of his illness. His only complaint was of mild headache and pain in the neck. A complete clinical examination showed no abnormal physical signs apart from a little neck stiffness. Radiography of the skull and chest and examination of the blood showed nothing abnormal.

Ventriculography (May 19) was performed with air and ethyl iodophenylundecylate. The ventricular pressure was 60 mm. There was moderate dilatation of all the ventricles and also a small irregular filling defect in the fourth ventricle on the left side.

The posterior fossa was explored on the same day. Both cerebellar tonsils were herniated through the foramen magnum. The right tonsil descended further than the left, reaching the lower border of the axis. It was very shrunken, frayed, and yellowish, and was adherent to the medulla oblongata. The dura mater was adherent to the right cerebellar hemisphere, which was also very shrunken and soft. An incision into it revealed yellowish diffuent tissue. Retraction of the right cerebellar lobe and tonsil showed atheromatous occlusion of the peripheral part of the posterior inferior cerebellar artery. The left side of the cerebellum was of normal appearance and consistency, and there was no atheroma of the left posterior cerebellar artery. The cerebello-pontine angle was clear on both sides and a fine catheter was passed upward into the fourth ventricle through the foramen of Magendie without obstruction. The small fragments of cerebellum taken from both hemispheres showed no histological evidence of tumour. It was concluded that infarction of the right cerebellar hemisphere had occurred from atheromatous occlusion of the distal part of the posterior inferior cerebellar artery (occlusion of the proximal part would have produced the classical syndrome of infarction of the medulla oblongata). The patient made a rapid and complete recovery and when last examined on January 11, 1956, was symptom-free and showed no abnormal physical signs.

Case 2

A man aged 69 was admitted to hospital on May 29, 1954, having the day before had an attack of giddiness and sweating followed by frontal headache and repeated vomiting. During the next 24 hours the headache became worse and was also felt in the occipital region. The vomiting continued. He had had two similar attacks 12 and 9 months previously, when repeated vomiting with slight headache had occurred.

R. A. STRUTHERS: POST-MENOPAUSAL OESTROGEN PRODUCTION

FIG. 1.—Grade A. Showing uniformly fully developed squames. ($\times 120$.)

FIG. 2.—Grade A. Illustrating large size of individual cells and small pyknotic nuclei. ($\times 300$.)

FIG. 3.—Grade B. Compared with grade A, both the size of cells and that of their nuclei vary. ($\times 120$.)

FIG. 4.—Grade B. The nuclei of these cells are much larger than those of grade A. ($\times 300$.)

FIG. 5.—Grade C. Showing mixture of navicular and parabasal cells with no fully developed squames. ($\times 120$.)

FIG. 6.—Grade C. Apart from navicular and parabasal cells, one degenerate squame can be seen. ($\times 300$.)

FIG. 7.—Grade D. Although most cells are of small parabasal type with large nuclei, there are three poorly developed squames and one navicular cell. ($\times 120$.)

FIG. 8.—Grade D. All cells parabasal apart from one navicular. One small cell is almost fully occupied by the nucleus and must be regarded as a most primitive form of parabasal cell. ($\times 300$.)