EMERGENCIES IN GENERAL PRACTICE**TREATMENT OF SEVERE HAEMOPTYSIS**

BY

F. H. YOUNG, M.D., F.R.C.P., D.P.H.*Physician, Brompton Hospital; Medical Officer in Charge of the Tuberculosis Dispensary, St. Bartholomew's Hospital; Consulting Physician, Charing Cross Hospital*

Few conditions give rise to such alarm among patients and their friends as haemorrhage. This is accentuated when the blood is vomited or coughed up, as such bleeding is associated in the average lay mind with sudden death. The alarm is all the greater because the actual amount of blood produced seems to be much greater than it actually is, owing to its admixture with gastric juice and sputum. Again, when the bleeding is from an open wound it can be covered easily; if it comes from the lungs or stomach it cannot be concealed. When haemoptysis is relatively small in extent and occurs on a single occasion it may be difficult to distinguish from haematemesis, but when it is severe little doubt can exist.

Aetiology

The approach to a patient who is undergoing a severe haemoptysis depends on whether the practitioner knows the probable cause from his knowledge of the patient's previous history and from his clinical examination. If the cause is known it is possible to deal with the situation from an intelligent though not necessarily effective standpoint; if not there is no alternative to placebo or palliative treatment until the additional information provided by x-ray examination is available, and this should be obtained as soon as possible.

The possible causes of haemoptysis are numerous, but many of them can be omitted from immediate consideration as only extremely rarely do they cause a severe haemorrhage. In this article only those conditions are discussed which I personally have seen cause severe haemorrhage. They are: pulmonary tuberculosis; bronchiectasis; neoplasms of the lungs, bronchi, and oesophagus; pulmonary suppuration and infarcts; aneurysms of the aorta; mitral stenosis; arteriovenous aneurysm; and trauma. It will be clear at once that this list includes some conditions in which, at the worst, death from haemoptysis will only anticipate death from other causes. Nothing can be done in such cases except to use one of the methods which are described below as "unproved" and to make the patient as comfortable as possible. In these cases morphine can be given without hesitation. Severe haemorrhages from malignant neoplasms of the lower respiratory tract, or of the oesophagus extending into the bronchus, and aneurysm of the aorta, come under this category.

Haemoptysis due to the other circulatory causes—mitral stenosis, infarcts, and arteriovenous aneurysm of the lungs—is relatively common, but, though it may occur often enough to cause alarm, severe haemorrhage is rare. For practical purposes it will almost certainly stop without treatment, so all that is necessary is confidence on the part of the doctor and reassurance of the patient. Sedatives or calcium should be used if the occasion calls for them (see below), but the cough reflex must not be damped down, for if the blood is not coughed up the patient may be suffocated or an aspiration pneumonia develop. However, as the blood is not infected, the latter will usually clear quickly. The same applies to non-malignant growths of the bronchus, in which very occasionally haemorrhage is severe. In the case of non-malignant growths of the bronchus and arteriovenous aneurysm a severe haemorrhage indicates the likelihood of a recurrence, hence a resection should be performed as soon as the patient's condition makes this possible without undue risk.

Of the other causes, pulmonary tuberculosis is the most common. Here, in addition to the dangers from the loss of blood itself, there is the risk that aspiration of blood containing tubercle bacilli into distant parts of the lung will lead to new areas of disease, so making the treatment of the pulmonary tuberculosis much more difficult.

Management of the Tuberculous Case

There is a feeling that if the patient coughs it is likely to keep up the haemorrhage, but I think that the risk is exaggerated. The cough can do little harm if the patient clears the main bronchi by an occasional effective cough, but an incessant suppressed cough neither rests the lung nor prevents aspiration. Accordingly I think that it is wrong to nurse the patient flat on his back. If he has lost so much blood that syncope is likely, it is best for him to lie on his side. Otherwise I prefer him to be nursed semi-recumbent, at such an angle that coughing is easy; but if this is done the cough reflex must not be unduly depressed. If the side from which the bleeding comes is known it is best to have that side dependent so far as is practicable. The source of a severe haemoptysis can usually be determined either from the history or the patient's story (if a patient is certain from which side the bleeding comes he is nearly always correct); in addition, bubbly rales can often be heard over the site of the haemorrhage.

The patient's anxiety should be alleviated, and he can be told with truth that haemorrhage occurring for the first time in tuberculosis is practically never fatal. As an immediate measure morphine, $\frac{1}{2}$ to $\frac{1}{4}$ gr. (11 to 16 mg.) subcutaneously, can be given, which is insufficient to interfere with the cough reflex. This should be followed by sedatives; I prefer the old-fashioned chloral hydrate and potassium bromide mixture (30 gr. (2 g.) of each, which may be repeated after eight hours), though a barbiturate can be given if desired. Amylobarbitone, 3 gr. (0.2 g.), is suitable and may be repeated after six hours, but not more than half this dose should be given till the effect of the morphine has passed off, else the respiratory centre may be unduly depressed.

A large number of substances have been prescribed to stop haemoptysis, and this is the best indication of their value. No statistical proof of their value has ever been forthcoming, and they must be classified as "unproved." Indeed, most haemoptyses stop whatever is done, so proof is unlikely to be forthcoming. However, it is quite clear that the doctor must do something in the face of persistent and repeated large haemorrhages, and if he has faith in these remedies so much the better. The easiest to use, and it is at least as good as any other, is amyl nitrite capsules. These should be crushed in a handkerchief and inhaled till a feeling of faintness is produced. Two doctor patients of mine who had recurrent haemoptyses were convinced of their effect, but even the theoretical basis of their action is suspect.

Calcium is used extensively. It may conveniently be given intravenously in doses of 10 ml. of 10% calcium gluconate, repeated in twelve hours and subsequently daily. Care must be taken to avoid injecting the material into the tissue around the vein, as it is irritating. Calcium laevulinate in the same dosage, which is less irritating, can be used if the veins are difficult, as sometimes happens when the patient is shocked. It can also be given intramuscularly in case of need. These preparations do at least get into the

circulation and get there quickly. Calcium by mouth is variable and slow in absorption, and I feel that it must be a credulous doctor who will place much faith in its effect. At one time congo red in the form of a 1% solution, which must be freshly prepared, given intravenously was popular, but lately it has gone out of fashion. Undesirable side-effects occur occasionally. Serum in the form of "haemoplastin" or "coagulen" can be given in doses of 5–10 ml. intramuscularly, and following their use the haemoptysis sometimes ceases. These "unproved" remedies can be given in cases of haemoptysis from any cause.

The use of blood transfusions may be considered; if the life of the patient is endangered as a result of the haemoptysis, a small transfusion of, say, 500 ml. may be given. It is useless to do this if a severe haemoptysis is in progress, but there is little evidence that a small transfusion will restart a haemoptysis which is abating. Accordingly, if the patient's condition is serious as a result of loss of blood, and the haemoptysis has ceased or shows signs of ceasing, a small transfusion should be given.

Collapse Therapy

A serious haemoptysis of tuberculous origin comes from an unsupported vessel running across or in the wall of a tuberculous cavity, so no drug is likely to have much effect in stopping it. The rational treatment would be to deal with the vessel mechanically by resection of the area from which the bleeding occurs, but unfortunately this is impracticable while the haemorrhage is in progress. The next best thing is to try to produce relaxation of the diseased areas. Unfortunately this is more easily said than done, as the disease is usually old-standing and pleural adhesions are numerous. However, if the doctor has some experience an artificial pneumothorax should be tried. A failure will do no harm. A very small amount of relaxation may be successful in stopping the haemorrhage. Air should be introduced if practicable up to 1,500–2,000 ml., or until an average pressure around zero is reached.

An alternative—and I regard this as the most satisfactory routine measure in severe and persistent haemoptysis—is to induce a pneumoperitoneum. It is less effective than a satisfactory pneumothorax, but is much safer for the less experienced physician. There is the additional advantage that more relaxation is usually automatically obtained on the more diseased side, so that it is not necessary to know beforehand on which side the haemorrhage is occurring.

The procedure is as follows. The patient is placed on his back. The skin around a point about 2 in. (5 cm.) below the tip of the left ninth costal cartilage is cleansed. This area is better supported than other places in common use. A Morland artificial pneumothorax needle or a fine trocar and cannula, preferably not less than 2½ in. (6.3 cm.) long, is sterilized. If these are not available a wide-bore aspirating needle can be used if the angle of the bevel is reduced with a file. Preferably the area is anaesthetized with a solution of 2% procaine or similar preparation, but this is not essential. The needle is introduced through the skin and subcutaneous tissue until it is thought that it is approaching the peritoneum. The trocar is removed and the cannula or needle connected up with a pressure apparatus, preferably a water artificial pneumothorax box. The simplest is the Lillingston-Pearson apparatus here illustrated. In an emergency an improvised model of this can be made from two Kilner jars and pieces of glass and rubber tubing, sealed with insulating tape and petroleum jelly. The tap is then turned on so that air can flow in at about 10 ml. of water pressure. The needle is then pushed in very slowly till air begins to flow in freely. It is useful to remember that pushing the needle through the peritoneum causes a stab of pain which helps to indicate the position of the needle.

A free flow of air denotes that the peritoneal space has been entered. If the air flows in slowly it is probable that the point of the needle is still in the subcutaneous tissues, and the point should be advanced till it is certain that the

peritoneal cavity has been reached. If there is still no flow the needle should be withdrawn slowly, with the operator's eye kept on the pressure apparatus. A free flow will often then be seen, indicating that the needle is in the peritoneal cavity and that the point had been obstructed by omentum on its way through. When this is achieved the pressure is raised to about 25 ml. of water, and air run in until no more

Lillingston-Pearson apparatus.

will enter at this pressure. The patient should then be placed on the affected side, lying nearly flat so as to produce the maximum rise of the diaphragm. Fresh insufflations of air should be given daily for four days, when the case should be reviewed.

The risks of pneumoperitoneum are few, introduction of air into a vein in the abdominal wall being the most serious. If a hypodermic syringe is attached to the needle to make certain that it is not in a vein before the needle is connected to the source of air, the chance of this trouble is very small. Where the pneumoperitoneum is ineffective, if the patient is in hospital and a surgeon with sufficient experience is available, a temporary phrenic crush should be performed to aid the relaxation, provided it is reasonably certain from which side the haemorrhage is coming.

As soon as the haemorrhage shows signs of ceasing a single dose of streptomycin sulphate, 1 g., should be given intramuscularly daily. Isoniazid, 100 mg., twice a day by mouth should be added as soon as the patient's stomach has ceased to be irritated by the swallowed blood. When the patient is reasonably fit a full assessment of the case should be made, as a further haemoptysis is likely and may prove fatal. Surgical treatment will often be essential, and should not be delayed.

Bronchiectasis, Lung Abscess, and the Undiagnosed Case

Haemorrhage from bronchiectasis is not often as severe as from a tuberculous cavity, but a considerable amount of blood may be brought up, and the haemorrhage is apt to recur. Usually it is associated with an exacerbation of infection. Accordingly, in addition to the treatment described above, antibiotics such as chlortetracycline or chloramphenicol should be given at once instead of streptomycin and isoniazid. If the bleeding persists a pneumoperitoneum should be instituted. A phrenic interruption should not be performed except as a last resort, because if, as sometimes happens, the paralysis does not pass off the bronchiectasis may be made worse. Pulmonary suppuration apart from bronchiectasis may also give rise to a

troublesome haemoptysis. Bleeding from an acute lung abscess is seldom severe, and ceases when the abscess is treated vigorously with penicillin in large doses. Chronic lung abscess is treated as for bronchiectasis, but greater emphasis should be laid on antibiotics.

The most difficult situation arises when with the means at his disposal the doctor cannot make a diagnosis. This usually happens when no radiograph of the lungs or mediastinum is possible. What is the correct course of action? Generally speaking, the treatment of the patient is by placebos and a confident attitude. This is based on probabilities. The chances are considerable that a severe haemoptysis occurring for the first time will stop of its own accord, irrespective of treatment, if it is from a cause that will not inevitably kill the patient. The fatal haemorrhages usually occur in patients who are already doomed. It must be emphasized that the above statement refers to a first haemoptysis, and is not applicable to those who have had previous bleeding; in these the probable diagnosis can usually be arrived at and treatment carried out on intelligent lines.

Admission to Hospital

Two practical questions often arise. First, should arrangements be made for the patient's immediate removal to hospital? Personally I think that this is a mistake if reasonable facilities are available in the home and the doctor can be regularly on call, but it is usually best to initiate arrangements so that the patient can be admitted for assessment as soon as the haemorrhage has ceased. Secondly, in view of the alarm caused by the haemorrhage, the doctor should consider offering to obtain a second opinion, although, as will be seen from this article, treatment is usually within the competence of the general practitioner. How easy it is to obtain a bed in hospital depends on the area. In a number of cases the local chest physician has a few emergency beds, but I see no reason why such a patient with severe haemoptysis should not be admitted into a medical ward of a general hospital. So long as the patient is in bed there is little risk of infection to anyone except the nursing staff, and they should be properly trained in their own protection.

Next article on Emergencies in General Practice.—"Bacillary Dysentery," by Dr. R. C. MacLeod.

Refresher Course Book.—Copies of the second volume of collected articles from the Refresher Course for General Practitioners are still available at 25s. (postage 1s.) each. The first volume is now sold out.

Clinical Pathology Book.—"Clinical Pathology in General Practice," a collection of 39 articles on clinical pathology that appeared in the *Journal* as part of the Refresher Course for General Practitioners, is now available, price 21s. (postage 9d.).

Both these volumes are obtainable from the Publishing Manager, B.M.A. House, Tavistock Square, London, W.C.1, or through any bookseller.

"In recent years there has arisen in this country a multiplicity of voluntary organizations for the welfare of one group or another of the physically disabled," commented Lord KILMAINE, secretary of the Pilgrim Trust, at the annual meeting of the National Association for the Paralysed. A lead should be given towards national co-ordination. Overlapping not only wasted money on appeal funds, it also created disunity when dealing with Government departments, where "a strong, concerted approach is likely to be more successful." Further, these divisions tended to divide the sufferers into "a number of distinct categories, even social groups, whereas surely the broad aim should be to bring them all so far as possible into the ordinary life of the normal community."

British Medical Association,
Canadian Medical Association, and
Ontario Medical Association

JOINT ANNUAL MEETING, TORONTO, 1955

SUMMARY OF PROCEEDINGS

THIRD GENERAL SESSION

Why do Cancer Patients Delay?

Dr. R. F. FARQUARSON (Toronto) took the chair at the Third General Session on June 24, when three papers were read. Dr. RALSTON PATERSON (Manchester), the first to speak, gave the Gordon Richards Memorial Lecture, entitled "Why do Cancer Patients Delay?" In his introductory remarks Dr. Paterson described Dr. Gordon Richards, who had been head of the department of radiology at Toronto General Hospital, as one of the great pioneers of radiotherapy. The work he did on the treatment of cancer put not only Canada but the whole world in his debt.

Dr. Paterson then turned to the main subject of his lecture—education of the public about cancer. In attempting to find an answer to whether cancer education paid, Dr. Paterson said they must first answer the question, Was early treatment of proved value? There were important voices which made claims to the contrary. It was important to realize that in this context "early" referred to the stage of advancement of the disease in relation to each patient. Many misconceptions had arisen by equating "early" purely to time interval from the first symptom, which disregarded the different rates of growth found in any sample of tumours of significant size. Of patients with treatable cancers those with growths of limited extent, whatever the length of time of their symptoms, had a vastly higher chance of being cured. Cure could not be promised to anyone however early they came for treatment, but by coming early their chances of cure were improved.

The two reasons why patients delayed going to the doctor were ignorance and fear. The first was the dominant cause in cancers of the cervix, mouth, or skin; the latter was the main cause with breast cancer. Fear still caused greater delay than ignorance, and in Great Britain this appears to be based on a widely held belief that cancer was still totally incurable. In an investigation carried out in Manchester and its neighbourhood numbers of women were questioned, and only one-third of them thought that cancer was ever cured. In similar investigations in the United States, however, two-thirds of the women questioned thought that cancer could sometimes be cured. Any public education scheme had therefore as its purpose the elimination of the fear of cancer before going much further in the direction of education against ignorance.

In cancer education schemes the emphasis should be on the theme that "some" cancers are curable in reasonable numbers, and not on the raw slogan "Cancer is curable." In the Manchester area an attempt had been made to measure the effectiveness of an educational campaign directed primarily towards breast and cervix cancer. It was found that the measures adopted—mainly sustained pressure through a continuing series of talks at small meetings—did lead to a shortening, to a moderate extent, of the interval before a patient sought advice. The Manchester experiments had shown that a modest but definite return for the effort expended could be expected. Hence public propaganda had some value. No evidence was yet available to show whether the patients influenced by the educational schemes had had their lives saved in consequence.

Public education at its best was bound to be costly, though in the long run the good results should be cumulative. Those working in this field were brought back to the unanswerable question, "What is the price of one human life?"