The remainder were investigated by Graham and Maxwell (1954); the criteria for diagnosis have been recorded. The age distribution at the first examination is recorded in Table I.

TABLE I.—Hypertensive Age Distribution

	Males		Females	
	No.	%	No.	%
Under 49	3	0.50	13	0.84
50-54	12	2.01	53	3.42
55-59	84	14.07	253	16.32
60-64	205	34.35	478	30.84
65-69	227	38.02	521	33.61
70+	66	11.05	232	14.97
Total	597		1,550	

The control series was provided by 7,418 consecutive new registrations of blood donors at the transfusion centre. Blood groups were all determined at this centre, using the methods described by the M.R.C. (1943, 1952).

Results

The distribution of groups in our series of 2,147 cases in comparison with the control series of 7,418 cases is shown in Table II.

TABLE II.—Basic Data

Blood Group	Control	Hypertension		
		Males	Females	Total
O	3,853	309	829	1,138
A	2,485	209	513	722
B	817	72	165	237
AB	263	7	43	50
Total	7,418	597	1,550	2,147

The percentage group frequencies and the percentage increase or decrease on control, together with the sex incidence in hypertensive patients, are given in Table III.

TABLE III.—Percentage Group Frequencies

Blood Group	Control	Hypertension	% Increase or Decrease on Control	Hypertension Sex Distribution	
				% Male	% Female
O	51.94	53.04	+2.12	51.76	53.48
A	33.50	33.62	+0.36	35.01	33.10
B	11.01	11.02	-0.72	12.06	10.65
AB	3.55	2.32	-34.65	1.17	2.77

From this it can be seen that there is little difference in distribution except in the AB group. However, the difference in relative frequency of the four blood groups between the hypertensive cases and the control cases is not significant ($\chi^2=7.815$, $0.05>P>0.01$). The difference is due almost entirely to a deficiency of hypertensive cases in the AB group. A slight excess of cases in Group O, 18 greater than expectation, is not significant.

The rhesus group frequencies are recorded in Table IV. The RhD-negative percentage of a smaller group of hypertensive patients is recorded, being those who presented between 1950 and 1954. The difference in Rh incidence between those and the control series of new registrations is not significant.

TABLE IV.—Rhesus Group Frequencies

	Rhesus-positive	Rhesus-negative	Total	% Rhesus-negative
Control	1,255	263	1,518	17.3
Hypertension	1,034	213	1,247	17.1

Discussion

Hypertension and peptic ulceration are believed to have several factors in common. The mechanisms of control of the blood pressure and the gastric secretion are similar. Eppinger and Hess (1915) linked hypertension with peptic

ulceration, suggesting a common basis of autonomic imbalance, and it is possible that stress produces a mainly vasomotor, or gastric secretomotor, response. In view of the Group O disadvantage in peptic ulceration, it would not have been surprising to find an ABO relationship in essential hypertension. However, as Aird *et al.* (1954) have commented, there must be an assumption that there is no association in the absence of strong evidence. The results of this survey show no significant difference in the O, A, and B distributions in the hypertensive and control series. The slight advantage in the AB group, particularly in males, is due probably to the relatively small number involved.

Conclusion

The distribution of blood groups in 2,147 patients with essential hypertension has been examined, and no significant difference in incidence in the O, A, and B distribution has been found in comparison with the distribution in a control series of 7,418 donors. An apparent slight advantage in the AB group is regarded as due to the limited sample. It is concluded that, although an inherited factor in essential hypertension is accepted, there is no association with ABO inheritance.

We are indebted to Dr. J. Wallace, regional director of the Blood Transfusion Service, for advice and access to records, and to Dr. J. Gibson Graham, senior physician, Royal Alexandra Infirmary, Paisley. Dr. J. B. St. V. Weir, Department of Physiology, University of Glasgow, provided statistical assistance.

REFERENCES

- Aird, I., Bentall, H. H., Mehigan, J. A., and Roberts, J. A. F. (1954). *British Medical Journal*, 2, 315.
 — and Roberts, J. A. F. (1953). *Ibid.*, 1, 799.
 Allan, T. M. (1953). *Brit. J. prev. soc. Med.*, 7, 220.
 Allbutt, T. C. (1915). *Diseases of the Arteries, including Angina Pectoris*, pp. 164, 412. Macmillan, London.
 Ayman, D. (1933). *New Engl. J. Med.*, 209, 194.
 Broadbent, W. H., and Broadbent, J. F. H. (1897). *Heart Disease*, p. 80. Wood, New York.
 Dieulafoy, M. (1876). *Gaz. hebdomadaire*, 13, 594.
 Eppinger, H., and Hess, L. (1915). *Vagotonia*. Nerv. and Ment. Dis. Pub. Co., New York.
 Graham, J. G., and Maxwell, R. D. H. (1954). *British Medical Journal*, 2, 1250.
 M.R.C. (1943). War Memoranda, No. 9.
 — (1952). *Ibid.*, No. 27.
 Platt, R. (1947). *Quart. J. Med.*, 16, 111.
 Raymond P. (1907). *Progr. méd. (Paris)*, 23, 196.
 Struthers, D. (1951). *Brit. J. prev. soc. Med.*, 5, 223.
 Thomas, C. B., and Cohen, B. H. (1955). *Ann. intern. Med.*, 42, 90.
 Waterhouse, J. A. H., and Hogben, L. (1947). *Brit. J. soc. Med.*, 1, 1.
 Weitz, W. (1923). *Z. klin. Med.*, 96, 151.

RUPTURED SPLEEN REPORT OF FIVE CASES

BY

A. G. ELLERKER, M.S., F.R.C.S.
 Royal Free Hospital, London

Rupture of the spleen is a clinical problem and valuable time may be wasted in carrying out unnecessary investigations, with a corresponding increase in the mortality rate. The clinical signs depend on the degree of damage to the splenic tissue and the rate of bleeding. Ballance's sign is regarded as a classical manifestation but is by no means always present, depending as it does on the development of a large perisplenic haematoma and free intraperitoneal blood. Perhaps the most valuable sign, taken in conjunction with the other physical findings, is the presence of left shoulder pain; this may be present spontaneously; it can be elicited by raising the foot of the bed or by pressure on the phrenic nerve in the neck.

Between October, 1952, and May, 1953, five cases of rupture of the spleen were admitted to the Royal Free Hospital. Splenectomy was performed successfully in all five cases.

Traumatic Rupture

Case 1.—Male aged 17. Road accident. Concussion with retrograde amnesia. On admission, conscious and fully co-operative. Examination revealed a superficial laceration of the scalp; no other positive physical signs. In view of the retrograde amnesia he was detained for observation. Three hours after the accident he experienced a sudden severe pain in the left hypochondrium and left shoulder region. Examination showed a rigid tender upper abdomen. Laparotomy through a left upper paramedian incision. Haemoperitoneum and ruptured spleen. Splenectomy.

Case 2.—Male aged 6. On admission shocked; pulse rate 150; blood pressure 80/60 mm. Hg. He complained of severe abdominal pain, and on examination there was marked guarding down the whole of the left side of the abdomen, which did not move on respiration. No external bruising. There was a fracture of the left tibia. After resuscitation, laparotomy through a left paramedian incision. Haemoperitoneum and a severely lacerated spleen. Splenectomy.

Case 3.—Male aged 40. Road accident. On admission shocked; blood pressure 90/70 mm. Hg; pulse rate 80. Multiple fractures. Tenderness in the left hypochondrium. Blood transfusions were given, but subsequent abdominal examinations showed progressive distension, dullness in both flanks, and absence of bowel sounds. The pulse rate had risen progressively, but the blood pressure remained steady at 106/66 mm. Hg. At laparotomy haemoperitoneum and ruptured spleen. Splenectomy.

Rupture in Infectious Mononucleosis

Case 4.—Male aged 19. Fell over while running for a bus and was unconscious for about five minutes. On examination he was found to have a superficial graze over the left frontal region. Blood pressure 120/80 mm. Hg; pulse regular, good volume, rate 98 a minute. No abnormality in the central nervous system, chest, or abdomen, but there were some small discrete posterior cervical glands. In view of his period of unconsciousness he was detained for observation. Two hours after admission, while defaecating, he suddenly developed acute upper left-sided abdominal pain which spread to both shoulders. His pulse rate rose to 106, the volume was poor, and his blood pressure fell to 60/0 mm. Hg. Examination showed him to be pale and restless; his extremities were cold but he was not sweating. There was tenderness below the left costal margin with rigidity down the whole of the left side. The peritoneal release sign was positive and there was dullness in the left flank. Bowel sounds were absent. Blood transfusion was started and operation carried out.

Operation (November 8, 1952).—Left upper paramedian incision. Haemoperitoneum. Spleen enlarged with gross tears across surface, compatible with a crushing force produced by the ribs. No other abnormality apart from enlarged nodes around the coeliac axis. Splenectomy and excision of a coeliac node for biopsy.

Post-operative convalescence was satisfactory until the third day, when he developed a sore throat and a temperature of 100.2° F. (37.9° C.). On examination marked suppurative tonsillitis and pharyngitis together with enlarged tender glands in both anterior and posterior triangles of the neck, the axillae, and groins.

Subsequently the following relevant investigations were carried out:—Throat swab: β -haemolytic streptococci. Haemoglobin, 78%; W.B.C. 19,450 per c.mm. (polymorphs 29%, lymphocytes 71%). X-ray examination of the chest normal. Paul-Bunnell test positive to 1:5,120; after absorption with guinea-pig kidney this result was not significantly changed. Absorption with ox red cells reduced the titre to 1:640.

The histological reports on the excised coeliac gland and spleen were compatible with a diagnosis of infectious mononucleosis and are reported elsewhere.

The course of his infectious mononucleosis was subsequently uneventful and he was discharged to convalescence three weeks after admission.

Spontaneous Rupture Due to Carcinoma of Colon

Case 5.—Female aged 61. Had been seized with sudden severe cramp-like abdominal pain while in bed. The pain was situated in the upper abdomen and was aggravated by movement.

She had been quite well until six months before admission, but since that time had suffered from indigestion, flatulence, poor appetite, and loss of weight. Two months previously she had a gripping pain in the left groin, relieved by passing flatus, and one month later a shivering attack associated with vomiting. Since then she had had "night sweats" and a feeling of tiredness. Her bowels had been loose but regular, and on occasion she had passed a little bright red blood. For two days before admission her bowels had not been open.

On examination a well-covered woman, pale, sweating, and in pain. Blood pressure 90/70 mm. Hg. Pulse 136. Temperature 97.6° F. (36.4° C.). Respirations 32. Tongue dry and furred. Abdomen: Generalized tenderness and guarding. Moderate distension. A large, hard mass arising from the pelvis. The percussion note was dull in both flanks, and a few bowel sounds were present. On raising the foot of the bed, pain was felt in both shoulders. Rectal examination revealed a large smooth swelling in the pouch of Douglas, and on vaginal examination this was continuous with the suprapubic mass. Catheterization yielded 2 oz. (57 ml.) of clear urine.

After resuscitation laparotomy was performed through a long left upper paramedian incision. There was a haemoperitoneum with a few large clots around the spleen, which had ruptured on its convex surface. The splenic flexure of the colon was the seat of an annular constricting carcinoma firmly adherent to the spleen, tail of the pancreas, and left kidney. The uterus contained one large intramural and several smaller fibroids. There were no metastases in the liver or the regional lymphatic nodes.

After mobilization, the left half of the transverse colon, great omentum, splenic flexure, and upper half of the descending colon, together with the adherent spleen, tail of pancreas, left suprarenal gland, and left kidney were removed *en bloc*. Intestinal continuity was re-established by end-to-end anastomosis of the colon. Subtotal hysterectomy was then carried out. A corrugated drain was placed in the splenic bed and brought out through a separate stab incision in the loin. After peritoneal toilet the abdomen was closed.

Post-operatively she developed a small left subphrenic abscess which drained spontaneously through the stab incision. She was subsequently discharged, and when last seen was perfectly well with no clinical evidence of recurrence 27 months after the operation.

Microscopical examination of the organs removed showed that the colon had become adherent to the splenic veins in the region of the hilum, leading to thrombosis and septic infarcts in the spleen; rupture had presumably taken place through one of these.

Summary

Five cases of rupture of the spleen are reported. Three were traumatic in origin. Case 4 is a further example of splenic rupture occurring in a case of infectious mononucleosis. Case 5 is believed to be an unusual example of spontaneous rupture of the spleen due to a carcinoma of the splenic flexure of the colon.

Cases 1, 2, and 5 were admitted under the care of Mr. Rodney H. Maingot, Case 3 under the care of Mr. Charles H. Gray, and Case 4 under the care of Mr. George Qvist. I wish to express my sincere thanks to these surgeons for allowing me to publish these cases and, in addition, to operate on Cases 1, 4, and 5.

BIBLIOGRAPHY

- Traumatic Rupture of the Spleen*
Byrne, R. V. (1950). *Arch. Surg. (Chicago)*, **61**, 273.
- Ruptured Spleen in Infectious Mononucleosis*
Attlee, W. (1932). *Lancet*, **1**, 291.
Belton, C. H. (1952). *N.Z. med. J.*, **51**, 11.
Brien, F. S. (1947). *Canad. med. Ass. J.*, **56**, 499.
Darley, W., Black, W. C., Smith, C., and Good, F. A. (1944). *Amer. J. med. Sci.*, **208**, 381.
Davis, J. S., MacFec, W., Wright, M., and Allyn, R. (1945). *Lancet*, **2**, 72.
Fisher, J. H. (1946). *Amer. J. Path.*, **22**, 651.
Friesleben, M. (1922). *Dtsch. Z. Chir.*, **173**, 45.
Gende, B. R., and Cottrell, J. E. (1948). *Amer. Practit. (Philad.)*, **2**, 472.
Guilben, S. (1950). *J. Amer. med. Wom. Ass.*, **5**, 488.
Hicks, S. P. (1948). *Nav. med. Bull. (Wash.)*, **48**, 460.
Hill, R. M. (1953). *British Medical Journal*, **2**, 1220.
Kass, E. H., and Robbins, S. L. (1950). *Arch. Path. (Chicago)*, **50**, 644.
King, R. B. (1941). *New Engl. J. Med.*, **224**, 1058.
Milne, J. (1945). *Ibid.*, **235**, 727.
Schrager, N., and Btsh, S. (1949). *Acta med. orient. (Tel-Aviv)*, **8**, 90.
Smith, E. B., and Custer, R. P. (1946). *Blood*, **1**, 317.
Spence, W. O. (1947). *British Medical Journal*, **2**, 910.
Sullivan, J. M., and Wasserman, S. E. (1947). *J. Amer. med. Ass.*, **134**, 144.
Timmes, J. J., Averill, J. H., and Metcalfe, J. (1948). *New Engl. J. Med.*, **239**, 173.
Tudor, R. W. (1953). *British Medical Journal*, **2**, 813.
Vaughan, S. L., Regan, J. S., and Terplan, K. (1946). *Blood*, **1**, 334.
Ziegler, E. E. (1944). *Arch. Path. (Chicago)*, **37**, 196.

Medical Memoranda

Encephalitis in Infants Following Vaccination with 17 D Yellow Fever Virus: Report of a Further Case

Attention has recently been drawn to the occurrence of encephalitis as a possible complication of the vaccination of infants with 17 D yellow fever virus. Following the initial report of Smith (1954), who described its occurrence in a child of 5 weeks, two further cases were reported. Haas (1954) described the condition in an infant of 8 weeks and Scott (1954) in an infant of 13 weeks. These are the only cases which have been reported in Great Britain, although Parish (1954, personal communication) states that he is aware of one other case, in an infant of 3 months.

A further case of meningo-encephalitis in an infant following vaccination with 17 D yellow fever virus is reported.

CASE REPORT

A child of 7 weeks was vaccinated with 17 D yellow fever virus on July 24, 1953, at the Immunization Clinic, Glasgow. Seventeen days after the vaccination the child, who had been well, was noticed to be cross and feverish. His mother, who was a trained nurse, was not unduly worried and did not seek medical advice. The child remained in this state for the next two days, having no feeding or bowel disorders. He became increasingly drowsy, however, and on the twentieth day after vaccination his eyes began to roll, his limbs began to twitch, and he had a frank convulsion. His temperature, taken by his mother, was 102° F. (38.9° C.), and, according to the mother, the anterior fontanelle was bulging. The family doctor was at last called and the child was admitted to hospital.

Two days before the onset of the signs of illness the infant had been vaccinated against smallpox, but this was unsuccessful. There was no reaction either at the site of the smallpox vaccination or at the site of the vaccination against yellow fever.

On admission to hospital the infant was noted to be well developed but restless and irritable. He had a pyrexia of 100.8° F. (38.2° C.). Apart from exaggerated tendon reflexes no other abnormality was found on physical examination. Shortly after his admission he developed three further convulsive seizures.

Investigations.—The haemoglobin was 106% and the leucocyte count 13,000 per c.mm. The cerebrospinal fluid was clear but under increased pressure; the protein was 100 mg. per 100 ml., with an increase in the globulin;

the sugar was 37.2 mg. per 100 ml.; and the chlorides 738 mg. per 100 ml. The cells were 55 per c.mm. (90% lymphocytes and 10% polymorphs). No acid-fast bacilli were found and there was no growth on culture.

Two days after admission the temperature had returned to normal, but further examination of the cerebrospinal fluid two, five, and eight days after admission revealed a raised protein content, reaching a maximum of 140 mg. per 100 ml. but falling to 30 mg. per 100 ml., with an increase in globulin. The cell counts remained elevated, reaching a maximum of 88 per c.mm. but falling to 11 in the final sample; the majority of the cells were lymphocytes. There was never any growth on culture and no acid-fast bacilli were demonstrated. The Mantoux test was negative on two occasions and a radiograph of the chest revealed no abnormality.

The infant settled very quickly without antibiotic therapy and was discharged from hospital sixteen days after admission. At a follow-up examination three weeks later the infant was well and no untoward signs had been noticed in the interval.

DISCUSSION

Although, according to Theiler (1948), 5% of all who are vaccinated with 17 D yellow fever virus have a mild reaction about the seventh day, meningitic symptoms are not usually experienced. With the exception of the series described in Brazil by Fox *et al.* (1942), when a substrain of the 17 D virus with an enhanced neurotropism was incriminated, all the described cases of encephalitis following vaccination with 17 D virus have occurred in infants. In the Brazilian series the most severely affected cases were children.

Yellow-fever vaccine is prepared from chick embryos infected with the living attenuated strain of pantropic (neurotropic and viscerotropic) virus (17 D), and, according to Fox *et al.* (1942), it is still capable of producing encephalitis in mice, although after a more prolonged incubation period. It is the practice not to modify the dose of yellow-fever vaccine (0.5 ml.) with age, and infants of even the first week of life receive the same dose as adults, although Theiler (1948) states that smaller doses of vaccine can produce adequate immunity. It may be that some modification of the dose in infants is desirable.

In this country it was recommended, at a conference at the Ministry of Health in 1948, to separate vaccination against yellow fever and primary vaccination against smallpox by an interval of not less than four days when the yellow-fever inoculation is given first, and twenty-one days when the primary inoculation against smallpox has already been done first (see Parish, 1954). This precaution is not carried out in all other countries—where, for instance, the French Dakar strain of yellow-fever virus is often used simultaneously with smallpox vaccination.

Of the three previously reported cases, only the child described by Haas had been vaccinated against smallpox, and then two days prior to the onset of illness (personal communications from the authors). While it is impossible to be dogmatic in the present state of our knowledge, it is unlikely that vaccination against smallpox plays any part in the occurrence of this complication.

The incidence of the complication is not known. In Glasgow over the period January 1, 1950, to December 31, 1954, 17,040 persons of all ages were vaccinated with 17 D yellow-fever virus with no more than an occasional report of malaise or sickness. Of these 17,040, however, only 103 were children of 3 months or younger and only 174 were under the age of 7 months. If this is a complication of young infants the incidence may be underestimated if it is taken in relation to the total numbers of persons vaccinated irrespective of age. The case described is the only untoward reaction to 17 D yellow-fever vaccination that has been brought to our attention.

Further information is still required. Whether the cases described are, in fact, complications of 17 D yellow-fever-virus vaccination or are, as Roodyn (1954) suggests, perhaps