EMERGENCIES IN GENERAL PRACTICE

BACILLARY DYSENTERY

BY

R. C. MacLEOD, M.D., D.P.H., D.T.M.&H.*Department of Infectious Diseases, University of Glasgow*

The classical description of bacillary dysentery which is still enshrined in many medical textbooks depicts an acute illness usually of sudden onset in which the dominant feature is frequent and painful diarrhoea. The stools contain or may consist solely of blood and mucus. Fever and abdominal pain are characteristic and a more or less severe prostration is common, so that the reader gains the impression that the disease is a severe one. So far as Great Britain is concerned such a picture is no longer true, and, although the actual proportion of cases which merit this description may vary from place to place, and from time to time, it must now represent a small minority, and in many patients none of these classical signs is present. Nevertheless bacillary dysentery still has emergency aspects in that outbreaks may disrupt the running of institutions, the infection if undetected in food-handlers may lead to explosive outbreaks, and debilitated infants and old people are likely to be seriously ill if infected.

Aetiology and Clinical Course

The infecting agent most commonly encountered in Britain is *Shigella sonnei*, which generally produces a mild illness, so mild indeed that many individuals never seek medical advice. The epidemiological picture is further complicated by the fact that many persons excrete the bacillus without suffering any clinical illness. The remainder of the infections are produced by one or other of the Flexner types, and with these, too, the resulting illness is usually mild. The more dangerous *Shigella dysenteriae* (Shiga's bacillus) is rarely encountered here, though localized outbreaks have been recorded.

The incubation period is two to three days. In a few patients nausea, slight fever, and abdominal discomfort may be experienced for 12 hours or more before the onset of diarrhoea, and in adults the diarrhoea may gradually increase in frequency during the first two days. The onset in children is more often abrupt and may be accompanied by spasms of abdominal colic and generalized lower abdominal discomfort. Loose motions may persist for a week, but a frequency of more than two or three stools a day seldom occurs after the first 48 hours. Actual diarrhoea indeed is not invariable and, particularly in young children, the only unusual occurrence may be the passage of one or two green loose stools without any evidence of blood or mucus. Blood is, in fact, observed in the stools of only about one-third of the cases, and even in those its appearance is often on one occasion only. The usual complaints, therefore, are of a mild character, and it is unusual to find any evidence of a systemic upset. For example, in a recent survey of 200 patients admitted to hospital only four could be regarded as really ill.

The recovery of the patient is equally rapid even when specific treatment is not given; unfortunately he may continue to excrete pathogens for a long time. Complications are rarely seen. However, when dysentery is encountered among severely debilitated infants, or in the aged, a wide variety of complications have been described, but these are more an index of the primary debility than true complications of the dysenteric infection.

Differential Diagnosis

When it is known that dysentery is prevalent or an epidemic occurs in a closed community, a presumptive diag-

nosis may be made with reasonable accuracy. Nevertheless, it is undoubtedly true that unless specimens are submitted for bacteriological examination the condition will often be missed. Rectal swabs of specimens of faeces, once taken, should be transmitted to the laboratory as quickly as possible. It is worth remembering that a few doses of a sulphonamide will not eliminate the bacilli from the bowel; therefore specimens may still be sent after treatment with these drugs has begun. In the mild type of dysentery now prevalent the classical bacillary exudate is absent and no information of value is obtained by microscopical examination.

Dysentery is, of course, in effect a form of food infection. Only by bacteriological examination will it be possible in many cases to distinguish it accurately from other forms of food infection such as those caused by *Salmonellae* or *Staphylococcus aureus*. Certain characteristics, however, may give a clue. The onset in salmonellosis is usually more abrupt and occurs within twenty-four hours of consuming the infected foodstuff. Vomiting is more often noted in salmonella infections, and the purging diarrhoea and accompanying tenesmus tend to be more severe. With *Staph. aureus* food-poisoning the incubation period is still shorter (four to six hours), and usually a number of persons are attacked almost simultaneously. Vomiting is an outstanding feature and the patients may rapidly become greatly prostrated from the profuse vomiting and diarrhoea so that they seem profoundly ill. The rapidity with which the infected material is voided may be a saving feature, for it is fortunately true that recovery usually ensues with equal rapidity.

Dysentery can also occur in infancy, and here the question of *Bact. coli* gastro-enteritis must be considered. The presence of blood in the stools is uncommon in gastro-enteritis, whereas in infants with dysentery blood is often present. Further, the vomiting which is usual with gastro-enteritis can produce severe signs of dehydration very rapidly. Infection of the infant with dysentery usually arises from family contact with an older child with the disease, whereas clinical evidence of *Bact. coli* infections of older siblings is rarely seen.

The two conditions of a non-infective character which may cause difficulty are intussusception and acute appendicitis. So far as intussusception is concerned the age and sex will be important guides, but the need to carry out a rectal examination where any doubt exists must be emphasized, for this is the condition which is most commonly sent to hospital in error. An inflamed pelvic appendix may cause diarrhoea early in the illness, but the pain and tenderness usually associated with an appendicitis is not often encountered in dysentery. Further, when pain or tenderness is complained of in dysentery it is rarely localized. Here again a rectal examination will often prove of great assistance.

Epidemiology

It must be confessed that the mildness of the disease now so prevalent has not served to encourage a very detailed study of its mode of spread in the community. The only index of the prevalence of the disease is the number of notifications, but clearly these do not represent the whole picture. Housing conditions appear to play some part in determining the incidence of the disease. Thus in a series of cases admitted to hospital in Liverpool I found that two-thirds came from the poorest type of accommodation, and, although the other third came from what would be

described as better housing, overcrowding and depressed standards of living were found. Similar findings are reported from Glasgow (T. S. Wilson, personal communication). Dysentery there was most prevalent, as judged by notifications, in the worst-housed areas of the city.

It would seem correct to regard the centre of infection as the family unit rather than the individual. Admittedly the highest proportion of notified cases is in the first five years of life, but when bacteriological examination is extended to the rest of the family many of the contacts, especially those of school age, will be found to be infected. These infected persons may excrete organisms for some time, the average period in untreated persons being placed by several observers at between two and four weeks. This must mean that every person in the family is repeatedly exposed to infection. It might indeed be said that the family environment becomes saturated with the organisms.

Treatment

A few days in bed should suffice. Plenty of fluids should be provided, and the patient, if of suitable age, should always have a drink within reach. Soft foods may be allowed according to appetite, but fried foods should be avoided. The appetite usually returns to normal within a few days, but in the few cases which are more than mild premature additions to the diet, particularly of roughage, may lead to a return of abdominal discomfort and diarrhoea, and this may result in a further period of excretion of organisms.

The introduction of the sulphonamides appeared to revolutionize the treatment of bacillary dysentery, but *Sh. sonnei* infections were soon found to be more refractory than the other forms. In many areas of Britain now all or most of the strains of *Sh. sonnei* are resistant to the sulphonamides, and the practitioner is advised to obtain advice from his local laboratory from time to time about the sensitivity of the prevalent strains.

When the organism is sensitive to the sulphonamides both succinylsulphathiazole and phthalylsulphathiazole will be found effective. Sulphaguanidine is less so, particularly against *Sh. sonnei*. Phthalylsulphathiazole is more useful in treating *Sh. sonnei* carriers than the succinyl preparation. The usual adult dosage of succinylsulphathiazole is 4 g. daily for one week, and for phthalylsulphathiazole 10 g. daily for the same period. The daily amount is given in four doses. Children from 4 to 14 years should receive half, and younger children and infants one-quarter, of the adult dose. Oxytetracycline, tetracycline, and chloramphenicol have all been shown to be effective in producing bacteriological cure in bacillary dysentery. Their use, however, is not without danger. A variety of toxic effects have been described and a severe and sometimes fatal enterocolitis due to *Staph. aureus* has followed treatment with them. Therefore these antibiotics are better not used in general practice for the treatment of what is after all usually a mild and self-limiting disease with no mortality.

Oral streptomycin is very poorly absorbed and may be used with good effect, although, *in vitro* at least, *Sh. sonnei* rapidly becomes resistant to it. Combined treatment with streptomycin and a sulphonamide will often prove effective, but here again advice from the laboratory on the sensitivity of the local bacillus will prove useful. The daily dosage of streptomycin for ages up to 4 years should be 1 g., for ages 4 to 14 years 2 g., and for ages above that 4 g., all given in four doses a day: 1 g. will dissolve in 1 dr. (3.5 ml.) of water, but it is better to make it up in 4 dr. (14 ml.), so that the infant will get a teaspoonful dose (0.25 g.). The course should last for five days only.

Outbreaks in Institutions

Many practitioners are in medical charge of institutions, both residential and non-residential. When dysentery breaks out in such places, not only the inmates but all staff members in whatever capacity they may function (resident, non-resident, permanent, temporary, or part-time) must be examined bacteriologically. Further, at least two negative

examinations must be recorded before an individual is judged to be non-infected. It is as well to arrange with the matron of non-residential institutions such as day nurseries that when a child is found to have diarrhoea a rectal swab should first be taken and then the child sent home. Cases or carriers should be excluded from non-residential establishments until three negative bacteriological reports have been received, and it is wise to obtain two more after the return of the child or staff member. Residential institutions will usually transfer infected persons to an isolation hospital. When this is not possible any isolation block in being, or created for the emergency, must be organized with the strictest division of staff and facilities from those of the rest of the institution. The outbreak must be cleared up as quickly as possible to prevent the spread of infection, and therefore any drug to which the organism is sensitive may be used. It will be found, should the tetracyclines be used, that much smaller doses than those generally recommended are effective. A dosage of 5 to 10 mg. of oxytetracycline per kg. is sufficient.

Conclusion

When a practitioner is called in to see a case of bacillary dysentery he is presented not with an infected individual but with an infected family. Should it be necessary for social reasons to admit a case to hospital—and these are generally the only reasons for so doing—the problem of the infected family will remain. While little or nothing can be done at present to prevent the infection passing from case to case and from carrier to case, something can be done to prevent explosive outbreaks due to cases and carriers handling food. The practitioner can do more useful work in preventing the spread of the infection than he can in attempting to cure it. Dysentery is a notifiable disease and practitioners should be scrupulous in notifying it. Where a member of the family of a patient is employed in the processing, distributing, cooking, or serving of food the practitioner should inform the medical officer of health by telephone. This will not only enable the latter to assess without delay the necessity for action at the food establishment, but he can agree with the practitioner which contacts are to be suspended from work. The medical officer of health alone can issue a certificate to the National Insurance Office covering the absence from work of contacts of communicable disease, thus enabling such persons to draw sickness benefit.

The practitioner should give simple advice on hygiene to the family. The importance of washing the hands after defaecation should be emphasized. The housewife should have explained to her the necessity for handwashing before handling food, on keeping visitors away for a period, and on making visits herself, particularly to homes with children. She should not handle food outside her own home. It may seem that much of this advice will fall on stony ground, but a grain may take root. Lastly, the practitioner should not only treat the case but also those members of the family who have, or recently have had, loose motions.

To many of us dysentery now occupies a similar position in our thoughts to the common cold, but its widespread prevalence in the community must imply low standards of hygiene, and for this reason alone it should receive more attention. Our knowledge of its natural behaviour will be advanced only by a co-operative effort by general practitioners, public health departments, epidemiological laboratories, and hospitals.

Next article on Emergencies in General Practice.—
"Senile Confusion," by Dr. Felix Post.

Refresher Course Book.—Copies of the second volume of collected articles from the Refresher Course for General Practitioners are still available at 25s. (postage 1s.) each from the Publishing Manager, B.M.A. House, Tavistock Square, London, W.C.1, or through any bookseller. The first volume is now sold out.