EMERGENCIES IN GENERAL PRACTICE

GASSING IN THE HOME

BY

A. THELWALL JONES, M.D., M.R.C.P., D.P.H.

Consultant Physician, United Liverpool Hospitals

The recent publication of the Home Office figures on accidents in the home¹ emphasizes that all practitioners should be equipped to deal with a surprising number of emergencies.

Estimates of the number of non-fatal accidents are difficult to obtain, but analysis of the commonest kinds of fatal accidents reveals the importance of gassing as a cause. Of all the fatal accidents occurring in 1951, 58% were falls, 13% were burns and scalds, 12% were caused by suffocation, and 8% were coal-gas poisoning. Two-thirds of the deaths from coal-gas poisoning occurred in people over 65 years, and 80% of the suffocations in children under 5 years.

Although carbon monoxide is by far the commonest and most important cause of gassing in the home, an increasing number of possible agents are in use and may be classified as follows:

Asphyxiants which produce a deficiency of oxygen but do not directly injure the respiratory tract. (a) Simple inert gases which displace oxygen from the respiratory tract—for example, suffocation from fumes of fires. (b) Chemical substances which interfere with the vascular respiratory exchange—for example, carbon monoxide.

Irritants.—(a) Acting on the upper respiratory tract—for example, ammonia. (b) Acting on the upper respiratory tract and lungs—for example, chlorine, sulphur dioxide.

Centrally acting poisons which are absorbed through the lungs and produce anaesthesia—for example, trichlorethylene, carbon tetrachloride, methyl chloride, methyl bromide.

It is apparent, therefore, that when sudden and sometimes obscure illness occurs in the home the practitioner must consider the possibility of absorption of toxic gases.

Carbon Monoxide Poisoning

The importance of this poison is measured by the yearly death roll of about 500. Keith Simpson² pointed out how easily the possibility of gassing can be overlooked and has drawn attention to the preponderance of elderly people who are gassed as a result of forgetfulness, infirmity, or loss of sense of smell. Cases arise from the accidental turning on of gas-taps, insufficient ventilation or the blockage of flues, the use of common meters, from coke ovens, or from the running of car engines in garages. Incomplete combustion of butane (calor gas) in ill-ventilated rooms, ships, or caravans carries similar hazards.

Carbon monoxide, which has no colour, vapour, or odour, is absorbed through the lungs at a rate in proportion to the concentration in the atmosphere. It rapidly produces a throbbing headache and dizziness, followed by weakness of the legs and loss of consciousness. These effects are due to loss of functioning haemoglobin, which has an affinity for carbon monoxide 300 times greater than for oxygen, and is converted to carboxyhaemoglobin. The body tissues become characteristically cherry-red in colour. A sample of diluted blood from a case of carbon monoxide poisoning remains pink as compared with the normal blood similarly diluted, which has a yellow colour. The spectrum of carboxyhaemoglobin is evident if the concentration is greater than 40%. Respiration fails at least 10 to 20 minutes before the circulation ceases, and, as the combination of carbon

monoxide and haemoglobin is reversible and no other toxic effects occur, rescue and treatment should be instituted without a moment's delay.

Rescue and Treatment

If called upon to enter a gas-filled room it is essential that proper precautions should be taken:

(1) Shut off the source of gas, if possible, but do not waste time looking for taps.

(2) Open all doors and windows.

(3) Take two or three deep breaths, then hold the breath as long as possible while rescuing the patient.

(4) Crouch if coal-gas is present, as this is lighter than air, but there is no point in doing this if carbon monoxide from other sources is present, as it is about the same weight and mixes thoroughly with air.

(5) A life-line should be attached to the rescuer, who should drag the patient out.

In some circumstances it may be necessary to obtain a proper breathing apparatus from the fire brigade or police before entry is possible. A wet handkerchief across the nose only prevents irritation from smoke and does not afford any protection from carbon monoxide. The normal civil defence or service respirators are not effective against carbon monoxide.

As soon as the patient is out of the dangerous atmosphere clear his mouth, pull his tongue forward, and swab his throat if necessary. If breathing has ceased, begin artificial respiration, using the Holger Nielsen method. Keep the patient warm with blankets and hot-water bottles, and administer oxygen, as soon as available, through a face-mask as shown in the illustration. It should be presumed that shallow, difficult breathing or loss of consciousness, with or without cyanosis, following gassing accidents indicates an interference with pulmonary ventilation which can be ameliorated by oxygen. Oxygen will pass through the alveoli into the blood at twice the normal rate if the concentration is doubled in the inspired air. A flow of 3–6 litres a minute is usually adequate.

The prompt use of artificial respiration and the administration of oxygen are by far the most important principles of treatment. A number of analeptics or stimulants of the central nervous system can be given, and it is now generally agreed that the following are effective to some extent and may be employed properly in emergency: injection of

¹ Report of the Standing Interdepartmental Committee on Accidents in the Home. H.M.S.O., London. 1953.

² Simpson, K. (1954). *British Medical Journal*, 2, 774.

picrotoxin (B.P.), 3 mg. in 1 ml. intravenously; injection of leptazol (B.P.), 100 mg. in 1 ml. intravenously; or injection of nikethamide (B.P.), 4 ml. intramuscularly.

The patient may be violent during the period of recovery, but should not be allowed to walk, and should be persuaded to rest for at least a day. The headache can be relieved by rest and oxygen, but aspirin and similar drugs should not be given; morphine and barbiturates are contraindicated. Venesection should be carried out only if there is marked engorgement or pulsation of the jugular veins.

Lung Irritants

Ammonia is occasionally a source of accidental gassing from its use as a solution in enclosed spaces. *Chlorine* has been generated and caused accidents by the combination of sodium bisulphite ("harpic") and hypochlorites ("chloros") when the use of one of these materials for cleaning has immediately followed the other. Prior to 1938 refrigerators contained ammonia and sulphur dioxide, but nowadays dichloro-difluoro-methane, which is non-toxic, is generally used.

The lung irritants produce immediate irritation of the upper respiratory tract and also much discomfort in the chest and throat. In mild cases the face is congested, and coughing painful and continuous; in more severe cases there may be respiratory embarrassment with distended cervical veins and cyanosis. Bronchial spasm is a feature, and medium rhonchi and rales are heard on auscultation of the chest. Recovery is usually rapid, and, although bronchitis may ensue, serious chest complications are unusual.

The essentials of treatment are rest and oxygen. The patient should be carried to a pure warm atmosphere, the clothing loosened and removed if contaminated. Coughing can be suppressed by encouragement and reassurance and by drinks of hot sweet tea or coffee. Dyspnoea or cyanosis can be relieved by oxygen. In cases with marked irritation of the upper respiratory tract, inhalation of steam from a quart (1.1 litres) of hot water containing two teaspoonfuls of compound tincture of benzoin B.P. (friars' balsam) is of benefit. It is not desirable to give morphine or atropine, and venesection should be performed only if there is embarrassment of the right side of the heart. Leptazol or nikethamide may be given as described under carbon monoxide poisoning.

Centrally Acting Poisons

Volatile solvents such as *trichlorethylene* and *carbon tetrachloride* are used for the cleaning of clothes, and the latter is also used in fire appliances. Exposure causes intoxication, followed by drowsiness, loss of consciousness, and respiratory failure. The vapours are usually heavy, and both the rescuer and patient should be removed from the floor as quickly as possible. Trichlorethylene is decomposed at red heat and lung-irritant gases are produced; it is not safe to smoke in or near its vapour. Carbon tetrachloride may produce symptoms of an "acute abdomen" with abdominal pain and vomiting. Later, tenderness in the loins develops and the urine is scanty and contains red blood cells. The after-effects of exposure to carbon tetrachloride are potentially serious—liver and renal damage occur—and the patient should be admitted to hospital.

Methyl chloride is a colourless liquid with an ethereal and sweetish smell and has been used as a refrigerant. In addition to narcosis it can produce convulsions and oliguria. *Methyl bromide* is a colourless gas with a vapour three to four times heavier than air. It has been used in fire extinguishers, and exposure to the vapour may produce burns of the skin and respiratory passages.

The treatment of casualties from these substances is absolute rest, and it is most important that no patient should be allowed to undergo exertion or resume work for 24 hours, even though apparently only slightly affected. Artificial respiration, oxygen, and analeptics are employed as for carbon monoxide.

Next article on Emergencies in General Practice.—
"Acute Alcoholic Poisoning," by Dr. J. A. Imrie.

[A.T.P., Zurich]

CARL GUSTAVE JUNG AT 80

[FROM A SPECIAL CORRESPONDENT]

Professor C. G. JUNG, the distinguished Swiss psychiatrist, was born in Switzerland on July 26, 1875. His eightieth birthday has been honoured in many countries and he has had greetings from all over the world. In Zurich, where Jung has lived for so many years, the event was marked, in Swiss fashion, by a reception in the morning and by a dinner in the evening. This happy arrangement made it possible for all the visitors—numbering hundreds—to meet Professor Jung and his wife. Amongst the guests from Switzerland (in addition to the staff of the C. G. Jung-Institute and the Zurich Analytical Psychology Club) were Professor W. KUHN, the Rector of Basle University; Professor W. BOVEN, Rector of Lausanne University; Professor T. WYRSCH, University of Berne; Professor ROCHEDIEU, Geneva University; Professor MANFRED BLEULER, Zurich University; Dr. A. L. VISCHER, representing the Swiss Academy of Medical Sciences; Professor W. PAULI, of Zurich, and Professor TH. REICHSTEIN, of Basle, two distinguished recipients of the Nobel Prize; Dr. jur. EMIL LANDOLT, the Mayor of Zurich; Professor E. ABEGG, Professor H. HEDIGER, Professor A. KELLER, all of Zurich; also Professor QUISPTEL, Utrecht, Professor H.-CH. PUECH, Paris. Guests from the United States included Dr. VIOLET DE LASZLO and Mr. FOWLER MCCORMICK; from England came Miss RUTH BAILEY and Drs. E. A. BENNET, MICHAEL FORDHAM, F. O'D. FINIGAN, and FRANCES SMART. A *Festschrift* in two volumes, containing essays written specially by former pupils and admirers of Jung, was presented to him at the reception.

Jung comes of an old Swiss family: his grandfather, a physician of note, held the chair of medicine at Basle. He—like his grandson later—was a pioneer in the treatment of mental illness, and he set aside beds in the town hospital for psychiatric patients. This was the nucleus of the now well-known psychiatric clinic at Basle. Jung's father, a parson, in addition to other duties, was the chaplain to the town hospital, and he, too, had a special interest in the sick in mind. It is not surprising, then, that C. G. Jung, even as a medical student, paid close attention to the workings of the human mind.

The philosophy of Kant and his successors fascinated him. But it gave few clues about mental sickness. The introduction to a work on psychiatry by Krafft-Ebbing gave Jung the key he sought, and there and then he decided