EMERGENCIES IN GENERAL PRACTICE**APOPLEXY**

BY

J. ST. C. ELKINGTON, M.D., F.R.C.P.*Physician in Charge of the Neurological Department, St. Thomas's Hospital, London*

For the purposes of this article apoplexy may be defined as a malady of sudden onset characterized by depression of consciousness and evidence of severe local damage to the brain.

The clinical picture thus defined is one that is familiar to all in general practice and is probably one of the commonest medical emergencies to which general practitioners are called. Sometimes the patient will be dead on the doctor's arrival, more often death will take place within a few hours and before the patient can be moved to hospital or a second opinion obtained, should either of these steps otherwise have been deemed desirable. In other cases the victim survives the immediate effects of the ictus and the doctor will be faced with decisions concerning management, treatment, and prognosis. These are problems of considerable difficulty, and they become more so as each decade adds its quota of advances in medicine and surgery. Before they can be solved in a rational manner it is essential that an accurate diagnosis should be made, and this diagnosis, in the first place, has to be made by the only person in a position to make it—the general practitioner.

There are two distinct stages in the diagnosis of apoplexy—namely, to distinguish apoplexy from other conditions causing loss of consciousness and then to determine which of the possible causes of apoplexy is responsible for the particular case.

Recognition of Apoplexy

Apoplexy very rarely causes sudden death in the sense that coronary thrombosis or rupture of the heart does. Even though consciousness is lost instantaneously it is almost unknown even for the most disastrous cerebral vascular accident to kill in less than five or ten minutes. Apoplexy, however, is a not uncommon cause of a patient being found dead in circumstances in which the mode of dying is unknown.

The principal group of conditions from which apoplexy has to be differentiated is that in which consciousness is lost through diffuse involvement of the brain in some toxic or infective process rather than through severe local damage to its structure. Loss of consciousness, the dominating event common to both groups, removes much of the evidence on which, in a conscious patient, one would rely to make the distinction. In these circumstances the doctor has to rely on prior knowledge of the patient, upon such history as is obtainable from third parties, and upon an objective examination akin to that in veterinary medicine.

Physical signs in an unconscious patient are difficult to elicit and treacherous to interpret. Any deeply unconscious person is likely to have a congested and cyanotic appearance and to breathe stertorously. The respiration may be rhythmic, but Cheyne-Stokes breathing is particularly common in apoplexy, and when associated with blowing out of one corner of the mouth it is very suggestive of unilateral brain damage. The pupils in unconscious patients are often inactive, and may be dilated or constricted. Marked inequality in size or dissimilarity in reaction between the two pupils indicates local brain damage. In any deeply unconscious person the limbs may be flaccid, the tendon reflexes absent, and the plantar responses extensor, but even in the absence of any co-operation it may be possible to demon-

strate a marked difference between the two sides of the body, and this is a point in favour of apoplexy. Fits may occur, though they are more common when there is local brain damage, and they may leave behind them transient signs which are apt to confuse the issue. Incontinence of urine or faeces may occur in any unconscious patient and is of no diagnostic assistance. Ophthalmoscopic examination may provide critical information. Advanced papilloedema indicates raised intracranial pressure of some standing and does not make its appearance in a matter of hours, though in subarachnoid haemorrhage it may develop in a day or two. Hypertensive retinopathy is readily recognizable and should direct attention immediately to the cardiovascular system. Diabetic retinitis can be pathognomonic of that disease.

After the nervous system has been examined attention should next be directed to the cardiovascular system. A high systolic pressure—200 mm. Hg or over—is not by any means proof that the patient's illness is due to hypertension. This, and even a raised diastolic pressure, may be the result of increasing intracranial pressure, but a diastolic tension of over 120 mm. Hg is strong evidence of pre-existing hypertension. The heaving apex beat displaced to the left, which is so characteristic of left ventricular hypertrophy, is the best confirmatory evidence that hypertension is of long standing.

A specimen of urine should be examined as soon as possible. It is often necessary to pass a catheter to obtain it and time may have to be allowed if the patient has recently been incontinent. Glycosuria and even ketonuria are not by themselves proof of diabetes. The former may be found as a transient event in any acute cerebral disaster and the latter may occur after vomiting. But in these circumstances the amounts are usually small. Heavy albuminuria is strong evidence of kidney disease, but light albuminuria and fever may occur in any comatose patient.

The odour of the breath should always be noted, and may reveal the presence of ketosis, uraemia, or alcohol. The head should be examined for evidence of injury.

An assessment of these clinical observations will usually enable the physician to recognize those cases in which loss of consciousness is due to apoplexy, even in the absence of a clear history. The latter, when available, will usually put the matter beyond doubt. The reader is referred to the article in this series by Dr. M. J. Parsonage* for a consideration of the differential diagnosis of other causes of coma such as exogenous and endogenous intoxications, head injuries, and epilepsy.

Cerebral Embolism

The great majority of cases of apoplexy are caused by one of the three varieties of major cerebral vascular accident—embolism, thrombosis, or haemorrhage.

Of all varieties of apoplexy, that due to a massive cerebral embolism is the most sudden in onset. The patient may fall in his tracks as if pole-axed. The stroke is likely to occur during activity, and the damage to cerebral function, including loss of consciousness, reaches its height immediately. If the patient survives the next twenty-four to forty-eight hours, improvement is likely to set in.

A smaller embolism may produce only transient or minor symptoms, but may set up a retrograde thrombosis of the vessel proximal to the obstruction, leading to a major apoplexy of thrombotic origin when the lumen of such an artery as the middle cerebral or even the internal carotid becomes occluded.

**British Medical Journal*, 1955, 1, 217.

Apart from these features cerebral embolism can usually be suspected by the presence of a recognizable source of emboli. Of these, a diseased heart is the most important. Embolism may result from chronic mitral disease with or without auricular fibrillation; from fibrillation, however caused; from infective endocarditis; or from detachment of portions of a mural thrombus after myocardial infarction. Rarely, a fragment of clot may be detached from an atheromatous ulcer in the lumen of the aorta or the common or internal carotid arteries, and under special circumstances both fat and air may give rise to cerebral emboli.

Cerebral Thrombosis

Cerebral thrombosis is usually ingravescent in onset, occurring particularly during sleep, at rest, in the presence of debilitating circumstances such as minor illnesses, fatigue, exposure to cold, or in states of privation. It is the commonest cause of apoplexy in the aged and it is not uncommon in the middle-aged. In most cases its essential cause is atheroma of the cerebral arteries, but the occurrence of a cerebral thrombosis in a person under 40 should always bring to mind the possibility of syphilis. Rare causes of cerebral thrombosis are the different varieties of arteritis, such as polyarteritis nodosa and thrombo-angiitis obliterans, and certain of the blood diseases such as polycythaemia.

In all such cases thrombosis may be precipitated by any circumstance leading to a slowing of the cerebral circulation. Some degree of consciousness is commonly retained throughout the episode, or, if consciousness is lost, the resulting coma develops gradually. The neurological picture conforms to a recognized pattern depending upon the identity of the vessel occluded—for example, the middle cerebral, the calcarine, or the posterior inferior cerebellar arteries. Because the patient is able to co-operate, it is often possible to observe the neurological signs with accuracy. The cerebrospinal fluid in cerebral thrombosis, as in embolism, is under normal pressure and is usually normal in composition. If the resultant softening reaches the superficial or ventricular surface of the hemisphere the fluid may contain an increased amount of protein and an excess of cells, of which up to 50% may be polymorphs.

In cases of thrombosis a general medical examination will usually disclose evidence of generalized atheroma. Often, though by no means invariably, there is an associated hypertension and occasionally evidence of renal impairment. But atheroma is essentially a patchy disorder, and just as the coronary arteries may be prematurely affected in one individual, so the cerebral arteries may be prematurely affected in another. For this reason lack of evidence of generalized atherosclerosis does not exclude the diagnosis of cerebral thrombosis from this cause.

Intracranial Haemorrhage

Apart from the two varieties of traumatic meningeal haemorrhage—that is, extradural and subdural haemorrhage—which do not fall within the scope of this article, spontaneous intracranial haemorrhage may be intracerebral or subarachnoid in position. Either may occur in the absence of the other, but if the bleeding is severe a primary intracerebral haemorrhage usually leads to effusion of blood into the subarachnoid space, and a severe subarachnoid haemorrhage is often accompanied by a tearing up of the brain tissue and the consequent development of an intracerebral clot.

In the middle-aged or elderly patient intracranial haemorrhage provides the most classic example of apoplexy. A plethoric, hypertensive subject, or a person known to be suffering from chronic renal disease or other disorder associated with hypertension, is suddenly stricken while in the active pursuit of his normal affairs. Brief, agonizing headache, often associated with vomiting, quickly gives way to deepening coma with stertorous breathing and flaccidity on

one side of the body. When first seen all reflexes are commonly absent, and deepening congestion may lead to death within half an hour of the first warning of disaster. The occurrence of neck stiffness may suggest the presence of blood in the subarachnoid space. Lumbar puncture in such a case usually yields a freely blood-stained fluid, indicating that the haemorrhage has burst through the superficial or ventricular surface of the brain. When this occurs death is seldom delayed beyond a day or so.

It is often supposed that survival after an intracerebral haemorrhage is rare and that those individuals who recover are more likely, in reality, to have suffered from a thrombosis. This viewpoint has probably been overstressed, and post-mortem examination as well as the results of surgical investigation of selected cases indicate that survival after massive intracerebral haemorrhage is by no means uncommon.

In healthy persons under the age of 35 intracranial haemorrhage is the most important cause of apoplexy and is usually predominantly if not entirely subarachnoid in type. Such disasters begin with sudden intense pain in the occipital or fronto-occipital regions, often accompanied by vomiting, followed rapidly by stupor and then coma. The patient on examination shows a varying combination of drowsiness and irritability, ranging from unrousable coma on the one hand to uncontrollable excitement on the other. Signs of meningeal irritation are prominent, but those of massive damage to the brain are often conspicuous by their absence. The optic fundi often show numerous recent haemorrhages, some of them subhyaloid. The C.S.F. is freely bloodstained and classically yields, on standing, a canary-yellow supernatant fluid. Such "spontaneous" subarachnoid haemorrhage is usually easily recognized, and in a previously healthy person it is almost invariably due to the rupture of some pre-existing vascular anomaly.

Since the recognition thirty years ago of the important part played by congenital berry aneurysms of the circle of Willis in the aetiology of spontaneous subarachnoid haemorrhage, it has become usual to attribute all such haemorrhages to the rupture of one of these structures. Time has shown this view to be an over-simplification, and the increasing use of arteriography has made it clear that a considerable proportion of these cases, indeed a majority in patients under 20, result from the rupture not of a congenital aneurysm but of a so-called angiomatous malformation. These peculiar structures, long recognized as occurring chiefly in the cerebral hemispheres, are evidently far commoner than was suspected before the days of arteriography. They consist of a tangle of thin-walled blood channels fed by one or more arteries of considerable size and draining into the neighbouring dural sinuses by thin-walled vessels often as large as one's little finger. They are not angiomas in the sense of being true neoplasms, but are vascular anomalies, almost certainly congenital in origin, which continue to develop as life advances. Clinically, these cases may present with epilepsy or migraine, often with rather atypical features; with progressive damage to the brain or with sudden subarachnoid or intracerebral bleeding following rupture. Frequently, though not invariably, such structures may be associated with a bruit audible on auscultation of the skull.

The distinction between a ruptured aneurysm and a ruptured angiomatous malformation as the cause of a spontaneous subarachnoid haemorrhage in a young person is always difficult and, by clinical means, often impossible. In the majority of cases the cause can be established with certainty only by arteriography, and this important subject is referred to below.

Rarer Causes of Apoplexy

Three relatively rare causes of apoplexy still need brief consideration—namely, the apoplectiform episodes of hypertensive encephalopathy, the congestive attacks of general paresis, and cerebral tumours.

Hypertensive Encephalopathy.—Patients suffering from extreme degrees of hypertension, whether primary or secondary in origin, are liable to recurrent episodes of cerebral disorder suggestive of a major cerebral vascular catastrophe, but which clear up in the course of a few days without serious sequelae. Such incidents rarely occur with a diastolic pressure of less than 130 mm. Hg, and become more frequent as the diastolic tension mounts towards 150–160 mm. Hg. They are usually preceded for a day or two by increasing drowsiness, mental confusion, headache, and sometimes vomiting. Examination may reveal a hemiplegia, monoplegia, dysphasia, hemianopia, or even total loss of sight. A serious vascular accident is suspected, but gradually during the course of a week or ten days the symptoms subside, consciousness returns, the evidence of local brain damage recedes, and the patient's condition reverts to what it was before the incident. The precise nature of the change in the brain during these episodes is unknown, but their transitory nature suggests that oedema rather than coarse changes in the circulation is responsible.

General Paresis.—It has long been known that the clinical course of G.P.I. may be punctuated by the occurrence of acute apoplectiform incidents which have been called "congestive attacks." Loss or severe impairment of speech is a common symptom in such attacks and may or may not be associated with a paresis of the face and arm or with frank hemiplegia. Mental confusion is the rule and consciousness may be diminished or lost. Recovery is usual in the course of a few days, but the patient is commonly left with increased mental impairment after each such episode, and the tremulous dysarthria so characteristic of G.P.I. is particularly in evidence after such episodes.

Cerebral Tumours.—Finally, a word needs to be said about cerebral tumours presenting in the guise of apoplexy. Generally speaking the clinical history of a case of cerebral tumour is one of gradual evolution, but occasionally acute episodes occur which may obscure the characteristic history, or may even be the first symptom of the illness. Such incidents are particularly common with secondary deposits of carcinoma from a primary growth in the bronchus, but may occur in other secondary growths as well as in the rapidly growing gliomas of elderly people. This possibility calls for a careful review of the history in atypical cases of apoplexy, with a view to eliciting previous symptoms suggestive of a cerebral tumour.

General Management

In days gone by the treatment of apoplexy resolved itself into the general management of the case with a view to preserving life and minimizing the effects of the permanent damage to the brain resulting from the stroke. There was no curative or preventive treatment. While modern advances have not made these general measures any less necessary, they have introduced possibilities both of improvement and of prevention that require discussion.

The immediate management of a case of apoplexy is that which applies generally to any unconscious patient. Where possible the trunk should be supported on pillows and the head should be turned to one side to minimize the risk of aspiration of stomach contents and secretions. Suction may be needed to keep the airway free, and initially nasal feeding is often necessary.

Attention to the back and hips to prevent bed-sores is that appropriate to any unconscious and doubly incontinent patient. Antibiotics may play a valuable part in the prevention and relief of chest infections.

As soon as consciousness has returned and the patient is able to co-operate, all joints of the affected limbs should be put through a full range of passive movements several times a day to minimize the danger of the development of "frozen" joints. In elderly subjects this risk is greatest in respect of the hip and shoulder. The wrist and ankle should be supported when at rest in postures of physiological dorsiflexion, but splints should be avoided, as they promote fixation. As soon as voluntary movement is regained in any

part it should be vigorously encouraged by both nurses and relatives as well as by the skilled physiotherapist by means of active movements under supervision and of occupational therapy.* The patient should be got up into a chair as soon as possible after it is clear that natural recovery has begun, and attempts at walking should be started within a day or two of the patient being out of bed. Electrotherapy has no useful part to play in the treatment of such cases, and it is doubtful if any form of actinotherapy has more than a psychobiological effect.

In addition to these general measures, which are applicable to all recovering cases of apoplexy, modern therapeutic advances have introduced at least three possibilities of more active treatment which require consideration in appropriate cases. These are: (a) the treatment of the underlying hypertension in cases where hypertensive disease is the cause of the stroke, (b) the use of anticoagulants in cases of cerebral thrombosis and embolism, and (c) surgery in cases of subarachnoid and intracerebral haemorrhage.

Treatment of Arterial Hypertension

In many cases apoplexy is no more than an event in the evolution of the underlying disorder of essential or symptomatic hypertension, and in cases that survive the remote prognosis is that of the underlying disease.

The treatment of hypertension is at present in a phase of active evolution. In the past ten years thoraco-lumbar sympathectomy, hypotensive drugs, especially the methonium compounds, salt-free diets, and total adrenalectomy have all claimed attention. In a disorder so liable to natural fluctuations in intensity and so long drawn out in its evolution as essential hypertension, it is not possible yet to arrive at any final decision on the best treatment. However, in cases where cerebral vascular accidents have already occurred a decision for or against treatment has often to be taken without delay, and for this reason certain provisional, if purely personal, conclusions appear to be permissible. In severe cases of hypertension, whether essential or symptomatic, any measure which effectively reduces the blood pressure, as for example nephrectomy in unilateral pyelonephritis, is likely to be followed by a gratifying improvement in the general cerebral symptoms of hypertension and by a reduction of the liability to further cerebral damage. In the commoner cases where at best treatment can hope only to achieve an improvement in the blood-pressure level without reducing it to normal limits, the cerebral symptoms are among those which undergo the greatest degree of amelioration, and I have seen nothing to make me think that the risks of thrombosis with a lower pressure exceed the dangers of haemorrhage with a higher one. I therefore conclude that in a given case of essential hypertension the occurrence or threat of a cerebral vascular accident should be regarded as an encouragement rather than as a contraindication to active treatment of the underlying arterial hypertension.

Anticoagulants in Cerebral Thrombosis and Embolism

The introduction of anticoagulants suitable for prolonged administration and their proved value in the results of coronary artery disease naturally raises the question of their possible usefulness in combating the results of cerebral vascular disease.

Theoretically, it is possible to envisage a number of advantages and disadvantages of such treatment. Anticoagulant therapy should decrease the liability to further thrombosis of the cerebral vessels in threatened persons, as in cases of coronary occlusion, but in the case of the brain the risk of haemorrhage exists which does not enter the picture in cases of myocardial infarction. Further, the danger of an extension or recurrence after a cerebral thrombosis is much less than after a coronary occlusion, and the differential diagnosis between a cerebral thrombosis and an intracerebral haemorrhage is often in doubt. Possibly from these

*For further details see under "Any Questions?" at p. 1220, the answer on physiotherapy after a stroke.

considerations, possibly from native conservatism, anticoagulant therapy does not seem to have been extensively employed in this country.

Surgery in Cerebral Haemorrhage

The introduction of cerebral angiography has made possible a precision in the diagnosis of certain cerebral vascular accidents which could never have been achieved by purely clinical means, and with this improvement in diagnosis has come the possibility of direct attack on the source and results of intracranial haemorrhage.

Angiography was first applied to the investigation of cases of spontaneous subarachnoid haemorrhage. Although it has long been possible to diagnose such cases with reasonable certainty, and although it was recognized that they were due in many cases to the rupture of a pre-existing anomaly, it was seldom possible to determine with accuracy the position of the offending aneurysm or angioma, and in consequence treatment could only be supportive. Now it is possible in many cases to visualize small aneurysms on the component vessels of the circle of Willis or on the middle or anterior cerebral arteries, and in suitable cases these may be eliminated surgically or the dangers of a further haemorrhage may at least be reduced by ligation of the appropriate carotid artery.

One of the surprising results of angiography as applied to a large number of cases of subarachnoid haemorrhage has been to show how many of these cases, particularly in adolescents and young adults, result from bleeding from angiomatous malformations rather than from aneurysms. These vascular anomalies usually lie on the surface of the brain and are eminently suitable for surgical removal. When once they are removed, the liability to recurrent bleeding is slight, as, unlike aneurysms, they are solitary. One thus reaches the conclusion that every case of subarachnoid haemorrhage in a young person without hypertension should be submitted to full investigation, including cerebral arteriography, as soon as possible after the initial haemorrhage. It is well to remember that there is often only a short interval between the initial bleeding and a subsequent one that may be fatal.

Nor is it only in cases of subarachnoid bleeding that modern methods of neurosurgical and neuroradiological investigation have introduced the possibility of more active treatment. The same is true of intracerebral haemorrhage. Regular study of post-mortem material makes it clear that many patients survive intracerebral haemorrhages of moderate size, provided these do not rupture into the ventricle or through the surface of the brain. Such haemorrhages give rise to intracerebral clots which separate the cortex from the underlying white matter or split the white matter apart without necessarily destroying its fibres.

It has been contended by various workers that the early evacuation of such clots permits a degree of recovery of function much greater than would occur if the clots were left to resolve by the natural processes of liquefaction and absorption. With this view I am in agreement. The practical application of this viewpoint clearly requires that we should be able to recognize cases of intracerebral haematoma and particularly to distinguish them from ischaemic softenings; on purely clinical grounds this is always difficult and often impossible. The associated circumstances, the mode of onset of the stroke, and the shape of the cerebral lesion may all give important indications, but are seldom sufficient to decide with certainty between a softening and an intracerebral haemorrhage in an elderly hypertensive subject. On the other hand, shift of the pineal shadow, deviation of the midline arteries towards the hemiplegic side, and distortion of the ventricular system may together give indisputable evidence of an intracerebral haemorrhage under tension and so justify an operation to evacuate the clot.

Such investigations can only be carried out in hospitals specially staffed and equipped, and a moment's reflection will make it clear that the acceptance of the viewpoint

expressed here will create a practical problem of considerable magnitude. However, it is one from which we should not shrink if careful and controlled study confirms that the degree of recovery in cases of intracerebral haemorrhage is thereby improved.

Conclusion

The main purpose of this article has been to advocate a more critical and constructive attitude towards the subject of apoplexy. This condition is one of the chief causes of death and disabling illness after middle age, and year by year its toll increases.

It must be feared that for the foreseeable future this will continue to be the case, but modern methods provide us with means of dealing effectively at least with the fringe of the problem. The first requirement, if such methods are to be employed, is that cases suitable for investigation and possibly for specialized treatment should be separated from those that are unsuitable.

This demands accurate diagnosis, which, in the nature of the problem, must initially be made in the patient's own home by those in general practice.

Next article on Emergencies in General Practice.—"Acute Psychosis," by Drs. J. Harrington and W. Mayer-Gross.

Refresher Course Book.—Copies of the second volume of collected articles from the Refresher Course for General Practitioners are still available at 25s. (postage 1s.) each. The first volume is now sold out.

Clinical Pathology Book.—"Clinical Pathology in General Practice," a collection of 39 articles on clinical pathology that appeared in the *Journal* as part of the Refresher Course for General Practitioners, is now available, price 21s. (postage 9d.).

Both these volumes are obtainable from the Publishing Manager, B.M.A. House, Tavistock Square, London, W.C.1, or through any bookseller.

VISIT OF THE RUSSIAN DOCTORS

The six distinguished Russian doctors who have been visiting Britain as guests of the British Medical Association concluded their three-week tour on Tuesday, November 8, when they left by plane for Russia.

London

The tour opened with the luncheon at B.M.A. House on Monday, October 17, at which Dr. E. A. GREGG, chairman of the B.M.A. Council, welcomed the visitors (see this *Journal*, October 22, p. 1021). At a reception at the Royal Society of Medicine that evening the president, Sir WILLIAM GILLIATT, received them. Most of the next day was spent at the Postgraduate Medical School at Hammersmith. Professors JOHN MCMICHAEL and IAN AIRD were guides, and Sir DESMOND MORTON, chairman of the board of governors, presided at luncheon. In the evening London University gave a dinner in honour of the visitors, the vice-chancellor, then Professor HAROLD ROBINSON, presiding. On Wednesday (October 19) the guests were received at the Wellcome Foundation's museums and laboratories in the Euston Road by Mr. M. W. PERRIN, chairman of the Foundation, and Sir HENRY DALE, chairman of the Wellcome Trustees. At the luncheon given by the Society of Apothecaries at their Hall in Black Friar's Lane, the master, Dr. N. S. FINZI, was host. The party then divided, some going to see the Glaxo Laboratories at Greenford, some to the surgical unit at St. Mary's Hospital, and some to the Chester Beatty Research Institute at the Royal Marsden Hospital. The National Institute for Medical Research at Mill Hill was visited on Thursday (October 20), where the director, Sir CHARLES