corresponding sigmoidoscopic improvement and the mucosa reverted to normal.

Some possible uses of this method of study are discussed.

REFERENCES

- Boston, F. K., and Macintosh, R. R. (1939). *Brit. dent. J.*, 66, 575.
 Gabriel, W. B., Dukes, C. E., and Bussey, H. J. R. (1951). *Brit. J. Surg.*, 38, 401.
 Levine, M. D., Kirsner, J. B., and Klotz, A. P. (1951). *Science*, 114, 552.
 Saher, N. F., and Salt, R. (1943). *British Medical Journal*, 1, 790.
 Truelove, S. C., and Witts, L. J. (1954). *Ibid.*, 2, 375.
 Warren, S., and Sommers, S. C. (1949). *Amer. J. Path.*, 25, 657.
 Wood, I. J., Doig, R. K., Motteram, R., and Hughes, A. (1949). *Lancet*, 1, 18.

GIANT-CELL ARTERITIS WITH ANEURYSMS

EFFECTS OF HORMONE THERAPY

BY

R. J. HARRISON, M.R.C.P.

Physician, St. James' Hospital, London

C. V. HARRISON, M.D.

*Professor of Morbid Anatomy, University of London,
Postgraduate Medical School of London*

AND

H. KOPELMAN, M.D., M.R.C.P.

Physician, St. Margaret's Hospital, Epping

[WITH SPECIAL PLATE]

Giant-cell or temporal arteritis is now a well-recognized entity with which most physicians and pathologists are familiar. The morphological changes were surveyed by one of us a few years ago (Harrison, 1948), and the clinical picture has recently been reviewed by Roux (1954), who traced over 200 cases in the literature. Our reasons for recording another two cases are: (1) to draw attention to some of the less common manifestations of the disease; and (2) to describe the effects of hormone therapy on the histology of the arterial lesions, our findings in this respect being different from those of previous workers.

Case 1

A 56-year-old housewife was well until December, 1953, when she complained of lassitude, gradual loss of weight, and night sweats. She also had a non-productive cough, for which her practitioner prescribed a linctus and phenobarbitone. In July, 1954, she noticed a painless swelling in the right side of her neck, and a week later a similar swelling appeared in the right infraclavicular region.

At this time there was a pulsating aneurysmal dilatation of the right external carotid artery $1\frac{1}{2}$ in. (3.75 cm.) long and 1 in. (2.5 cm.) across. Similar aneurysmal swellings were also present in the left and right infraclavicular regions and in the right axilla. The fingers of the right hand showed early clubbing, and this hand was warmer than the left. She was afebrile, and weighed 9 st. 12 lb. (62.6 kg.). There was no skin rash or muscle tenderness and no arthritis. Her blood pressure was 130/90 mm. Hg and her pulse 84. No other abnormality was found on clinical examination.

Blood culture was sterile on three occasions. Examination of the blood showed: Hb 75% (Haldane); W.B.C. 11,000 per c.mm. (differential count normal); albumin 3.2 g. and globulin 3.7 g. per 100 ml. (A./G. ratio 0.9); blood urea 19 mg. per 100 ml.; Wassermann reaction negative. Urine: no albumin; centrifuged deposit—no R.B.C. E.S.R. 80 mm. in one hour (Westergren).

Giant-cell arteritis was suspected, and 15 mm. of the affected left suprascapular artery was removed for examination. The specimen contained one small segment of artery

that was healthy, the rest being enlarged to twice the normal diameter. Microscopically the lumen in the thickened part of the artery was reduced to a slit by massive thickening of the intima by loose mucoid connective tissue infiltrated with inflammatory cells (Special Plate, Fig. 1). The media was virtually destroyed for about half its circumference, the elastica being broken into fragments. There was a heavy cellular infiltration consisting of polymorphonuclear leucocytes, macrophages, and lymphocytes. Giant cells were seen in relation to fragmented elastic tissue. The picture was typical of giant-cell arteritis in an active stage.

As soon as the diagnosis was confirmed oral administration of cortisone was started, 200 mg. being given daily during the first week. Thereafter the daily dosage was 150 mg. during the second, 100 mg. during the third, 50 mg. during the fourth, and 25 mg. during the fifth week. After the first week the erythrocyte sedimentation rate (E.S.R.) fell to normal and remained so; during and after the course of treatment the patient experienced a general feeling of well-being. No significant change in her weight or blood pressure occurred during treatment.

There was still some doubt whether the inflammatory processes were still active or not, and a second biopsy was performed five weeks after the first, the right acromio-thoracic axis being exposed by a transverse incision below the clavicle. The axis itself was dilated; a large fusiform aneurysm of the axillary artery at least 4 cm. in diameter was defined and palpated. The acromio-thoracic axis with its four branches was removed for section. On microscopical examination the main artery showed signs of healed arteritis (Plate, Fig. 2), but its branches were normal. The intima was thickened by loose mucoid connective tissue exactly like that of the earlier biopsy. For about half the circumference the elastica was fragmented and the media thinned (owing to loss of most of its muscle) and vascularized. In spite of these changes there was no inflammatory cellular infiltration.

For fear of recrudescence an oral maintenance dose of 25 mg. cortisone daily was continued. By December the aneurysmal swelling of the right external carotid artery had disappeared. In early January there was severe pain in the right arm with increased swelling in the right axillary region. This persisted, and after several days the right hand and fingers were numb and cold. On January 17 there was sudden onset of right wrist-drop with loss of sensation over the upper limb and especially marked in the right hand. The swelling in the axilla was no longer pulsatile and was very firm; no bruit was heard over it. The E.S.R. at this time was 11 mm. in one hour (Westergren). It was considered that there was a large thrombosed aneurysm in the axilla, and this was confirmed by arteriography. An operation to relieve nerve pressure was performed on January 25, cortisone treatment being discontinued.

At operation (Mr. Andrew M. Desmond) the main mass of the aneurysm was found to be false and occupying the lower part of the axilla. The smaller upper part of the aneurysm was a fusiform dilatation apparently of the subclavian artery just beyond the acromio-thoracic axis. There was no demonstrable communication between the two parts of the aneurysm. The false aneurysm was opened, the clot evacuated, and the bleeding controlled. A piece of the end of the aneurysm and the adjoining vessel was taken for histological examination.

Sections from the normal part of the vessel disclosed that it was a vein and not an artery as supposed. At this level the intima showed some thickening and there were a few minute foci of cellular infiltration in the intima and media; in other respects the part was healthy. A longitudinal section through the end of the aneurysm (Plate, Fig. 3) showed thinning-out of the vein wall without (in the tissue taken) any sign of medial destruction. An organizing thrombus was attached to one wall close to a venous valve; there was no sign of inflammatory activity. Fragments from the wall of the false aneurysm showed only fibrous tissue with no vascular residue.

Since the operation the patient has had physiotherapy daily, and there has been a fair degree of recovery of function of the right arm and hand. On March 30 the E.S.R. was 5 mm. in one hour.

The cortisone used for this case was supplied by the Medical Research Council for clinical trial.

Case 2

A retired policeman aged 76 was admitted on March 16, 1953, with severe throbbing headaches in the temporal regions extending to the top of the cranium, the angles of the jaw, and the neck. Although there had been slight improvement, the pain was severe and not relieved by the usual analgesics. Both temporal arteries appeared to be greatly thickened (Fig. A) and were extremely tender to

Case 2. Showing right temporal artery (A) before treatment, and (B) after treatment and immediately before second biopsy was taken.

palpation. Apart from this he was very fit for his age; his blood pressure was 170/90 mm. Hg, the fundi were normal, and there was no leucocytosis. His E.S.R. was 36 mm. in one hour.

On March 23 a biopsy and ligation of the left temporal artery was performed under local analgesia. The next day pain in the left temporal artery was much less and there was slight improvement on the opposite side, but by the 25th the pain had returned in the ligated artery proximal to the ligation. The distal part of the artery was, however, much less tender to palpation. The opposite artery remained unchanged. On the 27th A.C.T.H. gel was given by injection, 25 mg. three times a day. Within 24 hours the pain was completely relieved, and after 48 hours the artery became much less thickened, pulsatile, and no longer tender (Fig. B).

After eleven days' treatment a biopsy taken from the second artery showed the process to be inactive, and it was felt safe to reduce the dose of A.C.T.H. slowly over a period of seven days and finally to omit it. The patient has remained well and symptom-free since.

At the first biopsy the histology was typical of giant-cell arteritis (Plate, Fig. 4). The lumen of the artery was a minute slit. The inner half of the intima was composed of active fibroblastic connective tissue, and the outer half was invaded by vasa infiltrated by polymorphonuclear leucocytes and macrophages with a few eosinophil granulocytes and plasma cells. The internal elastic lamina was fragmented and there were giant cells among the fragments. The media was infiltrated by cells like those in the intima. The adventitia was thickened and infiltrated by a few cells, mostly plasma cells.

The histological appearances of the specimen taken after treatment were totally different (Plate, Fig. 5). The artery was almost exactly half the diameter of the earlier one. As before, the lumen was reduced to a slit and there was the same degree of intimal thickening, but there was now no cellular infiltration. The internal elastic lamina was frag-

mented but no longer the site of a giant-cell reaction. The media similarly had completely lost its cellular infiltration, though in places it was still traversed by enlarged vasa. Finally, the adventitia was of about normal thickness, and the only sign of previous involvement was the presence of a few groups of lymphocytes.

Discussion

The first case showed a number of unusual features: although for some months the patient had had lassitude, night sweats, and loss of weight, she did not suffer from headaches or from pain in any of the affected regions. The occurrence of saccular aneurysms is even more unusual. We have been able to trace only two other cases: the first was Gilmour's (1941) Case I, in which the patient died of a ruptured subclavian aneurysm, and the other the remarkable case of Gelfand (1955) of an aortic and a subclavian aneurysm in an African boy of 2 years. In addition to these two cases, Magarey (1950) described a fatal dissecting aneurysm of the aorta, and McMillan (1950) a partial aortic tear without dissection. Aneurysms occur rarely in a disease which causes such severe medial damage as giant-cell arteritis; when they do they appear to suggest a possible explanation for this. The stress in the wall of an artery is related not only to the pressure within, but also to the size of the lumen. In small arteries the lumen is rapidly reduced by the intimal overgrowth that is such a characteristic feature of this disease. In larger arteries like the subclavian, on the other hand, even a considerable degree of intimal thickening will not reduce the lumen sufficiently to protect a damaged media.

The use of either cortisone or A.C.T.H. is now well established in the treatment of this disease (Roux, 1954). Usually there has been prompt and dramatic relief of symptoms and slower improvement of physical signs, but the effect is suppressive and the disease is apt to flame up if treatment is not continued long enough. Occasionally treatment fails, and Fabian (1952) records a case confirmed by biopsy in which there was no clinical improvement. The effect of hormone treatment on the structural changes in the vessels is less certain, because only a few cases have been subjected to biopsy before and after treatment. We have found reports of four such cases. Siegert (1952) gave an injection of depot cortisone and took a second biopsy after 12 days. Höök and Jernelius (1952) gave 1 g. of cortisone over a period of a month. Oöstuizen and Van Wyk (1954) 20 mg. of A.C.T.H. four-hourly for 12 days, and Roux (1954) 13 g. of cortisone over 12 days, before taking a second biopsy. In all four cases the authors report clinical improvement, but in no case was there any significant change in histological appearance.

In both our own cases there was striking regression of the inflammatory reaction, and in fact we believe that they are the first cases reported in which serial biopsy has shown such structural change. We are unable to offer any satisfactory explanation. It might be argued that in the second biopsies the arteries had not been affected or had undergone spontaneous healing before treatment. The former argument seems to be disproved by the clinical change and the severe scarring that was present, and the second by the obvious clinical change that had taken place in an easily palpable artery. The dose of hormone used might be thought to explain the difference; our first case certainly received a much greater dose of cortisone than other published cases. In our second case, however, the dose of A.C.T.H. was less than that given, for example, by Oöstuizen and Van Wyk.

The clinical and histological evidence that we have is that cortisone and A.C.T.H. reduce the oedema and the inflammatory cellular reactions. This is probably the cause of the relief of symptoms. There is no evidence that the treatment has any effect on the scarring and intimal thickening that has already taken place; in fact in the first case thrombosis occurred despite treatment. It seems probable

R. J. HARRISON *ET AL.*: GIANT-CELL ARTERITIS WITH ANEURYSMS

FIG. 1.—Case 1: Left suprascapular artery before treatment. The intima is greatly thickened, leaving a slit-like lumen. The media is destroyed in the lower two-thirds and both coats are heavily infiltrated with inflammatory cells. (Haematoxylin and eosin. $\times 30$.)

FIG. 2.—Case 1: Acromio-thoracic axis after treatment. Intimal thickening unchanged. Media interrupted by fibrous scars, but both coats free from cellular infiltration. (The black crescent is operative haemorrhage in the vessel wall.) (Haematoxylin and eosin. $\times 30$.)

FIG. 3.—Case 1: Subclavian vein showing aneurysm. A mass of thrombus is present to the left of the valve. (Haematoxylin and eosin. $\times 19$.)

FIG. 4.—Case 2: Left temporal artery before treatment, showing intimal thickening and heavy cellular infiltration including giant cells. (Haematoxylin and eosin. $\times 73$.)

FIG. 5.—Case 2: Right temporal artery after treatment. The intimal thickening is unaltered, but the artery has returned to normal size and is totally free from cellular infiltration. Fragmentation of elastic tissue not visible in this section. (Haematoxylin and eosin. $\times 73$.)

that it is thrombus which causes the arterial obstruction and symptoms after cortisone has been given, and it might be argued that there is an indication for the use of anti-coagulants during and following cortisone therapy.

Summary

Two cases of giant-cell arteritis—one with the formation of saccular aneurysms—in a woman of 56 and a man of 76 are presented. Treatment was respectively with cortisone orally and A.C.T.H. by injection. Both patients recovered. The histological findings both before and after treatment are described and illustrated. A possible explanation of why aneurysms occur only in the larger vessels in giant-cell arteritis is offered. The advantages of a second biopsy after completion of treatment, with a view to assessing the degree of structural recovery, are pointed out.

REFERENCES

- Fabian, G. (1952). *Klin. Mbl. Augenheilk.*, 121, 80.
 Gelfand, M. (1955). *Brit. Heart J.*, 17, 264.
 Gilmour, J. R. (1941). *J. Path. Bact.*, 53, 263.
 Harrison, C. V. (1948). *J. clin. Path.*, 1, 197.
 Hook, O., and Jernelius, B. (1952). *Nord. Med.*, 48, 1224.
 McMillan, G. C. (1950). *Arch. Path. (Chicago)*, 49, 63.
 Magarey, F. R. (1950). *J. Path. Bact.*, 62, 445.
 Oosthuizen, W. J., and Van Wyk, F. A. K. (1954). *S. Afr. med. J.*, 28, 310.
 Roux, J. L. (1954). *Helv. med. Acta*, 21, Suppl. 34.
 Sievert, P. (1952). *Klin. Mbl. Augenheilk.*, 120, 254.

GIANT-CELL ARTERITIS OF THE LEGS

BY

R. FINLAYSON, M.B., B.Chir.

AND

J. O. ROBINSON, M.Chir., F.R.C.S.

From the Departments of Pathology and Surgery,
 St. Bartholomew's Hospital, London

[WITH SPECIAL PLATE]

Giant-cell or temporal arteritis was described clinically by Hutchinson in 1890, but it was not until 1934 that Horton *et al.* recognized the histological features. The condition was initially believed to be localized to the temporal arteries, but of recent years it has become apparent that it may be a generalized arterial disease. Visual disturbances and cerebral symptoms are frequent complications, but the development of gangrene in the limbs has rarely been mentioned, and it is for this reason that the following case is thought to be worth recording.

Case Report

A woman aged 75 was admitted to St. Bartholomew's Hospital on July 14, 1954, complaining of pain in both feet. She had suffered with cold feet and chilblains for many years, but these had caused her no serious inconvenience until three months before admission, when she developed a sudden severe pain in both legs, the left leg being the more painful. The legs became white, and were swollen to a level above her knees. A diagnosis of "thrombosis" was made, and after twelve days' confinement to bed the pain and swelling subsided and the colour returned to normal. When she became ambulant she noticed that her feet were colder than before, and that they were blue and stiff. The feet and legs became increasingly painful, but she persisted with her daily housework until two weeks before admission. Following the "thrombosis" she noted that her general health had deteriorated; she had felt a

little feverish towards the evening, her appetite had gradually worsened, and tiredness had become a prominent feature.

She was a pale woman with bilateral cataracts. The fundi showed choroidal pigmentation, the veins were normal, but the arteries were sclerosed. There was no evidence of temporal arteritis. Her heart was clinically not enlarged. Her arms were normal. B.P. 140/70; T. 99° F. (37.2° C.); P. 80; R. 22.

Legs.—*Left*: The foot was swollen and the skin waxen in appearance. The distal half of the foot was cold and dark blue, with patches of white on the toes. The remainder of the foot and limb was warm and of a normal colour. *Right*: The distal third of the foot was cold and dark blue apart from the toes, which were white; the remainder of the foot and limb was warm and of a normal colour; the femoral and popliteal pulses were present in both legs, but pulsation could not be felt in either foot.

Investigations.—Hb, 10.4 g. per 100 ml.; W.B.C., 16,000 per c.mm.; E.S.R., 25 mm. in one hour (Westergren); serum proteins, 6.7 g. per 100 ml. (albumin 3.9 g., globulin 2.8 g.); blood urea, 34 mg. per 100 ml. Urine: No albumin or sugar detected; a centrifuged deposit showed an occasional white blood cell, a few epithelial cells, and red blood corpuscles (one to two in an occasional 1/6 field). Wassermann and Kahn reactions were negative.

X-ray Examination.—Chest film: slight ventricular hypertrophy; lung fields clear. Intravenous pyelogram: both kidneys functioned promptly, with a good concentration. Leg films: there was evidence of a minor degree of vascular calcification.

Operation.—A below-knee amputation of the left leg was performed with a right lumbar sympathectomy on July 16.

Pathological Examination of Left Leg

Except for peripheral gangrene, the only pathological changes noted on dissection of the limb were in the larger arteries, which resembled firm cords. Atheromatous plaques were not evident on macroscopical examination, but the vessel walls showed multiple minute foci of calcification. The accompanying venae comitantes were small, but the vessels were fully patent except for a little recent blood clot in a small lateral plantar vein. The long and short saphenous veins were normal.

Microscopical examination of the anterior tibial, posterior tibial, and peroneal arteries showed essentially similar histological changes. The lumen was either completely obliterated or reduced to a narrow chink by thickening of the intima, resulting from a proliferation of loose, fairly vascular mesenchymal tissue infiltrated with small numbers of chronic inflammatory cells, predominantly lymphocytic in type (Special Plate, Fig. 1). Small foci of atheromatous fatty change were also present in some sections. The internal elastic lamina of the vessels was reduplicated and stained irregularly; the inner fibres, in general, stained differently, while the outer fibres appeared highly refractile, and in many areas were closely related to fine or heavy deposits of calcium. The elastica showed some fragmentation, but complete loss of a segment of the internal elastic lamina was seen in two sections only, and at both sites the deficiency was anatomically related to a calcified plaque in the adjacent inner media (Plate, Fig. 2).

There was an intense diffuse cellular infiltration of the media composed mainly of lymphocytes, but also including moderate numbers of histiocytes together with varying numbers of multinucleated giant cells, many with peripherally arranged nuclei, and showing, in some fields, a focal arrangement simulating tuberculous follicles (Plate, Fig. 3). Small to moderate numbers of plasma cells were also identified (Plate, Fig. 4), but neutrophil and eosinophil leucocytes either were absent or were present in very small numbers. The media showed increased vascularity, fibro-